

Hyperholomorphic sheaves and new examples of hyperkähler manifolds

Misha Verbitsky,

`verbit@thelema.dnttm.rssi.ru`, `verbit@ihes.fr`

Given a compact hyperkähler manifold M and a holomorphic bundle B over M , we consider a Hermitian connection ∇ on B which is compatible with all complex structures on M induced by the hyperkähler structure. Such a connection is unique, because it is Yang-Mills. We call the bundles admitting such connections hyperholomorphic bundles. A stable bundle is hyperholomorphic if and only if its Chern classes c_1, c_2 are $SU(2)$ -invariant, with respect to the natural $SU(2)$ -action on the cohomology. For several years, it was known that the moduli space of stable hyperholomorphic bundles is singular hyperkähler. More recently, it was proven that singular hyperkähler varieties admit a canonical hyperkähler desingularization. In the present paper, we show that a moduli space of stable hyperholomorphic bundles is compact, given some assumptions on Chern classes of B and hyperkähler geometry of M (we also require $\dim_{\mathbb{C}} M > 2$). Conjecturally, this leads to new examples of hyperkähler manifolds. We develop the theory of hyperholomorphic sheaves, which are (intuitively speaking) coherent sheaves compatible with hyperkähler structure. We show that hyperholomorphic sheaves with isolated singularities can be canonically desingularized by a blow-up. This theory is used to study degenerations of hyperholomorphic bundles.

Contents

1	Introduction	3
1.1	An overview	3
1.1.1	Examples of hyperkähler manifolds	3
1.1.2	Hyperholomorphic bundles	4
1.1.3	Hyperholomorphic sheaves	6
1.1.4	Deformations of hyperholomorphic sheaves	7
1.2	C -restricted complex structures: an introduction	9
1.3	Quaternionic-Kähler manifolds and Swann's formalism	10

1.4	Contents	13
2	Hyperkähler manifolds	15
2.1	Hyperkähler manifolds	15
2.2	Simple hyperkähler manifolds	17
2.3	Trianalytic subvarieties in hyperkähler manifolds.	18
2.4	Hyperholomorphic bundles	20
2.5	Stable bundles and Yang–Mills connections.	22
2.6	Twistor spaces	24
3	Hyperholomorphic sheaves	25
3.1	Stable sheaves and Yang–Mills connections	25
3.2	Stable and semistable sheaves over hyperkähler manifolds . .	27
3.3	Hyperholomorphic connection in torsion-free sheaves	29
3.4	Existence of hyperholomorphic connections	31
3.5	Tensor category of hyperholomorphic sheaves	35
4	Cohomology of hyperkähler manifolds	37
4.1	Algebraic induced complex structures	37
4.2	The action of $\mathfrak{so}(5)$ on the cohomology of a hyperkähler manifold	37
4.3	Structure of the cohomology ring	39
4.4	Cohomology classes of CA-type	43
5	C-restricted complex structures on hyperkähler manifolds	44
5.1	Existence of C -restricted complex structures	44
5.2	Hyperkähler structures admitting C -restricted complex structures	52
5.3	Deformations of coherent sheaves over manifolds with C -restricted complex structures	55
6	Desingularization of hyperholomorphic sheaves	56
6.1	Twistor lines and complexification	57
6.2	The automorphism $\Psi_{I,J}$ acting on hyperholomorphic sheaves	59
6.3	A $\mathbb{C}^*$ -action on a local ring of a hyperkähler manifold	62
6.4	Desingularization of $\mathbb{C}^*$ -equivariant sheaves	65
7	Twistor transform and quaternionic-Kähler geometry	66
7.1	Direct and inverse twistor transform	66
7.2	Twistor transform and Hermitian structures on vector bundles	68
7.3	B_2 -bundles on quaternionic-Kähler manifolds	70

7.4	Hyperkähler manifolds with special $\mathbb{H}^*$ -action and quaternionic-Kähler manifolds of positive scalar curvature	74
8	$\mathbb{C}^*$-equivariant twistor spaces	78
8.1	B_2 -bundles on quaternionic-Kähler manifolds and $\mathbb{C}^*$ -equivariant holomorphic bundles over twistor spaces	78
8.2	$\mathbb{C}^*$ -equivariant bundles and twistor transform	81
8.3	Twistor transform and the $\mathbb{H}^*$ -action	84
8.4	Hyperholomorphic sheaves and $\mathbb{C}^*$ -equivariant bundles over M_{fl}	86
8.5	Hyperholomorphic sheaves and stable bundles on $\mathbb{C}P^{2n+1}$. .	89
9	Moduli spaces of hyperholomorphic sheaves and bundles	90
9.1	Deformation of hyperholomorphic sheaves with isolated singularities	90
9.2	The Maruyama moduli space of coherent sheaves	95
9.3	Moduli of hyperholomorphic sheaves and C -restricted complexes structures	97
10	New examples of hyperkähler manifolds	99
10.1	Twistor paths	99
10.2	New examples of hyperkähler manifolds	104
10.3	How to check that we obtained new examples of hyperkähler manifolds?	105

1 Introduction

For an introduction to basic results and the history of hyperkähler geometry, see [Bes].

This Introduction is independent from the rest of this paper.

1.1 An overview

1.1.1 Examples of hyperkähler manifolds

A Riemannian manifold M is called **hyperkähler** if the tangent bundle of M is equipped with an action of quaternion algebra, and its metric is Kähler with respect to the complex structures I_ι , for all embeddings $\mathbb{C} \xhookrightarrow{\iota} \mathbb{H}$. The complex structures I_ι are called **induced complex structures**; the corresponding Kähler manifold is denoted by (M, I_ι) .

For a more formal definition of a hyperkähler manifold, see Definition 2.1. The notion of a hyperkähler manifold was introduced by E. Calabi ([C]).

Clearly, the real dimension of M is divisible by 4. For $\dim_{\mathbb{R}} M = 4$, there are only two classes of compact hyperkähler manifolds: compact tori and K3 surfaces.

Let M be a complex surface and $M^{(n)}$ be its n -th symmetric power, $M^{(n)} = M^n/S_n$. The variety $M^{(n)}$ admits a natural desingularization $M^{[n]}$, called **the Hilbert scheme of points**.

The manifold $M^{[n]}$ admits a hyperkähler metrics whenever the surface M is compact and hyperkähler ([Bea]). This way, Beauville constructed two series of examples of hyperkähler manifolds, associated with a torus (so-called “higher Kummer variety”) and a K3 surface. It was conjectured that all compact hyperkähler manifolds M with $H^1(M) = 0$, $H^{2,0}(M) = \mathbb{C}$ are deformationally equivalent to one of these examples. In this paper, we study the deformations of coherent sheaves over higher-dimensional hyperkähler manifolds in order to construct counterexamples to this conjecture. A different approach to the construction of new examples of hyperkähler manifolds is found in the recent paper of K. O’Grady, who studies the moduli of semistable bundles over a K3 surface and resolves the singularities using methods of symplectic geometry ([O’G]).

1.1.2 Hyperholomorphic bundles

Let M be a compact hyperkähler manifold, and I an induced complex structure. It is well known that the differential forms and cohomology of M are equipped with a natural $SU(2)$ -action (Lemma 2.5). In [V1], we studied the holomorphic vector bundles F on (M, I) which are compatible with a hyperkähler structure, in the sense that any of the following conditions hold:

- (i) The bundle F admits a Hermitian connection ∇ with a curvature $\Theta \in \Lambda^2(M, \text{End}(F))$ which is of Hodge type $(1,1)$ with respect to any of induced complex structures.
- (ii) The bundle F is a direct sum of stable bundles, and its Chern classes $c_1(F)$, $c_2(F)$ are $SU(2)$ -invariant. (1.1)

These conditions are equivalent (Theorem 2.27). Moreover, the connection ∇ of (1.1) (i) is Yang-Mills (Proposition 2.25), and by Uhlenbeck–Yau theorem (Theorem 2.24), it is unique.

A holomorphic vector bundle satisfying any of the conditions of (1.1) is called **hyperholomorphic** ([V1]).

Clearly, a stable deformation of a hyperholomorphic bundle is again a hyperholomorphic bundle. In [V1], we proved that a deformation space of hyperholomorphic bundles is a singular hyperkähler variety. A recent development in the theory of singular hyperkähler varieties ([V-d], [V-d2], [V-d3]) gave a way to desingularize singular hyperkähler manifolds, in a canonical way. It was proven (Theorem 2.16) that a normalization of a singular hyperkähler variety (taken with respect to any induced complex structure I) is a smooth hyperkähler manifold.

This suggested a possibility of constructing new examples of compact hyperkähler manifolds, obtained as deformations of hyperholomorphic bundles. Two problems arise.

Problem 1. The deformation space of hyperholomorphic bundles is *a priori* non-compact and must be compactified.

Problem 2. The geometry of deformation spaces is notoriously hard to study. Even the dimension of a deformation space is difficult to compute, in simplest examples. How to find, for example, the dimension of the deformation space of a tangent bundle, on a Hilbert scheme of points on a K3 surface? The Betti numbers are even more difficult to compute. Therefore, there is no easy way to distinguish a deformation space of hyperholomorphic bundles from already known examples of hyperkähler manifolds.

In this paper, we address Problem 1. Problem 2 can be solved by studying the algebraic geometry of moduli spaces. It turns out that, for a generic deformation of a complex structure, the Hilbert scheme of points on a K3 surface has no closed complex subvarieties ([V5]; see also Theorem 2.17). It is possible to find a 21-dimensional family of deformations of the moduli space $\text{Def}(B)$ of hyperholomorphic bundles, with all fibers having complex subvarieties (Lemma 10.28). Using this observation, it is possible to show that $\text{Def}(B)$ is a new example of a hyperkähler manifold. Details of this approach are given in Subsection 10.3, and the complete proofs will be given in a forthcoming paper.

It was proven that a Hilbert scheme of a generic K3 surface has no tri-analytic subvarieties.¹ Given a hyperkähler manifold M and an appropriate hyperholomorphic bundle B , denote the deformation space of hyperholomorphic connections on B by $\text{Def}(B)$. Then the moduli of complex structures

¹Trianalytic subvariety (Definition 2.9) is a closed subset which is complex analytic with respect to any of induced complex structures.

on M are locally embedded to a moduli of complex structures on $\text{Def}(B)$ (Claim 10.26). Since the dimension of the moduli of complex structures on $\text{Def}(B)$ is equal to its second Betti number minus 2 (Theorem 5.9), the second Betti number of $\text{Def}(B)$ is no less than the second Betti number of M . The Betti numbers of Beauville's examples of simple hyperkähler manifolds are 23 (Hilbert scheme of points on a K3 surface) and 7 (generalized Kummer variety). Therefore, for M a generic deformation of a Hilbert scheme of points on K3, $\text{Def}(B)$ is either a new manifold or a generic deformation of a Hilbert scheme of points on K3. It is easy to construct trianalytic subvarieties of the varieties $\text{Def}(B)$, for hyperholomorphic B (see [V2], Appendix for details). This was the motivation of our work on trianalytic subvarieties of the Hilbert scheme of points on a K3 surface ([V5]).

For a generic complex structure on a hyperkähler manifold, all stable bundles are hyperholomorphic ([V2]). Nevertheless, hyperholomorphic bundles over higher-dimensional hyperkähler manifolds are in short supply. In fact, the only example to work with is the tangent bundle and its tensor powers, and their Chern classes are not prime. Therefore, there is no way to insure that a deformation of a stable bundle will remain stable (like it happens, for instance, in the case of deformations of stable bundles of rank 2 with odd first Chern class over a K3 surface). Even worse, a new kind of singularities may appear which never appears for 2-dimensional base manifolds: a deformation of a stable bundle can have a singular reflexization. We study the singularities of stable coherent sheaves over hyperkähler manifolds, using Yang-Mills theory for reflexive sheaves developed by S. Bando and Y.-T. Siu ([BS]).

1.1.3 Hyperholomorphic sheaves

A compactification of the moduli of hyperholomorphic bundles is the main purpose of this paper. We require the compactification to be singular hyperkähler. A natural approach to this problem requires one to study the coherent sheaves which are compatible with a hyperkähler structure, in the same sense as hyperholomorphic bundles are holomorphic bundles compatible with a hyperkähler structure. Such sheaves are called **hyperholomorphic sheaves** (Definition 3.11). Our approach to the theory of hyperholomorphic sheaves uses the notion of admissible Yang-Mills connection on a coherent sheaf ([BS]).

The equivalence of conditions (1.1) (i) and (1.1) (ii) is based on Uhlenbeck–Yau theorem (Theorem 2.24), which states that every stable bundle F with $\deg c_1(F) = 0$ admits a unique Yang–Mills connection, that is, a

connection ∇ satisfying $\Lambda\nabla^2 = 0$ (see Subsection 2.5 for details). S. Bando and Y.-T. Siu developed a similar approach to the Yang–Mills theory on (possibly singular) coherent sheaves. Consider a coherent sheaf F and a Hermitian metric h on a locally trivial part of $F|_U$. Then h is called admissible (Definition 3.5) if the curvature ∇^2 of the Hermitian connection on $F|_U$ is square-integrable, and the section $\Lambda\nabla^2 \in \text{End}(F|_U)$ is uniformly bounded. The admissible metric is called **Yang–Mills** if $\Lambda\nabla^2 = 0$ (see Definition 3.6 for details). There exists an analogue of Uhlenbeck–Yau theorem for coherent sheaves (Theorem 3.8): a stable sheaf admits a unique admissible Yang–Mills metric, and conversely, a sheaf admitting a Yang–Mills metric is a direct sum of stable sheaves with the first Chern class of zero degree.

A coherent sheaf F is called **reflexive** if it is isomorphic to its second dual sheaf F^{**} . The sheaf F^{**} is always reflexive, and it is called a **reflexization** of F (Definition 3.1).

Applying the arguments of Bando and Siu to a reflexive coherent sheaf F over a hyperkähler manifold (M, I) , we show that the following conditions are equivalent (Theorem 3.19).

- (i) The sheaf F is stable and its Chern classes $c_1(F)$, $c_2(F)$ are $SU(2)$ -invariant
- (ii) F admits an admissible Yang–Mills connection, and its curvature is of type $(1,1)$ with respect to all induced complex structures.

A reflexive sheaf satisfying any of these conditions is called **reflexive stable hyperholomorphic**. An arbitrary torsion-free coherent sheaf is called **stable hyperholomorphic** if its reflexization is hyperholomorphic, and its second Chern class is $SU(2)$ -invariant, and **semistable hyperholomorphic** if it is a successive extension of stable hyperholomorphic sheaves (see Definition 3.11 for details).

This paper is dedicated to the study of hyperholomorphic sheaves.

1.1.4 Deformations of hyperholomorphic sheaves

By Proposition 2.14, for an induced complex structure I of general type, **all** coherent sheaves are hyperholomorphic. However, the complex structures of general type are never algebraic, and in complex analytic situation, the moduli of coherent sheaves are, generally speaking, non-compact. We study the flat deformations of hyperholomorphic sheaves over (M, I) , where I is an algebraic complex structure.

A priori, a flat deformation of a hyperholomorphic sheaf will be no longer hyperholomorphic. We show that for some algebraic complex structures, called **C -restricted complex structures**, a flat deformation of a hyperholomorphic sheaf remains hyperholomorphic (Theorem 5.14). This argument is quite convoluted, and takes two sections (Sections 4 and 5).

Further on, we study the local structure of stable reflexive hyperholomorphic sheaves with isolated singularities. We prove the Desingularization Theorem for such hyperholomorphic sheaves (Theorem 6.1). It turns out that such a sheaf can be desingularized by a single blow-up. The proof of this result is parallel to the proof of Desingularization Theorem for singular hyperkähler varieties (Theorem 2.16).

The main idea of the desingularization of singular hyperkähler varieties ([V-d2]) is the following. Given a point x on a singular hyperkähler variety M and an induced complex structure I , the complex variety (M, I) admits a local $\mathbb{C}^*$ -action which preserves x and acts as a dilatation on the Zariski tangent space of x . Here we show that any stable hyperholomorphic sheaf F is equivariant with respect to this $\mathbb{C}^*$ -action (Theorem 6.6, Definition 6.11). Then an elementary algebro-geometric argument (Proposition 6.12) implies that F is desingularized by a blow-up.

Using the desingularization of hyperholomorphic sheaves, we prove that a hyperholomorphic deformation of a hyperholomorphic bundle is again a bundle (Theorem 9.3), assuming that it has isolated singularities. The proof of this result is conceptual but quite difficult, it takes 3 sections (Sections 7–9), and uses arguments of quaternionic-Kähler geometry ([Sw], [N2]) and twistor transform ([KV]).

In our study of deformations of hyperholomorphic sheaves, we usually assume that a deformation of a hyperholomorphic sheaf over (M, I) is again hyperholomorphic, i. e. that an induced complex structure I is C -restricted, for C sufficiently big (Definition 5.1). Since C -restrictness is a tricky condition, it is preferable to get rid of it. For this purpose, we use the theory of twistor paths, developed in [V3-bis], to show that the moduli spaces of hyperholomorphic sheaves are real analytic equivalent for different complex structures I on M (Theorem 10.14). This is done as follows.

A hyperkähler structure on M admits a 2-dimensional sphere of induced complex structures. This gives a rational curve in the moduli space $Comp$ of complex structures on M , so-called **twistor curve**. A sequence of such rational curves connect any two points of $Comp$ (Theorem 10.4). A sequence of connected twistor curves is called a **twistor path**. If the intersection points of these curves are generic, the twistor path is called **admissible** (Definition 10.6). It is known (Theorem 10.8) that an admissible twistor

path induces a real analytic isomorphism of the moduli spaces of hyperholomorphic bundles. There exist admissible twistor paths connecting any two complex structures (Claim 10.13). Thus, if we prove that the moduli of deformations of hyperholomorphic bundles are compact for one generic hyperkähler structure, we prove a similar result for all generic hyperkähler structures (Theorem 10.14). Applying this argument to the moduli of deformations of a tangent bundle, we obtain the following theorem.

Theorem 1.1: Let M be a Hilbert scheme of points on a K3 surface, $\dim_{\mathbb{H}}(M) > 1$ and $\mathcal{H}$ a generic hyperkähler structure on M . Assume that for all induced complex structures I , except at most a finite many of, all semistable bundle deformations of the tangent bundle $T(M, I)$ are stable. Then, for all complex structures J on M and all generic polarizations ω on (M, J) , the deformation space $\mathcal{M}_{J,\omega}(T(M, J))$ is singular hyperkähler and compact, and admits a smooth compact hyperkähler desingularization.

Proof: This is Theorem 10.20. ■

In the course of this paper, we develop the theory of C -restricted complex structures (Sections 4 and 5) and another theory, which we called **the Swann's formalism for vector bundles** (Sections 7 and 8). These themes are of independent interest. We give a separate introduction to C -restricted complex structures (Subsection 1.2) and Swann's formalism (Subsection 1.3).

1.2 C -restricted complex structures: an introduction

This part of the Introduction is highly non-precise. Our purpose is to clarify the intuitive meaning of C -restricted complex structure.

Consider a compact hyperkähler manifold M , which is **simple** (Definition 2.7), that is, satisfies $H^1(M) = 0$, $H^{2,0}(M) = \mathbb{C}$.

A **reflexive hyperholomorphic sheaf** is by definition a semistable sheaf which has a filtration of stable sheaves with $SU(2)$ -invariant c_1 and c_2 . A **hyperholomorphic sheaf** is a torsion-free sheaf which has hyperholomorphic reflexization and has $SU(2)$ -invariant c_2 (Definition 3.11). If the complex structure I is of general type, all coherent sheaves are hyperholomorphic (Definition 2.13, Proposition 2.14), because all integer (p, p) -classes are $SU(2)$ -invariant. However, for generic complex structures I , the corresponding complex manifold (M, I) is never algebraic. If we wish to compactify the moduli of holomorphic bundles, we need to consider algebraic complex structures, and if we want to stay in hyperholomorphic cat-

egory, the complex structures must be generic. This paradox is reconciled by considering the C -restricted complex structures (Definition 5.1).

Given a generic hyperkähler structure $\mathcal{H}$, consider an algebraic complex structure I with $\text{Pic}(M, I) = \mathbb{Z}$. The group of rational (p, p) -cycles has form

$$\begin{aligned} H_I^{p,p}(M, \mathbb{Q}) = & H^{2p}(M, \mathbb{Q})^{SU(2)} \oplus a \cdot H^{2p}(M, \mathbb{Q})^{SU(2)} \\ & a^2 \cdot \oplus H^{2p}(M, \mathbb{Q})^{SU(2)} \oplus \dots \end{aligned} \quad (1.2)$$

where a is a generator of $\text{Pic}(M, I) \subset H_I^{p,p}(M, \mathbb{Z})$ and $H^{2p}(M, \mathbb{Q})^{SU(2)}$ is the group of rational $SU(2)$ -invariant cycles. This decomposition follows from an explicit description of the algebra of cohomology given by Theorem 4.6. Let

$$\Pi : H_I^{p,p}(M, \mathbb{Q}) \longrightarrow a \cdot H^{2p}(M, \mathbb{Q})^{SU(2)} \oplus a^2 \cdot H^{2p}(M, \mathbb{Q})^{SU(2)} \oplus \dots$$

be the projection onto non- $SU(2)$ -invariant part. Using Wirtinger's equality, we prove that a fundamental class $[X]$ of a complex subvariety $X \subset (M, I)$ is $SU(2)$ -invariant unless $\deg \Pi([X]) \neq 0$ (Proposition 2.11). A similar result holds for the second Chern class of a stable bundle (Corollary 3.24).

A C -restricted complex structure is, heuristically, a structure for which the decomposition (1.2) holds, and $\deg a > C$. For a C -restricted complex structure I , and a fundamental class $[X]$ of a complex subvariety $X \subset (M, I)$ of complex codimension 2, we have $\deg[X] > C$ or X is trianalytic. A version of Wirtinger's inequality for vector bundles (Corollary 3.24) implies that a stable vector bundle B over (M, I) is hyperholomorphic, unless $|\deg c_2(B)| > C$. Therefore, over a C -restricted (M, I) , all torsion-free semistable coherent sheaves with bounded degree of the second Chern class are hyperholomorphic (Theorem 5.14).

The utility of C -restricted induced complex structures is that they are algebraic, but behave like generic induced complex structures with respect to the sheaves F with low $|\deg c_2(F)|$ and $|\deg c_1(F)|$.

We prove that a generic hyperkähler structure admits C -restricted induced complex structures for all C , and the set of C -restricted induced complex structures is dense in the set of all induced complex structures (Theorem 5.13). We prove this by studying the algebro-geometric properties of the moduli of hyperkähler structures on a given hyperkähler manifold (Subsection 5.2).

1.3 Quaternionic-Kähler manifolds and Swann's formalism

Quaternionic-Kähler manifolds (Subsection 7.3) are a beautiful subject of Riemannian geometry. We are interested in these manifolds because they

are intimately connected with singularities of hyperholomorphic sheaves. A stable hyperholomorphic sheaf is equipped with a natural connection, which is called **hyperholomorphic connection**. By definition, a hyperholomorphic connection on a torsion-free coherent sheaf is a connection ∇ defined outside of singularities of F , with square-integrable curvature ∇^2 which is an $SU(2)$ -invariant 2-form (Definition 3.15). We have shown that a stable hyperholomorphic sheaf admits a hyperholomorphic connection, and conversely, a reflexive sheaf admitting a hyperholomorphic connection is a direct sum of stable hyperholomorphic sheaves (Theorem 3.19).

Consider a reflexive sheaf F over (M, I) with an isolated singularity in $x \in M$. Let ∇ be a hyperholomorphic connection on F . We prove that F can be desingularized by a blow-up of its singular set. In other words, for $\pi : \widetilde{M} \longrightarrow (M, I)$ a blow-up of $x \in M$, the pull-back π^*F is a bundle over $\widetilde{M}$.

Consider the restriction $\pi^*F|_C$ of π^*F to the blow-up divisor

$$C = \mathbb{P}T_x M \cong \mathbb{C}P^{2n-1}.$$

To be able to deal with the singularities of F effectively, we need to prove that the bundle $\pi^*F|_C$ is semistable and satisfies $c_1(\pi^*F|_C) = 0$.

The following intuitive picture motivated our work with bundles over quaternionic-Kähler manifolds. The manifold $C = \mathbb{P}T_x M$ has a quaternionic structure, which comes from the $SU(2)$ -action on $T_x M$. We know that bundles which are compatible with a hyperkähler structure (hyperholomorphic bundles) are (semi-)stable. If we were able to prove that the bundle $\pi^*F|_C$ is in some way compatible with quaternionic structure on C , we could hope to prove that it is (semi-)stable.

To give a precise formulation of these heuristic arguments, we need to work with the theory of quaternionic-Kähler manifolds, developed by Berard-Bergery and Salamon ([Sal]). A quaternionic-Kähler manifold (Definition 7.8) is a Riemannian manifold Q equipped with a bundle W of algebras acting on its tangent bundle, and satisfying the following conditions. The fibers of W are (non-canonically) isomorphic to the quaternion algebra, the map $W \hookrightarrow \text{End}(TQ)$ is compatible with the Levi-Civita connection, and the unit quaternions $h \in W$ act as orthogonal automorphisms on TQ . For each quaternion-Kähler manifold Q , one has a twistor space $\text{Tw}(Q)$ (Definition 7.10), which is a total space of a spherical fibration consisting of all $h \in W$ satisfying $h^2 = -1$. The twistor space is a complex manifold ([Sal]), and it is Kähler unless W is flat, in which case Q is hyperkähler. Further on,

we shall use the term “quaternionic-Kähler” for manifolds with non-trivial W .

Consider the twistor space $\text{Tw}(M)$ of a hyperkähler manifold M , equipped with a natural map

$$\sigma : \text{Tw}(M) \longrightarrow M.$$

Let (B, ∇) be a bundle over M equipped with a hyperholomorphic connection. A pullback $(\sigma^*B, \sigma^*\nabla)$ is a holomorphic bundle on $\text{Tw}(M)$ (Lemma 7.2), that is, the operator $\sigma^*\nabla^{0,1}$ is a holomorphic structure operator on σ^*B . This correspondence is called **the direct twistor transform**. It is invertible: from a holomorphic bundle $(\sigma^*B, \sigma^*\nabla^{0,1})$ on $\text{Tw}(M)$ it is possible to reconstruct (B, ∇) , which is unique ([KV]; see also Theorem 7.3).

A similar construction exists on quaternionic-Kähler manifolds, due to T. Nitta ([N1], [N2]). A bundle (B, ∇) on a quaternionic-Kähler manifold Q is called a **B_2 -bundle** if its curvature ∇^2 is invariant with respect to the adjoint action of $\mathbb{H}^*$ on $\Lambda^2(M, \text{End}(B))$ (Definition 7.12). An analogue of direct and inverse transform exists for B_2 -bundles (Theorem 7.14). Most importantly, T. Nitta proved that on a quaternionic-Kähler manifold of positive scalar curvature a twistor transform of a B_2 -bundle is a Yang-Mills bundle on $\text{Tw}(Q)$ (Theorem 7.17). This implies that a twistor transform of a Hermitian B_2 -bundle is a direct sum of stable bundles with $\deg c_1 = 0$.

In the situation described in the beginning of this Subsection, we have a manifold $C = \mathbb{P}T_x M \cong \mathbb{C}P^{2n-1}$ which is a twistor space of a quaternionic projective space

$$\mathbb{P}_{\mathbb{H}}T_x M = \left(T_x M \setminus \{0\} \right) / \mathbb{H}^* \cong \mathbb{H}P^n.$$

To prove that $\pi^*F|_C$ is stable, we need to show that $\pi^*F|_C$ is obtained as twistor transform of some Hermitian B_2 -bundle on $\mathbb{P}_{\mathbb{H}}T_x M$.

This is done using an equivalence between the category of quaternionic-Kähler manifolds of positive scalar curvature and the category of hyperkähler manifolds equipped with a special type of $\mathbb{H}^*$ -action, constructed by A. Swann ([Sw]). Given a quaternionic-Kähler manifold Q , we consider a principal bundle $\mathcal{U}(Q)$ consisting of all quaternion frames on Q (7.4). Then $\mathcal{U}(Q)$ is fibered over Q with a fiber $\mathbb{H}/\{\pm 1\}$. It is easy to show that $\mathcal{U}(Q)$ is equipped with an action of quaternion algebra in its tangent bundle. A. Swann proved that if Q has with positive scalar curvature, then this action of quaternion algebra comes from a hyperkähler structure on $\mathcal{U}(M)$ (Theorem 7.24).

The correspondence $Q \longrightarrow \mathcal{U}(Q)$ induces an equivalence of appropriately defined categories (Theorem 7.25). We call this construction **Swann's formalism**.

The twistor space $\mathrm{Tw}(\mathcal{U}(Q))$ of the hyperkähler manifold $\mathcal{U}(Q)$ is equipped with a holomorphic action of $\mathbb{C}^*$. Every B_2 -bundle corresponds to a $\mathbb{C}^*$ -invariant holomorphic bundle on $\mathrm{Tw}(\mathcal{U}(Q))$ and this correspondence induces an equivalence of appropriately defined categories, called **Swann's formalism for bundles** (Theorem 8.5). Applying this equivalence to the $\mathbb{C}^*$ -equivariant sheaf obtained as an associate graded sheaf of a hyperholomorphic sheaf, we obtain a B_2 bundle on $\mathbb{P}_{\mathbb{H}}T_x M$, and $\pi^*F|_C$ is obtained from this B_2 -bundle by a twistor transform.

The correspondence between B_2 -bundles on Q and $\mathbb{C}^*$ -invariant holomorphic bundles on $\mathrm{Tw}(\mathcal{U}(Q))$ is an interesting geometric phenomenon which is of independent interest. We construct it by reduction to $\dim Q = 0$, where it follows from an explicit calculation involving 2-forms over a flat manifold of real dimension 4.

1.4 Contents

The paper is organized as follows.

- Section 1 is an introduction. It is independent from the rest of this paper.
- Section 2 is an introduction to the theory of hyperkähler manifolds. We give a compendium of results from hyperkaehler geometry which are due to F. Bogomolov ([Bo]) and A. Beauville ([Bea]), and give an introduction to the results of [V1], [V-d3], [V2(II)].
- Section 3 contains a definition and basic properties of hyperholomorphic sheaves. We prove that a stable hyperholomorphic sheaf admits a hyperholomorphic connection, and conversely, a reflexive sheaf admitting a hyperholomorphic connection is stable hyperholomorphic (Theorem 3.19). This equivalence is constructed using Bando-Siu theory of Yang–Mills connections on coherent sheaves.

We prove an analogue of Wirtinger's inequality for stable sheaves (Corollary 3.24), which states that for any induced complex structure $J \neq \pm I$, and any stable reflexive sheaf F on (M, I) , we have

$$\deg_I \left(2c_2(F) - \frac{r-1}{r}c_1(F)^2 \right) \geq \left| \deg_J \left(2c_2(F) - \frac{r-1}{r}c_1(F)^2 \right) \right|,$$

and the equality holds if and only if F is hyperholomorphic.

- Section 4 contains the preliminary material used for the study of C -restricted complex structures in Section 5. We give an exposition of various algebraic structures on the cohomology of a hyperkähler manifold, which were discovered in [V0] and [V3]. In the last Subsection, we apply the Wirtinger's inequality to prove that the fundamental classes of complex subvarieties and c_2 of stable reflexive sheaves satisfy a certain set of axioms. Cohomology classes satisfying these axioms are called CA-classes. This definition simplifies the work on C -restricted complex structures in Section 5.
- In Section 5 we define C -restricted complex structures and prove the following. Consider a compact hyperkähler manifold and an $SU(2)$ -invariant class $a \in H^4(M)$. Then for all C -restricted complex structures I , with $C > \deg a$, and all semistable sheaves F on (M, I) with $c_2(F) = a$, the sheaf F is hyperholomorphic (Theorem 5.14). This is used to show that a deformation of a hyperholomorphic sheaf is again hyperholomorphic, over (M, I) with I a C -restricted complex structure, $c > \deg c_2(F)$.

We define the moduli space of hyperkähler structures, and show that for a dense set $\mathcal{C}$ of hyperkähler structures, all $\mathcal{H} \in \mathcal{C}$ admit a dense set of C -induced complex structures, for all $C \in \mathbb{R}$ (Theorem 5.13).

- In Section 6 we give a proof of Desingularization Theorem for stable reflexive hyperholomorphic sheaves with isolated singularities (Theorem 6.1). We study the natural $\mathbb{C}^*$ -action on a local ring of a hyperkähler manifold defined in [V-d2]. We show that a sheaf F admitting a hyperholomorphic connection is equivariant with respect to this $\mathbb{C}^*$ -action. Then F can be desingularized by a blow-up, because any $\mathbb{C}^*$ -equivariant sheaf with an isolated singularity can be desingularized by a blow-up (Proposition 6.12).
- Section 7 is a primer on twistor transform and quaternionic-Kähler geometry. We give an exposition of the works of A. Swann ([Sw]), T. Nitta ([N1], [N2]) on quaternionic-Kähler manifolds and explain the direct and inverse twistor transform over hyperkähler and quaternionic-Kähler manifolds.
- Section 8 gives a correspondence between B_2 -bundles on a quaternionic-Kähler manifold, and $\mathbb{C}^*$ -equivariant holomorphic bundles on the

twistor space of the corresponding hyperkähler manifold constructed by A. Swann. This is called “Swann’s formalism for vector bundles”. We use this correspondence to prove that an associate graded sheaf of a hyperholomorphic sheaf is equipped with a natural connection which is compatible with quaternions. This implies polystability of the bundle $\pi^*F|_C$ (see Subsection 1.3).

- In Section 9, we use the polystability of the bundle $\pi^*F|_C$ to show that a hyperholomorphic deformation of a hyperholomorphic bundle is again a bundle. Together with results on C -restricted complex structures and Maruyama’s compactification ([Ma2]), this implies that the moduli of semistable bundles are compact, under conditions of C -restrictness and non-existence of trianalytic subvarieties (Theorem 9.11).
- In Section 10, we apply these results to the hyperkähler geometry. Using the desingularization theorem for singular hyperkähler manifolds (Theorem 2.16), we prove that the moduli of stable deformations of a hyperholomorphic bundle has a compact hyperkähler desingularization (Theorem 10.17). We give an exposition of the theory of twistor paths, which allows one to identify the categories of stable bundles for different Kähler structures on the same hyperkähler manifold (Theorem 10.8). These results allow one to weaken the conditions necessary for compactness of the moduli spaces of vector bundles. Finally, we give a conjectural exposition of how these results can be used to obtain new examples of compact hyperkähler manifolds.

2 Hyperkähler manifolds

2.1 Hyperkähler manifolds

This subsection contains a compression of the basic and best known results and definitions from hyperkähler geometry, found, for instance, in [Bes] or in [Bea].

Definition 2.1: ([Bes]) A **hyperkähler manifold** is a Riemannian manifold M endowed with three complex structures I , J and K , such that the following holds.

- (i) the metric on M is Kähler with respect to these complex structures and

- (ii) I , J and K , considered as endomorphisms of a real tangent bundle, satisfy the relation $I \circ J = -J \circ I = K$.

The notion of a hyperkähler manifold was introduced by E. Calabi ([C]).

Clearly, a hyperkähler manifold has a natural action of the quaternion algebra $\mathbb{H}$ in its real tangent bundle TM . Therefore its complex dimension is even. For each quaternion $L \in \mathbb{H}$, $L^2 = -1$, the corresponding automorphism of TM is an almost complex structure. It is easy to check that this almost complex structure is integrable ([Bes]).

Definition 2.2: Let M be a hyperkähler manifold, and L a quaternion satisfying $L^2 = -1$. The corresponding complex structure on M is called **an induced complex structure**. The M , considered as a Kähler manifold, is denoted by (M, L) . In this case, the hyperkähler structure is called **compatible with the complex structure L** .

Let M be a compact complex variety. We say that M is **of hyperkähler type** if M admits a hyperkähler structure compatible with the complex structure.

Definition 2.3: Let M be a complex manifold and Θ a closed holomorphic 2-form over M such that $\Theta^n = \Theta \wedge \Theta \wedge \dots$, is a nowhere degenerate section of a canonical class of M ($2n = \dim_{\mathbb{C}}(M)$). Then M is called **holomorphically symplectic**.

Let M be a hyperkähler manifold; denote the Riemannian form on M by $\langle \cdot, \cdot \rangle$. Let the form $\omega_I := \langle I(\cdot), \cdot \rangle$ be the usual Kähler form which is closed and parallel (with respect to the Levi-Civita connection). Analogously defined forms ω_J and ω_K are also closed and parallel.

A simple linear algebraic consideration ([Bes]) shows that the form $\Theta := \omega_J + \sqrt{-1}\omega_K$ is of type $(2, 0)$ and, being closed, this form is also holomorphic. Also, the form Θ is nowhere degenerate, as another linear algebraic argument shows. It is called **the canonical holomorphic symplectic form of a manifold M** . Thus, for each hyperkähler manifold M , and an induced complex structure L , the underlying complex manifold (M, L) is holomorphically symplectic. The converse assertion is also true:

Theorem 2.4: ([Bea], [Bes]) Let M be a compact holomorphically symplectic Kähler manifold with the holomorphic symplectic form Θ , a Kähler class $[\omega] \in H^{1,1}(M)$ and a complex structure I . Let $n = \dim_{\mathbb{C}} M$. Assume that $\int_M \omega^n = \int_M (\operatorname{Re} \Theta)^n$. Then there is a unique hyperkähler structure $(I, J, K, (\cdot, \cdot))$ over M such that the cohomology class of the symplectic form $\omega_I = (\cdot, I\cdot)$ is equal to $[\omega]$ and the canonical symplectic form $\omega_J + \sqrt{-1} \omega_K$ is equal to Θ .

Theorem 2.4 follows from the conjecture of Calabi, proven by Yau ([Y]).

■

Let M be a hyperkähler manifold. We identify the group $SU(2)$ with the group of unitary quaternions. This gives a canonical action of $SU(2)$ on the tangent bundle, and all its tensor powers. In particular, we obtain a natural action of $SU(2)$ on the bundle of differential forms.

Lemma 2.5: The action of $SU(2)$ on differential forms commutes with the Laplacian.

Proof: This is Proposition 1.1 of [V2(II)]. ■

Thus, for compact M , we may speak of the natural action of $SU(2)$ in cohomology.

Further in this article, we use the following statement.

Lemma 2.6: Let ω be a differential form over a hyperkähler manifold M . The form ω is $SU(2)$ -invariant if and only if it is of Hodge type (p, p) with respect to all induced complex structures on M .

Proof: This is [V1], Proposition 1.2. ■

2.2 Simple hyperkähler manifolds

Definition 2.7: ([Bea]) A connected simply connected compact hyperkähler manifold M is called **simple** if M cannot be represented as a product of two hyperkähler manifolds:

$$M \neq M_1 \times M_2, \text{ where } \dim M_1 > 0 \text{ and } \dim M_2 > 0$$

Bogomolov proved that every compact hyperkähler manifold has a finite covering which is a product of a compact torus and several simple hyperkähler manifolds. Bogomolov's theorem implies the following result ([Bea]):

Theorem 2.8: Let M be a compact hyperkähler manifold. Then the following conditions are equivalent.

- (i) M is simple
- (ii) M satisfies $H^1(M, \mathbb{R}) = 0$, $H^{2,0}(M) = \mathbb{C}$, where $H^{2,0}(M)$ is the space of $(2, 0)$ -classes taken with respect to any of induced complex structures.

■

2.3 Trianalytic subvarieties in hyperkähler manifolds.

In this subsection, we give a definition and basic properties of trianalytic subvarieties of hyperkähler manifolds. We follow [V2(II)], [V-d2].

Let M be a compact hyperkähler manifold, $\dim_{\mathbb{R}} M = 2m$.

Definition 2.9: Let $N \subset M$ be a closed subset of M . Then N is called **trianalytic** if N is a complex analytic subset of (M, L) for any induced complex structure L .

Let I be an induced complex structure on M , and $N \subset (M, I)$ be a closed analytic subvariety of (M, I) , $\dim_{\mathbb{C}} N = n$. Consider the homology class represented by N . Let $[N] \in H^{2m-2n}(M)$ denote the Poincaré dual cohomology class, so called **fundamental class** of N . Recall that the hyperkähler structure induces the action of the group $SU(2)$ on the space $H^{2m-2n}(M)$.

Theorem 2.10: Assume that $[N] \in H^{2m-2n}(M)$ is invariant with respect to the action of $SU(2)$ on $H^{2m-2n}(M)$. Then N is trianalytic.

Proof: This is Theorem 4.1 of [V2(II)]. ■

The following assertion is the key to the proof of Theorem 2.10 (see [V2(II)] for details).

Proposition 2.11: (Wirtinger's inequality) Let M be a compact hyperkähler manifold, I an induced complex structure and $X \subset (M, I)$ a closed complex subvariety for complex dimension k . Let J be an induced complex structure, $J \neq \pm I$, and ω_I, ω_J the associated Kähler forms. Consider the

numbers

$$\deg_I X := \int_X \omega_I^k, \quad \deg_J X := \int_X \omega_J^k$$

Then $\deg_I X \geq |\deg_J X|$, and the inequality is strict unless X is trianalytic.

■

Remark 2.12: Trianalytic subvarieties have an action of quaternion algebra in the tangent bundle. In particular, the real dimension of such subvarieties is divisible by 4.

Definition 2.13: Let M be a complex manifold admitting a hyperkähler structure $\mathcal{H}$. We say that M is **of general type** or **generic** with respect to $\mathcal{H}$ if all elements of the group

$$\bigoplus_p H^{p,p}(M) \cap H^{2p}(M, \mathbb{Z}) \subset H^*(M)$$

are $SU(2)$ -invariant. We say that M is **Mumford–Tate generic** if for all $n \in \mathbb{Z}^{>0}$, all the cohomology classes

$$\alpha \in \bigoplus_p H^{p,p}(M^n) \cap H^{2p}(M^n, \mathbb{Z}) \subset H^*(M^n)$$

are $SU(2)$ -invariant. In other words, M is Mumford–Tate generic if for all $n \in \mathbb{Z}^{>0}$, the n -th power M^n is generic. Clearly, Mumford–Tate generic implies generic.

Proposition 2.14: Let M be a compact manifold, $\mathcal{H}$ a hyperkähler structure on M and S be the set of induced complex structures over M . Denote by $S_0 \subset S$ the set of $L \in S$ such that (M, L) is Mumford–Tate generic with respect to $\mathcal{H}$. Then S_0 is dense in S . Moreover, the complement $S \setminus S_0$ is countable.

Proof: This is Proposition 2.2 from [V2(II)] ■

Theorem 2.10 has the following immediate corollary:

Corollary 2.15: Let M be a compact holomorphically symplectic manifold. Assume that M is of general type with respect to a hyperkähler structure $\mathcal{H}$. Let $S \subset M$ be closed complex analytic subvariety. Then S is trianalytic with respect to $\mathcal{H}$.

■

In [V-d3], [V-d], [V-d2], we gave a number of equivalent definitions of a singular hyperkähler and hypercomplex variety. We refer the reader to [V-d2] for the precise definition; for our present purposes it suffices to say that all trianalytic subvarieties are hyperkähler varieties. The following Desingularization Theorem is very useful in the study of trianalytic subvarieties.

Theorem 2.16: ([V-d2]) Let M be a hyperkähler or a hypercomplex variety, I an induced complex structure. Let

$$(\widetilde{M}, I) \xrightarrow{n} (M, I)$$

be the normalization of (M, I) . Then $(\widetilde{M}, I)$ is smooth and has a natural hyperkähler structure $\mathcal{H}$, such that the associated map $n : (\widetilde{M}, I) \rightarrow (M, I)$ agrees with $\mathcal{H}$. Moreover, the hyperkähler manifold $\widetilde{M} := (\widetilde{M}, I)$ is independent from the choice of induced complex structure I .

■

Let M be a K3 surface, and $M^{[n]}$ be a Hilbert scheme of points on M . Then $M^{[n]}$ admits a hyperkähler structure ([Bea]). In [V5], we proved the following theorem.

Theorem 2.17: Let M be a complex K3 surface without automorphisms. Assume that M is Mumford-Tate generic with respect to some hyperkähler structure. Consider the Hilbert scheme $M^{[n]}$ of points on M . Pick a hyperkähler structure on $M^{[n]}$ which is compatible with the complex structure. Then $M^{[n]}$ has no proper trianalytic subvarieties.

■

2.4 Hyperholomorphic bundles

This subsection contains several versions of a definition of hyperholomorphic connection in a complex vector bundle over a hyperkähler manifold. We follow [V1].

Let B be a holomorphic vector bundle over a complex manifold M , ∇ a connection in B and $\Theta \in \Lambda^2 \otimes \text{End}(B)$ be its curvature. This connection is called **compatible with a holomorphic structure** if $\nabla_X(\zeta) = 0$ for any holomorphic section ζ and any antiholomorphic tangent vector field

$X \in T^{0,1}(M)$. If there exists a holomorphic structure compatible with the given Hermitian connection then this connection is called **integrable**.

One can define a **Hodge decomposition** in the space of differential forms with coefficients in any complex bundle, in particular, $End(B)$.

Theorem 2.18: Let ∇ be a Hermitian connection in a complex vector bundle B over a complex manifold. Then ∇ is integrable if and only if $\Theta \in \Lambda^{1,1}(M, End(B))$, where $\Lambda^{1,1}(M, End(B))$ denotes the forms of Hodge type $(1,1)$. Also, the holomorphic structure compatible with ∇ is unique.

Proof: This is Proposition 4.17 of [Ko], Chapter I. ■

This result has the following more general version:

Proposition 2.19: Let ∇ be an arbitrary (not necessarily Hermitian) connection in a complex vector bundle B . Then ∇ is integrable if and only if its $(0,1)$ -part has square zero.

■

This proposition is a version of Newlander-Nirenberg theorem. For vector bundles, it was proven by Atiyah and Bott.

Definition 2.20: Let B be a Hermitian vector bundle with a connection ∇ over a hyperkähler manifold M . Then ∇ is called **hyperholomorphic** if ∇ is integrable with respect to each of the complex structures induced by the hyperkähler structure.

As follows from Theorem 2.18, ∇ is hyperholomorphic if and only if its curvature Θ is of Hodge type $(1,1)$ with respect to any of complex structures induced by a hyperkähler structure.

As follows from Lemma 2.6, ∇ is hyperholomorphic if and only if Θ is a $SU(2)$ -invariant differential form.

Example 2.21: (Examples of hyperholomorphic bundles)

- (i) Let M be a hyperkähler manifold, and TM be its tangent bundle equipped with the Levi-Civita connection ∇ . Consider a complex structure on TM induced from the quaternion action. Then ∇ is a Hermitian connection which is integrable with respect to each induced complex structure, and hence, is Yang–Mills.

- (ii) For B a hyperholomorphic bundle, all its tensor powers are also hyperholomorphic.
- (iii) Thus, the bundles of differential forms on a hyperkähler manifold are also hyperholomorphic.

2.5 Stable bundles and Yang–Mills connections.

This subsection is a compendium of the most basic results and definitions from the Yang–Mills theory over Kähler manifolds, concluding in the fundamental theorem of Uhlenbeck–Yau [UY].

Definition 2.22: Let F be a coherent sheaf over an n -dimensional compact Kähler manifold M . We define $\deg(F)$ as

$$\deg(F) = \int_M \frac{c_1(F) \wedge \omega^{n-1}}{\text{vol}(M)}$$

and $\text{slope}(F)$ as

$$\text{slope}(F) = \frac{1}{\text{rank}(F)} \cdot \deg(F).$$

The number $\text{slope}(F)$ depends only on a cohomology class of $c_1(F)$.

Let F be a coherent sheaf on M and $F' \subset F$ its proper subsheaf. Then F' is called **destabilizing subsheaf** if $\text{slope}(F') \geq \text{slope}(F)$

A coherent sheaf F is called **stable**¹ if it has no destabilizing subsheaves. A coherent sheaf F is called **semistable** if for all destabilizing subsheaves $F' \subset F$, we have $\text{slope}(F') = \text{slope}(F)$.

Later on, we usually consider the bundles B with $\deg(B) = 0$.

Let M be a Kähler manifold with a Kähler form ω . For differential forms with coefficients in any vector bundle there is a Hodge operator $L : \eta \longrightarrow \omega \wedge \eta$. There is also a fiberwise-adjoint Hodge operator Λ (see [GH]).

Definition 2.23: Let B be a holomorphic bundle over a Kähler manifold M with a holomorphic Hermitian connection ∇ and a curvature $\Theta \in \Lambda^{1,1} \otimes \text{End}(B)$. The Hermitian metric on B and the connection ∇ defined by this metric are called **Yang–Mills** if

$$\Lambda(\Theta) = \text{constant} \cdot \text{Id} \Big|_B,$$

¹In the sense of Mumford–Takemoto

where Λ is a Hodge operator and $\text{Id}|_B$ is the identity endomorphism which is a section of $\text{End}(B)$.

Further on, we consider only these Yang–Mills connections for which this constant is zero.

A holomorphic bundle is called **indecomposable** if it cannot be decomposed onto a direct sum of two or more holomorphic bundles.

The following fundamental theorem provides examples of Yang–Mills bundles.

Theorem 2.24: (Uhlenbeck-Yau) Let B be an indecomposable holomorphic bundle over a compact Kähler manifold. Then B admits a Hermitian Yang–Mills connection if and only if it is stable, and this connection is unique.

Proof: [UY]. ■

Proposition 2.25: Let M be a hyperkähler manifold, L an induced complex structure and B be a complex vector bundle over (M, L) . Then every hyperholomorphic connection ∇ in B is Yang–Mills and satisfies $\Lambda(\Theta) = 0$ where Θ is a curvature of ∇ .

Proof: We use the definition of a hyperholomorphic connection as one with $SU(2)$ -invariant curvature. Then Proposition 2.25 follows from the

Lemma 2.26: Let $\Theta \in \Lambda^2(M)$ be a $SU(2)$ -invariant differential 2-form on M . Then $\Lambda_L(\Theta) = 0$ for each induced complex structure L .²

Proof: This is Lemma 2.1 of [V1]. ■

Let M be a compact hyperkähler manifold, I an induced complex structure. For any stable holomorphic bundle on (M, I) there exists a unique Hermitian Yang–Mills connection which, for some bundles, turns out to be hyperholomorphic. It is possible to tell when this happens.

Theorem 2.27: Let B be a stable holomorphic bundle over (M, I) , where M is a hyperkähler manifold and I is an induced complex structure over M . Then B admits a compatible hyperholomorphic connection if and only if the first two Chern classes $c_1(B)$ and $c_2(B)$ are $SU(2)$ -invariant.³

²By Λ_L we understand the Hodge operator Λ associated with the Kähler complex structure L .

³We use Lemma 2.5 to speak of action of $SU(2)$ in cohomology of M .

Proof: This is Theorem 2.5 of [V1]. ■

2.6 Twistor spaces

Let M be a hyperkähler manifold. Consider the product manifold $X = M \times S^2$. Embed the sphere $S^2 \subset \mathbb{H}$ into the quaternion algebra $\mathbb{H}$ as the subset of all quaternions J with $J^2 = -1$. For every point $x = m \times J \in X = M \times S^2$ the tangent space $T_x X$ is canonically decomposed $T_x X = T_m M \oplus T_J S^2$. Identify $S^2 = \mathbb{C}P^1$ and let $I_J : T_J S^2 \rightarrow T_J S^2$ be the complex structure operator. Let $I_m : T_m M \rightarrow T_m M$ be the complex structure on M induced by $J \in S^2 \subset \mathbb{H}$.

The operator $I_x = I_m \oplus I_J : T_x X \rightarrow T_x X$ satisfies $I_x \circ I_x = -1$. It depends smoothly on the point x , hence defines an almost complex structure on X . This almost complex structure is known to be integrable (see [Sal]).

Definition 2.28: The complex manifold $\langle X, I_x \rangle$ is called *the twistor space* for the hyperkähler manifold M , denoted by $\text{Tw}(M)$. This manifold is equipped with a real analytic projection $\sigma : \text{Tw}(M) \rightarrow M$ and a complex analytic projection $\pi : \text{Tw}(M) \rightarrow \mathbb{C}P^1$.

The twistor space $\text{Tw}(M)$ is not, generally speaking, a Kähler manifold. For M compact, it is easy to show that $\text{Tw}(M)$ does not admit a Kähler metric. We consider $\text{Tw}(M)$ as a Hermitian manifold with the product metric.

Definition 2.29: Let X be an n -dimensional Hermitian manifold and let $\sqrt{-1}\omega$ be the imaginary part of the metric on X . Thus ω is a real $(1, 1)$ -form. Assume that the form ω satisfies the following condition of Li and Yau ([LY]).

$$\omega^{n-2} \wedge d\omega = 0. \quad (2.1)$$

Such Hermitian metrics are called **metrics satisfying the condition of Li–Yau**.

For a closed real 2-form η let

$$\deg \eta = \int_X \omega^{n-1} \wedge \eta.$$

The condition (2.1) ensures that $\deg \eta$ depends only on the cohomology class of η . Thus it defines a degree functional $\deg : H^2(X, \mathbb{R}) \rightarrow \mathbb{R}$. This

functional allows one to repeat verbatim the Mumford-Takemoto definitions of stable and semistable bundles in this more general situation. Moreover, the Hermitian Yang-Mills equations also carry over word-by-word. Li and Yau proved a version of Uhlenbeck–Yau theorem in this situation ([LY]; see also Theorem 3.8).

Proposition 2.30: Let M be a hyperkähler manifold and $\mathrm{Tw}(M)$ its twistor space, considered as a Hermitian manifold. Then $\mathrm{Tw}(M)$ satisfies the conditions of Li–Yau.

Proof: [KV], Proposition 4.5. ■

3 Hyperholomorphic sheaves

3.1 Stable sheaves and Yang-Mills connections

In [BS], S. Bando and Y.-T. Siu developed the machinery allowing one to apply the methods of Yang-Mills theory to torsion-free coherent sheaves. In the course of this paper, we apply their work to generalise the results of [V1]. In this Subsection, we give a short exposition of their results.

Definition 3.1: Let X be a complex manifold, and F a coherent sheaf on X . Consider the sheaf $F^* := \mathcal{H}om_{\mathcal{O}_X}(F, \mathcal{O}_X)$. There is a natural functorial map $\rho_F : F \longrightarrow F^{**}$. The sheaf F^{**} is called a **reflexive hull**, or **reflex-ization** of F . The sheaf F is called **reflexive** if the map $\rho_F : F \longrightarrow F^{**}$ is an isomorphism.

Remark 3.2: For all coherent sheaves F , the map $\rho_{F^*} : F^* \longrightarrow F^{***}$ is an isomorphism ([OSS], Ch. II, the proof of Lemma 1.1.12). Therefore, a reflexive hull of a sheaf is always reflexive.

Claim 3.3: Let X be a Kähler manifold, and F a torsion-free coherent sheaf over X . Then F (semi)stable if and only if F^{**} is (semi)stable.

Proof: This is [OSS], Ch. II, Lemma 1.2.4. ■

Definition 3.4: Let X be a Kähler manifold, and F a coherent sheaf over X . The sheaf F is called **polystable** if F is a direct sum of stable sheaves.

The admissible Hermitian metrics, introduced by Bando and Siu in [BS], play the role of the ordinary Hermitian metrics for vector bundles.

Let X be a Kähler manifold. In Hodge theory, one considers the operator $\Lambda : \Lambda^{p,q}(X) \longrightarrow \Lambda^{p-1,q-1}(X)$ acting on differential forms on X , which is adjoint to the multiplication by the Kähler form. This operator is defined on differential forms with coefficient in every bundle. Considering a curvature Θ of a bundle B as a 2-form with coefficients in $\text{End}(B)$, we define the expression $\Lambda\Theta$ which is a section of $\text{End}(B)$.

Definition 3.5: Let X be a Kähler manifold, and F a reflexive coherent sheaf over X . Let $U \subset X$ be the set of all points at which F is locally trivial. By definition, the restriction $F|_U$ of F to U is a bundle. An **admissible metric** on F is a Hermitian metric h on the bundle $F|_U$ which satisfies the following assumptions

- (i) the curvature Θ of (F, h) is square integrable, and
- (ii) the corresponding section $\Lambda\Theta \in \text{End}(F|_U)$ is uniformly bounded.

Definition 3.6: Let X be a Kähler manifold, F a reflexive coherent sheaf over X , and h an admissible metric on F . Consider the corresponding Hermitian connection ∇ on $F|_U$. The metric h and the connection ∇ are called **Yang-Mills** if its curvature satisfies

$$\Lambda\Theta \in \text{End}(F|_U) = c \cdot \text{id}$$

where c is a constant and id the unit section $\text{id} \in \text{End}(F|_U)$.

Further in this paper, we shall only consider Yang-Mills connections with $\Lambda\Theta = 0$.

Remark 3.7: By Gauss-Bonnet formule, the constant c is equal to $\deg(F)$, where $\deg(F)$ is the degree of F (Definition 2.22).

One of the main results of [BS] is the following analogue of Uhlenbeck–Yau theorem (Theorem 2.24).

Theorem 3.8: Let M be a compact Kähler manifold, or a compact Hermitian manifold satisfying conditions of Li-Yau (Definition 2.29), and F

a coherent sheaf without torsion. Then F admits an admissible Yang–Mills metric if and only if F is polystable. Moreover, if F is stable, then this metric is unique, up to a constant multiplier.

Proof: In [BS], Theorem 3.8 is proved for Kähler M ([BS], Theorem 3). It is easy to adapt this proof for Hermitian manifolds satisfying conditions of Li–Yau. ■

Remark 3.9: Clearly, the connection, corresponding to a metric on F , does not change when the metric is multiplied by a scalar. The Yang–Mills metric on a polystable sheaf is unique up to a componentwise multiplication by scalar multipliers. Thus, the Yang–Mills connection of Theorem 3.8 is unique.

Another important theorem of [BS] is the following.

Theorem 3.10: Let (F, h) be a holomorphic vector bundle with a Hermitian metric h defined on a Kähler manifold X (not necessarily compact nor complete) outside a closed subset S with locally finite Hausdorff measure of real co-dimension 4. Assume that the curvature tensor of F is locally square integrable on X . Then F extends to the whole space X as a reflexive sheaf $\mathcal{F}$. Moreover, if the metric h is Yang–Mills, then h can be smoothly extended as a Yang–Mills metric over the place where $\mathcal{F}$ is locally free.

Proof: This is [BS], Theorem 2. ■

3.2 Stable and semistable sheaves over hyperkähler manifolds

Let M be a compact hyperkähler manifold, I an induced complex structure, F a torsion-free coherent sheaf over (M, I) and F^{**} its reflexization. Recall that the cohomology of M are equipped with a natural $SU(2)$ -action (Lemma 2.5). The motivation for the following definition is Theorem 2.27 and Theorem 3.8.

Definition 3.11: Assume that the first two Chern classes of the sheaves F , F^{**} are $SU(2)$ -invariant. Then F is called **stable hyperholomorphic** if F is stable, and **semistable hyperholomorphic** if F can be obtained as a successive extension of stable hyperholomorphic sheaves.

Remark 3.12: The slope of a hyperholomorphic sheaf is zero, because a degree of an $SU(2)$ -invariant 2-form is zero (Lemma 2.26).

Claim 3.13: Let F be a semistable coherent sheaf over (M, I) . Then the following conditions are equivalent.

- (i) F is stable hyperholomorphic
- (ii) Consider the support S of the sheaf F^{**}/F as a complex subvariety of (M, I) . Let $X_1, \dots, X_n$ be the set of irreducible components of S of codimension 2. Then X_i is trianalytic for all i , and the sheaf F^{**} is stable hyperholomorphic.

Proof: Consider an exact sequence

$$0 \longrightarrow F \longrightarrow F^{**} \longrightarrow F^{**}/F \longrightarrow 0.$$

Let $[F/F^{**}] \in H^4(M)$ be the fundamental class of the union of all codimension-2 components of support of the sheaf F/F^{**} , taken with appropriate multiplicities. Then, $c_2(F^{**}/F) = -[F/F^{**}]$. From the product formula for Chern classes, it follows that

$$c_2(F) = c_2(F_i^{**}) + c_2(F^{**}/F) = c_2(F_i^{**}) - [F/F^{**}]. \quad (3.1)$$

Clearly, if all X_i are trianalytic then the class $[F/F^{**}]$ is $SU(2)$ -invariant. Thus, if the sheaf F^{**} is hyperholomorphic and all X_i are trianalytic, then the second Chern class of F is $SU(2)$ -invariant, and F is hyperholomorphic. Conversely, assume that F is hyperholomorphic. We need to show that all X_i are trianalytic. By definition,

$$[F/F^{**}] = \sum_i \lambda_i [X_i]$$

where $[X_i]$ denotes the fundamental class of X_i , and λ_i is the multiplicity of F/F^{**} at X_i . By (3.1), (F hyperholomorphic) implies that the class $[F/F^{**}]$ is $SU(2)$ -invariant. Since $[F/F^{**}]$ is $SU(2)$ -invariant, we have

$$\sum_i \lambda_i \deg_J(X_i) = \sum_i \lambda_i \deg_I(X_i).$$

By Wirtinger's inequality (Proposition 2.11),

$$\deg_J(X_i) \leq \deg_I(X_i),$$

and the equality is reached only if X_i is trianalytic. By definition, all the numbers λ_i are positive. Therefore,

$$\sum_i \lambda_i \deg_J(X_i) \leq \sum_i \lambda_i \deg_I(X_i).$$

and the equality is reached only if all the subvarieties X_i are trianalytic. This finishes the proof of Claim 3.13. ■

Claim 3.14: Let M be a compact hyperkähler manifold, and I an induced complex structure of general type. Then all torsion-free coherent sheaves over (M, I) are semistable hyperholomorphic.

Proof: Let F be a torsion-free coherent sheaf over (M, I) . Clearly from the definition of induced complex structure of general type, the sheaves F and F^{**} have $SU(2)$ -invariant Chern classes. Now, all $SU(2)$ -invariant 2-forms have degree zero (Lemma 2.26), and thus, F is semistable. ■

3.3 Hyperholomorphic connection in torsion-free sheaves

Let M be a hyperkähler manifold, I an induced complex structure, and F a torsion-free sheaf over (M, I) . Consider the natural $SU(2)$ -action in the bundle $\Lambda^i(M, B)$ of the differential i -forms with coefficients in a vector bundle B . Let $\Lambda_{inv}^i(M, B) \subset \Lambda^i(M, B)$ be the bundle of $SU(2)$ -invariant i -forms.

Definition 3.15: Let $X \subset (M, I)$ be a complex subvariety of codimension at least 2, such that $F|_{M \setminus X}$ is a bundle, h be an admissible metric on $F|_{M \setminus X}$ and ∇ the associated connection. Then ∇ is called **hyperholomorphic** if its curvature

$$\Theta_\nabla = \nabla^2 \in \Lambda^2 \left(M, \text{End} \left(F|_{M \setminus X} \right) \right)$$

is $SU(2)$ -invariant, i. e. belongs to $\Lambda_{inv}^2 \left(M, \text{End} \left(F|_{M \setminus X} \right) \right)$.

Claim 3.16: The singularities of a hyperholomorphic connection form a trianalytic subvariety in M .

Proof: Let J be an induced complex structure on M , and U the set of all points of (M, I) where F is non-singular. Clearly, (F, ∇) is a bundle with admissible connection on (U, J) . Therefore, the holomorphic structure on $F|_{(U, J)}$ can be extended to (M, J) . Thus, the singular set of F is holomorphic with respect to J . This proves Claim 3.16. ■

Proposition 3.17: Let M be a compact hyperkähler manifold, I an induced complex structure and F a reflexive sheaf admitting a hyperholomorphic connection. Then F is a polystable hyperholomorphic sheaf.

Proof: By Remark 3.20 and Theorem 3.8, F is polystable. We need only to show that the Chern classes $c_1(F)$ and $c_2(F)$ are $SU(2)$ -invariant. Let $U \subset M$ be the maximal open subset of M such that $F|_U$ is locally trivial. By Theorem 3.10, the metric h and the connection ∇ can be extended to U . Let $\text{Tw} U \subset \text{Tw} M$ be the corresponding twistor space, and $\sigma : \text{Tw} U \rightarrow U$ the standard map. Consider the bundle $\sigma^*F|_U$, equipped with a connection $\sigma^*\nabla$. It is well known¹ that $\sigma^*F|_U$ is a bundle with an admissible Yang-Mills metric (we use Yang-Mills in the sense of Li-Yau, see Definition 2.29). By Theorem 3.10, $\sigma^*F|_U$ can be extended to a reflexive sheaf $\mathcal{F}$ on $\text{Tw} M$. Clearly, this extension coincides with the push-forward of $\sigma^*F|_U$. The singular set $\tilde{S}$ of $\mathcal{F}$ is a pull-back of the singular set S of F . Thus, S is trianalytic. By desingularization theorem (Theorem 2.16), S can be desingularized to a hyperkähler manifold in such a way that its twistors form a desingularization of $\mathcal{S}$. From the exact description of the singularities of $\mathcal{S}$, provided by the desingularization theorem, we obtain that the standard projection $\pi : \mathcal{S} \rightarrow \mathbb{C}P^1$ is flat. By the following lemma, the restriction of $\mathcal{F}$ to the fiber $(M, I) = \pi^{-1}(\{I\})$ of π coincides with F .

Lemma 3.18: Let $\pi : X \rightarrow Y$ be a map of complex varieties, and $S \hookrightarrow X$ a subvariety of X of codimension at least 2, which is flat over Y . Denote by $U \xrightarrow{j} X$ the complement $U = (X \setminus S)$. Let F be a vector bundle over U , and j_*F its push-forward. Then the restriction of j_*F to the fibers of π is reflexive.

Proof: Let $Z = \pi^{-1}(\{y\})$ be a fiber of π . Since S is flat over Y and of codimension at least 2, we have $j_*(\mathcal{O}_{Z \cap U}) = \mathcal{O}_Z$. Clearly, for an open embedding $\gamma : T_1 \rightarrow T_2$ and coherent sheaves A, B on T_1 , we have $\gamma_*(A \otimes B) = \gamma_*A \otimes \gamma_*B$. Thus, for all coherent sheaves A on U , we have

$$j_*A \otimes \mathcal{O}_Z = j_*(A \otimes \mathcal{O}_{Z \cap U}). \quad (3.2)$$

This implies that $j_*(F|_Z) = j_*F|_Z$. It is well known ([OSS], Ch. II, 1.1.12; see also Lemma 9.2) that a push-forward of a reflexive sheaf under an open embedding γ is reflexive, provided that the complement of the image of γ

¹See for instance the section “Direct and inverse twistor transform” in [KV].

has codimension at least 2. Therefore, $j_*F|_Z$ is a reflexive sheaf over Z . This proves Lemma 3.18. ■

Return to the proof of Proposition 3.17. Consider the sheaf $\mathcal{F}$ on the twistor space constructed above. Since $\mathcal{F}$ is reflexive, its singularities have codimension at least 3 ([OSS], Ch. II, 1.1.10). Therefore, $\mathcal{F}$ is flat in codimension 2, and the first two Chern classes of $F = \mathcal{F}|_{\pi^{-1}(I)}$ can be obtained by restricting the first two Chern classes of $\mathcal{F}$ to the subvariety $(M, I) = \pi^{-1}(I) \subset \text{Tw}(M)$. It remains to show that such restriction is $SU(2)$ -invariant. Clearly, $H^2((M, I)) = H^2((M, I) \setminus S)$, and $H^4((M, I)) = H^4((M, I) \setminus S)$. Therefore,

$$c_1(\mathcal{F}|_{(M, I)}) = c_1(\mathcal{F}|_{(M, I) \setminus S})$$

and

$$c_2(\mathcal{F}|_{(M, I)}) = c_2(\mathcal{F}|_{(M, I) \setminus S}).$$

On the other hand, the restriction $\mathcal{F}|_{\text{Tw}(M) \setminus S}$ is a bundle. Therefore, the classes

$$c_1(\mathcal{F}|_{(M, I) \setminus S}), \quad c_2(\mathcal{F}|_{(M, I) \setminus S})$$

are independent from $I \in \mathbb{CP}^1$. On the other hand, these classes are of type (p, p) with respect to all induced complex structures $I \in \mathbb{CP}^1$. By Lemma 2.6, this implies that the classes $c_1(\mathcal{F}|_{(M, I)})$, $c_1(\mathcal{F}|_{(M, I)})$ are $SU(2)$ -invariant. As we have shown above, these two classes are equal to the first Chern classes of F . Proposition 3.17 is proven. ■

3.4 Existence of hyperholomorphic connections

The following theorem is the main result of this section.

Theorem 3.19: Let M be a compact hyperkähler manifold, I an induced complex structure and F a reflexive sheaf on (M, I) . Then F admits a hyperholomorphic connection if and only if F is polystable hyperholomorphic in the sense of Definition 3.11.

Remark 3.20: From Lemma 2.26, it is clear that a hyperholomorphic connection is always Yang-Mills. Therefore, such a connection is unique (Theorem 3.8).

The “only if” part of Theorem 3.19 is Proposition 3.17. The proof of “if” part of Theorem 3.19 takes the rest of this subsection.

Let I be an induced complex structure. We denote the corresponding Hodge decomposition on differential forms by $\Lambda^*(M) = \oplus \Lambda_I^{p,q}(M)$, and the standard Hodge operator by $\Lambda_I : \Lambda_I^{p,q}(M) \longrightarrow \Lambda_I^{p-1,q-1}(M)$. All these structures are defined on the differential forms with coefficients in a bundle. Let $\deg_I \eta := \int_M \text{Tr}(\Lambda_I)^k(\eta)$, for $\eta \in \Lambda^k(M, \text{End } B)$. The following claim follows from an elementary linear-algebraic computation.

Claim 3.21: Let M be a hyperkähler manifold, B a Hermitian vector bundle over M , and Θ a 2-form on M with coefficients in $\mathfrak{su}(B)$. Assume that

$$\Lambda_I \Theta = 0, \quad \Theta \in \Lambda_I^{1,1}(M, \text{End } B)$$

for some induced complex structure I . Assume, moreover, that Θ is square-integrable. Let J be another induced complex structure, $J \neq \pm I$. Then

$$\deg_I \Theta^2 \geq |\deg_J \Theta^2|,$$

and the equality is reached only if Θ is $SU(2)$ -invariant.

Proof: The following general argument is used.

Sublemma 3.22: Let M be a Kähler manifold, B a Hermitian vector bundle over M , and Ξ a square-integrable 2-form on M with coefficients in $\mathfrak{su}(B)$. Then:

(i) For

$$\Lambda_I \Xi = 0, \quad \Xi \in \Lambda_I^{1,1}(M, \text{End } B)$$

we have

$$\deg_I \Xi^2 = C \int_M |\Xi|^2 \text{Vol } M,$$

where $C = (4\pi^2 n(n-1))^{-1} M$.

(ii) For

$$\Xi \in \Lambda_I^{2,0}(M, \text{End } B) \oplus \Lambda_I^{0,2}(M, \text{End } B)$$

we have

$$\deg_I \Xi^2 = -C \int_M |\Xi|^2 \text{Vol } M,$$

where C is the same constant as appeared in (i).

Proof: The proof is based on a linear-algebraic computation (so-called Lübcke-type argument). The same computation is used to prove Hodge-Riemann bilinear relations. ■

Return to the proof of Claim 3.21. Let $\Theta = \Theta_J^{1,1} + \Theta_J^{2,0} + \Theta_J^{0,2}$ be the Hodge decomposition associated with J . The following Claim shows that $\Theta_J^{1,1}$ satisfies conditions of Sublemma 3.22 (i).

Claim 3.23: Let M be a hyperkähler manifold, I, L induced complex structures and Θ a 2-form on M satisfying

$$\Lambda_I \Theta = 0, \quad \Theta \in \Lambda_I^{1,1}(M).$$

Let $\Theta_L^{1,1}$ be the $(1,1)$ -component of Θ taken with respect to L . Then $\Lambda_L \Theta_L^{1,1} = 0$.

Proof: Clearly, $\Lambda_L \Theta_L^{1,1} = \Lambda_L \Theta$. Consider the natural Hermitian structure on the space of 2-forms. Since Θ is of type $(1,1)$ with respect to I , Θ is fiberwise orthogonal to the holomorphic symplectic form $\Omega = \omega_J + \sqrt{-1}\Omega_K \in \Lambda_I^{2,0}(M)$. By the same reason, Θ is orthogonal to $\bar{\Omega}$. Therefore, Θ is orthogonal to ω_J and ω_K . Since $\Lambda_I \Theta = 0$, Θ is also orthogonal to ω_I . The map Λ_L is a projection to the form ω_L which is a linear combination of ω_I, ω_J and ω_K . Since Θ is fiberwise orthogonal to ω_L , we have $\Lambda_L \Theta = 0$. ■

By Sublemma 3.22, we have

$$\deg_J \left(\Theta_J^{1,1} \right)^2 = C \int_M |\Theta_J^{1,1}|^2$$

and

$$\deg_J \left(\Theta_J^{2,0} + \Theta_J^{0,2} \right)^2 = -C \int_M |\Theta_J^{2,0} + \Theta_J^{0,2}|^2.$$

Thus,

$$\deg_J \Theta^2 = C \int_M \left| \Theta_J^{1,1} \right|^2 - C \int_M \left| \Theta_J^{2,0} + \Theta_J^{0,2} \right|^2.$$

On the other hand,

$$\deg_I \Theta^2 = C \int_M |\Theta|^2 = C \int_M \left| \Theta_J^{1,1} \right|^2 + C \int_M \left| \Theta_J^{2,0} + \Theta_J^{0,2} \right|^2.$$

Thus, $\deg_I \Theta^2 > |\deg_J \Theta^2|$ unless $\Theta_J^{2,0} + \Theta_J^{0,2} = 0$. On the other hand, $\Theta_J^{2,0} + \Theta_J^{0,2} = 0$ means that Θ is of type $(1,1)$ with respect to J . Consider

the standard $U(1)$ -action on differential forms associated with the complex structures I and J . These two $U(1)$ -actions generate the whole Lie group $SU(2)$ acting on $\Lambda^2(M)$ (here we use that $I \neq \pm J$). Since Θ is of type $(1,1)$ with respect to I and J , this form is $SU(2)$ -invariant. This proves Claim 3.21. ■

Return to the proof of Theorem 3.19. Let ∇ be the admissible Yang-Mills connection in F , and Θ its curvature. Recall that the form $Tr\Theta^2$ represents the cohomology class $2c_2(F) - \frac{r-1}{r}c_1(F)^2$, where c_i are Chern classes of F . Since the form $Tr\Theta^2$ is square-integrable, the integral

$$\deg_J \Theta^2 = \int_M Tr\Theta^2 \omega_J^{n-2}$$

makes sense. In [BS], it was shown how to approximate the connection ∇ by smooth connections, via the heat equation. This argument, in particular, was used to show that the value of integrals like $\int_M Tr\Theta^2 \omega_J^{n-2}$ can be computed through cohomology classes and the Gauss–Bonnet formula

$$Tr\Theta^2 = 2c_2(F) - \frac{r-1}{r}c_1(F)^2.$$

Since the classes $c_2(F)$, $c_1(F)$ are $SU(2)$ -invariant, we have

$$\deg_I \Theta^2 = \deg_J \Theta^2$$

for all induced complex structures I, J . By Claim 3.21, this implies that Θ is $SU(2)$ -invariant. Theorem 3.19 is proven. ■

The same argument implies the following corollary.

Corollary 3.24: Let M be a compact hyperkähler manifold, I an induced complex structure, F a stable reflexive sheaf on (M, I) , and J be an induced complex structure, $J \neq \pm I$. Then

$$\deg_I \left(2c_2(F) - \frac{r-1}{r}c_1(F)^2 \right) \geq \left| \deg_J \left(2c_2(F) - \frac{r-1}{r}c_1(F)^2 \right) \right|,$$

and the equality holds if and only if F is hyperholomorphic.

■

3.5 Tensor category of hyperholomorphic sheaves

This subsection is extraneous. Further on, we do not use the tensor structure on the category of hyperholomorphic sheaves. However, we need the canonical identification of the categories of hyperholomorphic sheaves associated with different induced complex structures.

From Bando-Siu (Theorem 3.8) it follows that on a compact Kähler manifold a tensor product of stable reflexive sheaves is polystable. Similarly, Theorem 3.19 implies that a tensor product of polystable hyperholomorphic sheaves is polystable hyperholomorphic. We define the following category.

Definition 3.25: Let M be a compact hyperkähler manifold and I an induced complex structure. Let $\mathcal{F}_{st}(M, I)$ be a category with objects reflexive polystable hyperholomorphic sheaves and morphisms as in category of coherent sheaves. This category is obviously additive. The tensor product on $\mathcal{F}_{st}(M, I)$ is induced from the tensor product of coherent sheaves.

Claim 3.26: The category $\mathcal{F}_{st}(M, I)$ is abelian. Moreover, it is a Tannakian tensor category.

Proof: Let $\varphi : F_1 \longrightarrow F_2$ be a morphism of hyperholomorphic sheaves. In Definition 2.22, we introduced a **slope** of a coherent sheaf. Clearly, $sl(F_1) \leqslant sl(\operatorname{im} \varphi) \leqslant sl(F_2)$. All hyperholomorphic sheaves have slope 0 by Remark 3.12. Thus, $sl(\operatorname{im} \varphi) = 0$ and the subsheaf $\operatorname{im} \varphi \subset F_2$ is destabilizing. Since F_2 is polystable, this sheaf is decomposed:

$$F_2 = \operatorname{im} \varphi \oplus \operatorname{coker} \varphi.$$

A similar argument proves that $F_1 = \ker \varphi \oplus \operatorname{coim} \varphi$, with all summands hyperholomorphic. This proves that $\mathcal{F}_{st}(M, I)$ is abelian. The Tannakian properties are clear. ■

The category $\mathcal{F}_{st}(M, I)$ does not depend from the choice of induced complex structure I :

Theorem 3.27: Let M be a compact hyperkähler manifold, I_1, I_2 induced complex structures and $\mathcal{F}_{st}(M, I_1), \mathcal{F}_{st}(M, I_2)$ the associated categories of polystable reflexive hyperholomorphic sheaves. Then, there exists a natural equivalence of tensor categories

$$\Phi_{I_1, I_2} : \mathcal{F}_{st}(M, I_1) \longrightarrow \mathcal{F}_{st}(M, I_2).$$

Proof: Let $F \in \mathcal{F}_{st}(M, I_1)$ be a reflexive polystable hyperholomorphic sheaf and ∇ the canonical admissible Yang-Mills connection. Consider the sheaf $\mathcal{F}$ on the twistor space $\mathrm{Tw}(M)$ constructed as in the proof of Proposition 3.17. Restricting $\mathcal{F}$ to $\pi^{-1}(I_2) \subset \mathrm{Tw}(M)$, we obtain a coherent sheaf F' on (M, I_2) . As we have shown in the proof of Proposition 3.17, the sheaf $(F')^{**}$ is polystable hyperholomorphic. Let $\Phi_{I_1, I_2}(F) := (F')^{**}$. It is easy to check that thus constructed map of objects gives a functor

$$\Phi_{I_1, I_2} : \mathcal{F}_{st}(M, I_1) \longrightarrow \mathcal{F}_{st}(M, I_2),$$

and moreover, $\Phi_{I_1, I_2} \circ \Phi_{I_2, I_1} = \mathrm{Id}$. This shows that Φ_{I_1, I_2} is an equivalence. Theorem 3.27 is proven. ■

Definition 3.28: By Theorem 3.27, the category $\mathcal{F}_{st}(M, I_1)$ is independent from the choice of induced complex structure. We call this category **the category of polystable hyperholomorphic reflexive sheaves on M** and denote it by $\mathcal{F}(M)$. The objects of $\mathcal{F}(M)$ are called **hyperholomorphic sheaves on M** . For a hyperholomorphic sheaf on M , we denote by F_I the corresponding sheaf from $\mathcal{F}_{st}(M, I_1)$.

Remark 3.29: Using the same argument as proves Theorem 10.8 (ii), it is easy to check that the category $\mathcal{F}(M)$ is a deformational invariant of M . That is, for two hyperkähler manifolds M_1, M_2 which are deformationally equivalent, the categories $\mathcal{F}(M_i)$ are also equivalent, assuming that $\mathrm{Pic}(M_1) = \mathrm{Pic}(M_2) = 0$. The proof of this result is essentially contained in [V3-bis].

Remark 3.30: As Deligne proved ([D]), for a each Tannakian category $\mathcal{C}$ equipped with a fiber functor, there exists a natural pro-algebraic group G such that $\mathcal{C}$ is a group of representations of G . For $\mathcal{F}(M)$, there are several natural fiber functors. The simplest one is defined for each induced complex structure I such that (M, I) is algebraic (such complex structures always exist, as proven in [F]; see also [V-a] and Subsection 4.1). Let $\mathcal{K}(M, I)$ is the space of rational functions on (M, I) . For $F \in \mathcal{F}_{st}(M, I)$, consider the functor $F \longrightarrow \eta_I(F)$, where $\eta_I(F)$ is the space of global sections of $F \otimes \mathcal{K}(M, I)$. This is clearly a fiber functor, which associates to $\mathcal{F}(M)$ the group G_I . The corresponding pro-algebraic group G_I is a deformational, that is, topological, invariant of the hyperkähler manifold.

4 Cohomology of hyperkähler manifolds

This section contains a serie of preliminary results which are used further on to define and study the C -restricted complex structures.

4.1 Algebraic induced complex structures

This subsection contains a recapitulation of results of [V-a].

A more general version of the following theorem was proven by A. Fujiki ([F], Theorem 4.8 (2)).

Theorem 4.1: Let M be a compact simple hyperkähler manifold and $\mathcal{R}$ be the set of induced complex structures $\mathcal{R} \cong \mathbb{C}P^1$. Let $\mathcal{R}_{alg} \subset \mathcal{R}$ be the set of all algebraic induced complex structures. Then $\mathcal{R}_{alg}$ is countable and dense in $\mathcal{R}$.

Proof: This is [V-a], Theorem 2.2. ■

In the proof of Theorem 4.1, the following important lemma was used.

Lemma 4.2:

- (i) Let $\mathcal{O} \subset H^2(M, \mathbb{R})$ be the set of all cohomology classes which are Kähler with respect to some induced complex structure. Then $\mathcal{O}$ is open in $H^2(M, \mathbb{R})$. Moreover, for all $\omega \in \mathcal{O}$, the class ω is *not* $SU(2)$ -invariant.
- (ii) Let $\eta \in H^2(M, \mathbb{R})$ be a cohomology class which is not $SU(2)$ -invariant. Then there exists a unique up to a sign induced complex structure $I \in \mathcal{R}/\{\pm 1\}$ such that η belongs to $H_I^{1,1}(M)$.

Proof: This statement is a form of [V-a], Lemma 2.3. ■

4.2 The action of $\mathfrak{so}(5)$ on the cohomology of a hyperkähler manifold

This subsection is a recollection of data from [V0] and [V2(II)].

Let M be a hyperkähler manifold. For an induced complex structure R over M , consider the Kähler form $\omega_R = (\cdot, R\cdot)$, where $(\cdot, \cdot)$ is the Riemannian form. As usually, L_R denotes the operator of exterior multiplication by ω_R , which is acting on the differential forms $A^*(M, \mathbb{C})$ over M . Consider the adjoint operator to L_R , denoted by Λ_R .

One may ask oneself, what algebra is generated by L_R and Λ_R for all induced complex structures R ? The answer was given in [V0], where the following theorem was proven.

Theorem 4.3: ([V0]) Let $M, \mathcal{H}$ be a hyperkähler manifold, and $\mathfrak{a}_{\mathcal{H}}$ be a Lie algebra generated by L_R and Λ_R for all induced complex structures R over M . Then the Lie algebra $\mathfrak{a}_{\mathcal{H}}$ is isomorphic to $\mathfrak{so}(4, 1)$. ■

The following facts about a structure of $\mathfrak{a}_{\mathcal{H}}$ were also proven in [V0].

Let I, J and K be three induced complex structures on M , such that $I \circ J = -J \circ I = K$. For an induced complex structure R , consider an operator adR on cohomology, acting on (p, q) -forms as a multiplication by $(p - q)\sqrt{-1}$. The operators adR generate a 3-dimensional Lie algebra $\mathfrak{g}_{\mathcal{H}}$, which is isomorphic to $\mathfrak{su}(2)$. This algebra coincides with the Lie algebra associated to the standard $SU(2)$ -action on $H^*(M)$. The algebra $\mathfrak{a}_{\mathcal{H}}$ contains $\mathfrak{g}_{\mathcal{H}}$ as a subalgebra, as follows:

$$[\Lambda_J, L_K] = [L_J, \Lambda_K] = ad I \quad (\text{etc}). \quad (4.1)$$

The algebra $\mathfrak{a}_{\mathcal{H}}$ is 10-dimensional. It has the following basis: $L_R, \Lambda_R, ad R$ ($R = I, J, K$) and the element $H = [L_R, \Lambda_R]$. The operator H is a standard Hodge operator; it acts on r -forms over M as multiplication by a scalar $n - r$, where $n = \dim_{\mathbb{C}} M$.

Definition 4.4: Let $\mathfrak{g}$ be a semisimple Lie algebra, V its representation and $V = \oplus V_{\alpha}$ a $\mathfrak{g}$ -invariant decomposition of V , such that for all α , V_{α} is a direct sum of isomorphic finite-dimensional representations W_{α} of V , and all W_{α} are distinct. Then the decomposition $V = \oplus V_{\alpha}$ is called **the isotypic decomposition of V** .

It is clear that for all finite-dimensional representations, isotypic decomposition always exists and is unique.

Let M be a compact hyperkähler manifold. Consider the cohomology space $H^*(M)$ equipped with the natural action of $\mathfrak{a}_{\mathcal{H}} = \mathfrak{so}(5)$. Let $H_o^* \subset H^*(M)$ be the isotypic component containing $H^0(M) \subset H^*(M)$. Using the root system written explicitly for $\mathfrak{a}_{\mathcal{H}}$ in [V0], [V3], it is easy to check that $H_o^*(M)$ is an irreducible representation of $\mathfrak{so}(5)$. Let $p : H^*(M) \longrightarrow H_o^*(M)$ be the unique $\mathfrak{so}(5)$ -invariant projection, and $i : H_o^*(M) \hookrightarrow H^*(M)$ the natural embedding.

Let M be a compact hyperkähler manifold, I an induced complex structure, and ω_I the corresponding Kähler form. Consider the **degree map** $\deg_I : H^{2p}(M) \longrightarrow \mathbb{C}$, $\eta \longmapsto \int_M \eta \wedge \omega_I^{n-p}$, where $n = \dim_{\mathbb{C}} M$.

Proposition 4.5: The space

$$H_o^*(M) \subset H^*(M)$$

is a subalgebra of $H^*(M)$, which is invariant under the $SU(2)$ -action. Moreover, for all induced complex structures I , the degree map

$$\deg_I : H^*(M) \longrightarrow \mathbb{C}$$

satisfies

$$\deg_I(\eta) = \deg_I(i(p(\eta))),$$

where $i : H_o^*(M) \hookrightarrow H^*(M)$, $p : H^*(M) \longrightarrow H_o^*(M)$ are the $\mathfrak{so}(5)$ -invariant maps defined above. And finally, the projection $p : H^*(M) \longrightarrow H_o^*(M)$ is $SU(2)$ -invariant.

Proof: The space $H_o^*(M)$ is generated from $\mathbf{1} \in H^0(M)$ by operators L_R, Λ_R . To prove that $H_o^*(M)$ is closed under multiplication, we have to show that $H_o^*(M)$ is generated (as a linear space) by expressions of type $L_{r_1} \circ L_{r_2} \circ \dots \circ \mathbf{1}$. By (4.1), the commutators of L_R, Λ_R map such expressions to linear combinations of such expressions. On the other hand, the operators Λ_R map $\mathbf{1}$ to zero. Thus, the operators Λ_R map expressions of type $L_{r_1} \circ L_{r_2} \circ \dots \circ \mathbf{1}$ to linear combinations of such expressions. This proves that $H_o^*(M)$ is closed under multiplication. The second statement of Proposition 4.5 is clear (see, e. g. [V2(II)], proof of Proposition 4.5). It remains to show that $H_o^*(M) \subset H^*(M)$ is an $SU(2)$ -invariant subspace and that $p : H^*(M) \longrightarrow H_o^*(M)$ is compatible with the $SU(2)$ -action. From (4.1), we obtain that the Lie group G_A associated with $\mathfrak{a}_{\mathcal{H}} \cong \mathfrak{so}(1, 4)$ contains $SU(2)$ acting in a standard way on $H^*(M)$. Since the map $p : H^*(M) \longrightarrow H_o^*(M)$ commutes with G_A -action, p also commutes with $SU(2)$ -action. We proved Proposition 4.5. ■

4.3 Structure of the cohomology ring

In [V3] (see also [V3-bis]), we have computed explicitly the subalgebra of cohomology of M generated by $H^2(M)$. This computation can be summed up as follows.

Theorem 4.6: ([V3], Theorem 15.2) Let M be a compact hyperkähler manifold, $H^1(M) = 0$, $\dim_{\mathbb{C}} M = 2n$, and $H_r^*(M)$ the subalgebra of cohomology of M generated by $H^2(M)$. Then

$$\begin{cases} H_r^{2i}(M) \cong S^i H^2(M) & \text{for } i \leq n, \text{ and} \\ H_r^{2i}(M) \cong S^{2n-i} H^2(M) & \text{for } i \geq n \end{cases}$$

■

Theorem 4.7: Let M be a simple hyperkähler manifold. Consider the group G generated by a union of all $SU(2)$ for all hyperkähler structures on M . Then the Lie algebra of G is isomorphic to $\mathfrak{so}(H^2(M))$, for a certain natural integer bilinear symmetric form on $H^2(M)$, called Bogomolov-Beauville form.

Proof: [V3] (see also [V3-bis]). ■

The key element in the proof of Theorem 4.6 and Theorem 4.7 is the following algebraic computation.

Theorem 4.8: Let M be a simple hyperkähler manifold, and $\mathcal{H}$ a hyperkähler structure on M . Consider the Lie subalgebra

$$\mathfrak{a}_{\mathcal{H}} \subset \text{End}(H^*(M)), \quad \mathfrak{a}_{\mathcal{H}} \cong \mathfrak{so}(1, 4),$$

associated with the hyperkähler structure (Subsection 4.2). Let

$$\mathfrak{g} \subset \text{End}(H^*(M))$$

be the Lie algebra generated by subalgebras $\mathfrak{a}_{\mathcal{H}} \subset \text{End}(H^*(M))$, for all hyperkähler structures $\mathcal{H}$ on M . Then

- (i) The algebra $\mathfrak{g}$ is naturally isomorphic to the Lie algebra $\mathfrak{so}(V \oplus \mathfrak{H})$, where V is the linear space $H^2(M, \mathbb{R})$ equipped with the Bogomolov-Beauville pairing, and $\mathfrak{H}$ is a 2-dimensional vector space with a quadratic form of signature $(1, -1)$.
- (ii) The space $H_r^*(M)$ is invariant under the action of $\mathfrak{g}$. Moreover,

$$H_r^*(M) \subset H^*(M)$$

is an isotypic¹ component of the space $H^*(M)$ considered as a representation of $\mathfrak{g}$.

¹See Definition 4.4 for the definition of isotypic decomposition.

Proof: [V3] (see also [V3-bis]). ■

As one of the consequences of Theorem 4.6, we obtain the following lemma, which will be used further on in this paper.

Lemma 4.9: Let M be a simple hyperkähler manifold, $\dim_{\mathbb{H}} M = n$, and $p : H^*(M) \longrightarrow H_o^*(M)$ the map defined in Subsection 4.2. Then, for all $x, y \in H_r^*(M)$, we have

$$p(x)p(y) = p(xy), \text{ whenever } xy \in \bigoplus_{i \leq 2n} H^i(M).$$

Proof: Let $\omega_I, \omega_J, \omega_K, x_1, \dots, x_n$ be an orthonormal basis in $H^2(M)$. Clearly, the vectors $x_1, \dots, x_k$ are $SU(2)$ -invariant. Therefore, these vectors are highest vectors of the corresponding $\mathfrak{a}_{\mathcal{H}}$ -representations, with respect to the root system and Cartan subalgebra for $\mathfrak{a}_{\mathcal{H}}$ which is written in [V0] or [V3]. We obtain that the monomials

$$P_{k_1, k_2, k_3, \{n_i\}} = \omega_I^{k_1} \omega_J^{k_2} \omega_K^{k_3} \prod x_i^{n_i}, \sum n_i = N, \quad P_{k_1, k_2, k_3, \{n_i\}} \in \bigoplus_{i \leq 2n} H^i(M)$$

belong to the different isotypical components for different N 's. By Theorem 4.6, a product of two such monomials $P_{k_1, k_2, k_3, \{n_i\}}$ and $P_{k'_1, k'_2, k'_3, \{n'_i\}}$ is equal to $P_{k_1+k'_1, k_2+k'_2, k_3+k'_3, \{n_i+n'_i\}}$, assuming that

$$P_{k_1, k_2, k_3, \{n_i\}} P_{k'_1, k'_2, k'_3, \{n'_i\}} \in \bigoplus_{i \leq 2n} H^i(M).$$

Thus, the isotypical decomposition associated with the $\mathfrak{a}_{\mathcal{H}}$ -action is compatible with multiplicative structure on $H^*(M)$, for low-dimensional cycles. This implies Lemma 4.9. ■

We shall use the following corollary of Lemma 4.9.

Corollary 4.10: Let M be a simple hyperkähler manifold, $\dim_{\mathbb{H}} M > 1$, and $\omega_1, \omega_2 \in H^2(M)$ cohomology classes which are $SU(2)$ -invariant. Then, for all induced complex structures I , we have $\deg_I(\omega_1 \omega_2) = 0$.

Proof: By definition, the classes ω_1, ω_2 satisfy $\omega_i \in \ker p$. By Lemma 4.9, we have $\omega_1 \omega_2 \in \ker p$. By Proposition 4.5, $\deg_I \omega_1 \omega_2 = 0$. ■

Let ω be a rational Kähler form. The corresponding $\mathfrak{sl}(2)$ -action on $H^*(M)$ is clearly compatible with the rational structure on $H^*(M)$. It is easy

to see (using, for instance, Lemma 4.2) that $\mathfrak{g}$ is generated by $\mathfrak{sl}(2)$ -triples associated with rational Kähler forms ω . Therefore, the action of $\mathfrak{g}$ on $H^*(M)$ is compatible with the rational structure on $H^*(M)$. Using the isotypic decomposition, we define a natural $\mathfrak{g}$ -invariant map $r : H^*(M) \longrightarrow H_r^*(M)$. Further on, we shall use the following properties of this map.

Claim 4.11:

- (i) The map $r : H^*(M) \longrightarrow H_r^*(M)$ is compatible with the rational structure on $H^*(M)$.
- (ii) For every $x \in \ker r$, and every hyperkähler structure $\mathcal{H}$, the corresponding map $p : H^*(M) \longrightarrow H_o^*(M)$ satisfies $p(x) = 0$.
- (iii) For every $x \in \ker r$, every hyperkähler structure $\mathcal{H}$, and every induced complex structure I on M , we have $\deg_I x = 0$.

Proof: Claim 4.11 (i) is clear, because the action of $\mathfrak{g}$ on $H^*(M)$ is compatible with the rational structure on $H^*(M)$. To prove Claim 4.11 (ii), we notice that the space $H_r^*(M)$ is generated from $H^0(M)$ by the action of $\mathfrak{g}$, and $H_o^*(M)$ is generated from $H^0(M)$ by the action of $\mathfrak{a}_{\mathcal{H}}$. Since $\mathfrak{a}_{\mathcal{H}}$ is by definition a subalgebra in $\mathfrak{g}$, we have $H_o^*(M) \subset H_r^*(M)$. The isotypic projection $r : H^*(M) \longrightarrow H_r^*(M)$ is by definition compatible with the $\mathfrak{g}$ -action. Since $\mathfrak{a}_{\mathcal{H}} \subset \mathfrak{g}$, the map r is also compatible with the $\mathfrak{a}_{\mathcal{H}}$ -action. Therefore, $\ker r \subset \ker p$. Claim 4.11 (iii) is implied by Claim 4.11 (ii) and Proposition 4.5. ■

Let x_i be an basis in $H^2(M, \mathbb{Q})$ which is rational and orthonormal with respect to Bogomolov-Beauville pairing, $(x_i, x_i)_{\mathcal{B}} = \varepsilon_i = \pm 1$. Consider the cohomology class $\theta' := \varepsilon_i x_i^2 \in H^4(M, \mathbb{Q})$. Let $\theta \in H^4(M, \mathbb{Z})$ be a non-zero integer cohomology class which is proportional to θ' . From results of [V3] (see also [V3-bis]), the following proposition can be easily deduced.

Proposition 4.12: The cohomology class $\theta \in H^4(M, \mathbb{Z})$ is $SU(2)$ -invariant for all hyperkähler structures on M . Moreover, for a generic hyperkähler structure, the group of $SU(2)$ -invariant integer classes $\alpha \in H_r^4(M)$ has rank one, where $H_r^*(M)$ is the subalgebra of cohomology generated by $H^2(M)$.

Proof: Clearly, if an integer class α is $SU(2)$ -invariant for a generic hyperkähler structure, then α is G -invariant, where G is the group de-

defined in Theorem 4.7. On the other hand, $H_r^4(M) \cong S^2(H^2(M))$, as follows from Theorem 4.6. Clearly, the vector $\theta \in H_r^4(M) \cong S^2(H^2(M))$ is $\mathfrak{so}(H^2(M))$ -invariant. Moreover, the space of $\mathfrak{so}(H^2(M))$ -invariant vectors in $S^2(H^2(M))$ is one-dimensional. Finally, from an explicit computation of G it follows that G acts on $H^4(M)$ as $SO(H^2(M))$, and thus, the Lie algebra invariants coincide with invariants of G . We found that the space of G -invariants in $H_r^4(M)$ is one-dimensional and generated by θ . This proves Proposition 4.12. ■

Remark 4.13: It is clear how to generalize Proposition 4.12 from dimension 4 to all dimensions. The space $H_r^{2d}(M)^G$ of G -invariants in $H_r^{2d}(M)$ is 1-dimensional for d even and zero-dimensional for d odd.

4.4 Cohomology classes of CA-type

Let M be a compact hyperkähler manifold, and I an induced complex structure. All cohomology classes which appear as fundamental classes of complex subvarieties of (M, I) satisfy certain properties. Classes satisfying these properties are called classes of CA-type, from Complex Analytic. Here is the definition of CA-type.

Definition 4.14: Let $\eta \in H_I^{2,2}(M) \cap H^4(M, \mathbb{Z})$ be an integer (2,2)-class. Assume that for all induced complex structures J , satisfying $I \circ J = -J \circ I$, we have $\deg_I(\eta) \geq \deg_J(\eta)$, and the equality is reached only if η is $SU(2)$ -invariant. Assume, moreover, that $\deg_I(\eta) \geq |\deg_J(\eta)|$. Then η is called a **class of CA-type**.

Theorem 4.15: Let M be a simple hyperkähler manifold, of dimension $\dim_{\mathbb{H}} M > 1$, I an induced complex structure, and $\eta \in H_I^{2,2}(M) \cap H^4(M, \mathbb{Z})$ an integer (2,2)-class. Assume that one of the following conditions holds.

- (i) There exists a complex subvariety $X \subset (M, I)$ such that η is the fundamental class of X
- (ii) There exists a stable coherent torsion-free sheaf F over (M, I) , such that the first Chern class of F is zero, and $\eta = c_2(F)$.

Then η is of CA-type.

Proof: Theorem 4.15 (i) is a direct consequence of Wirtinger's inequality (Proposition 2.11). It remains to prove Theorem 4.15 (ii).

We assume, temporarily, that F is reflexive. By Corollary 3.24, we have

$$\deg_I(2c_2(F) - \frac{r-1}{r}c_1(F)^2) \geq \left| \deg_J(2c_2(F) - \frac{r-1}{r}c_1(F)^2) \right|, \quad (4.2)$$

and the equality happens only if F is hyperholomorphic. Since $c_1(F)$ is $SU(2)$ -invariant, we have $\deg_I(c_1(F)^2) = \deg_J(c_1(F)^2) = 0$ (Corollary 4.10). Thus, (4.2) implies that

$$\deg_I 2c_2(F) \geq |\deg_J 2c_2(F)|$$

and the inequality is strict unless F is hyperholomorphic, in which case, the class $c_2(F)$ is $SU(2)$ -invariant by definition. We have proven Theorem 4.15 (ii) for the case of reflexive F .

For F not necessary reflexive sheaf, we have shown in the proof of Claim 3.13 that

$$c_2(F) = c_2(F^{**}) + \sum n_i [X_i],$$

where n_i are positive integers, and $[X_i]$ are the fundamental classes of irreducible components of support of the sheaf F^{**}/F . Therefore, the class $c_2(F)$ is a sum of classes of CA-type. Clearly, a sum of cohomology classes of CA-type is again a class of CA-type. This proves Theorem 4.15. ■

5 C -restricted complex structures on hyperkähler manifolds

5.1 Existence of C -restricted complex structures

We assume from now till the end of this section that the hyperkähler manifold M is simple (Definition 2.7). This assumption can be avoided, but it simplifies notation.

We assume from now till the end of this section that the hyperkähler manifold M is compact of real dimension $\dim_{\mathbb{R}} M \geq 8$, i. e. $\dim_{\mathbb{H}} M \geq 2$. This assumption is absolutely necessary. The case of hyperkähler surfaces with $\dim_{\mathbb{H}} M = 1$ (torus and K3 surface) is trivial and for our purposes not interesting. It is not difficult to extend our definitions and results to the case of a compact hyperkähler manifold which is a product of simple hyperkähler manifolds with $\dim_{\mathbb{H}} M \geq 2$.

Definition 5.1: Let M be a compact hyperkähler manifold, and I an induced complex structure. As usually, we denote by $\deg_I : H^{2p}(M) \rightarrow \mathbb{C}$ the associated degree map, and by $H^*(M) = \oplus H_I^{p,q}(M)$ the Hodge decomposition. Assume that I is algebraic. Let C be a positive real number. We say that the induced complex structure I is **C -restricted** if the following conditions hold.

- (i) For all non- $SU(2)$ -invariant cohomology classes $\eta \in H_I^{1,1}(M) \cap H^2(M, \mathbb{Z})$, we have $|\deg_I(\eta)| > C$.
- (ii) Let $\eta \in H_I^{2,2}(M)$ be a cohomology class of CA-type which is not $SU(2)$ -invariant. Then $|\deg_I(\eta)| > C$.

The heuristic (completely informal) meaning of this definition is the following. The degree map plays the role of the metric on the cohomology. Cohomology classes with small degrees are “small”, the rest is “big”. Under reasonably strong assumptions, there are only finitely many “small” integer classes, and the rest is “big”. For each non- $SU(2)$ -invariant cohomology class η there exists at most two induced complex structures for which η is of type (p, p) . Thus, for most induced complex structures, all non- $SU(2)$ -invariant integer (p, p) classes are “big”. Intuitively, the C -restriction means that all non- $SU(2)$ -invariant integer $(1, 1)$ and $(2, 2)$ -cohomology classes are “big”. This definition is needed for the study of first and second Chern classes of sheaves. The following property of C -restricted complex structures is used (see Theorem 4.15): for every subvariety $X \subset (M, I)$ of complex codimension 2, either X is trianalytic or $\deg_I(X) > C$.

Definition 5.2: Let M be a compact manifold, and $\mathcal{H}$ a hyperkähler structure on M . We say that $\mathcal{H}$ **admits C -restricted complex structures** if for all $C > 0$, the set of all C -restricted algebraic complex structures is dense in the set $R_{\mathcal{H}} = \mathbb{C}P^1$ of all induced complex structures.

Proposition 5.3: Let M be a compact simple hyperkähler manifold, $\dim_{\mathbb{H}}(M) > 1$, and $r : H^4(M) \rightarrow H_r^4(M)$ be the map defined in Claim 4.11. Assume that for all algebraic induced complex structures I , the group $H_I^{1,1}(M) \cap H^2(M, \mathbb{Z})$ has rank one, and the group

$$H_I^{2,2}(M) \cap H^4(M, \mathbb{Z}) / (\ker r)$$

has rank 2. Then M admits C -restricted complex structures.

Proof: The proof of Proposition 5.3 takes the rest of this section.

Denote by $\mathcal{R}$ the set $\mathcal{R} \cong \mathbb{C}P^1$ of all induced complex structures on M . Consider the set $\mathcal{R}/\{\pm 1\}$ of induced complex structures up to a sign (Lemma 4.2). Let $\alpha \in H^2(M)$ be a cohomology class which is not $SU(2)$ -invariant. According to Lemma 4.2, there exists a unique element $c(\alpha) \in \mathcal{R}/\{\pm 1\}$ such that $\alpha \in H_{c(\alpha)}^{1,1}(M)$. This defines a map

$$c : (H^2(M, \mathbb{R}) \setminus H_{inv}^2(M)) \longrightarrow \mathcal{R}/\{\pm 1\},$$

where $H_{inv}^2(M) \subset H^2(M)$ is the set of all $SU(2)$ -invariant cohomology classes. For induced complex structures I and $-I$, and $\eta \in H^{2p}(M)$, the degree maps satisfy

$$\deg_I(\eta) = (-1)^p \deg_{-I}(\eta). \quad (5.1)$$

Thus, the number $|\deg_I(\eta)|$ is independent from the sign of I .

Let $\eta \in H^*(M, \mathbb{Z})$ be a cohomology class. The **largest divisor** of η is the biggest positive integer number k such that the cohomology class ηk is also integer.

Let $\alpha \in H^2(M, \mathbb{Z})$ be an integer cohomology class, which is not $SU(2)$ -invariant, k its largest divisor and $\tilde{\alpha} := \alpha k$ the corresponding integer class. Denote by $\widetilde{\deg}(\alpha)$ the number

$$\widetilde{\deg}(\alpha) := \left| \deg_{c(\alpha)}(\tilde{\alpha}) \right|.$$

The induced complex structure $c(\alpha)$ is defined up to a sign, but from (5.1) it is clear that $\widetilde{\deg}(\alpha)$ is independent from the choice of a sign.

Lemma 5.4: Let M be a compact hyperkähler manifold, and I be an algebraic induced complex structure, such that the group $H_I^{1,1}(M) \cap H^2(M, \mathbb{Z})$ has rank one, and the group $H_I^{2,2}(M) \cap H^4(M, \mathbb{Z})/(\ker r)$ has rank 2. Denote by α the generator of $H_I^{1,1}(M) \cap H^2(M, \mathbb{Z})$. Since the class α is proportional to a Kähler form, α is not $SU(2)$ -invariant (Lemma 4.2, (i)). Let $d := \widetilde{\deg} \alpha$. Then, there exists a positive real constant A depending on volume of M , its topology and its dimension, such that I is $d \cdot A$ -restricted.

Proof: This lemma is a trivial calculation based on results of [V3] (see also [V3-bis] and Subsection 4.3).

Since $H_I^{1,1}(M) \cap H^2(M, \mathbb{Z})$ has rank one, for all $\eta \in H_I^{1,1}(M) \cap H^2(M, \mathbb{Z})$, $\eta \neq 0$, we have $|\deg_I \eta| \geq d$. This proves the first condition of Definition 5.1.

Let θ be the $SU(2)$ -invariant integer cycle $\theta \in H^4(M)$ defined in Proposition 4.12. By Lemma 2.6, $\theta \in H_I^{2,2}(M)$. Consider $\alpha^2 \in H_I^{2,2}(M)$, where α is the generator of $H_I^{1,1}(M) \cap H^2(M, \mathbb{Z})$.

Sublemma 5.5: Let J be an induced complex structure, $J \circ I = -J \circ I$, and $\deg_I, \deg_J$ the degree maps associated with I, J . Then

$$\deg_I \alpha^2 > 0, \deg_J \alpha^2 = 0, \deg_I \theta = \deg_J \theta > 0.$$

Proof: Since α is a Kähler class with respect to I , we have $\deg_I \alpha^2 > 0$. Since the cohomology class θ is $SU(2)$ -invariant, and $SU(2)$ acts transitively on the set of induced complex structures, we have $\deg_I \theta = \deg_J \theta$. It remains to show that $\deg_J \alpha^2 = 0$ and $\deg_J \theta > 0$. The manifold M is by our assumptions simple; thus, $\dim H^{2,0}(M) = 1$ ([Bes]). Therefore, in the natural $SU(2)$ -invariant decomposition

$$H^2(M) = H_{inv}^2(M) \oplus H_+^2(M), \quad (5.2)$$

we have $\dim H_+^2(M) = 3$. In particular, the intersection $H_+^2(M) \cap H_I^{1,1}(M)$ is 1-dimensional. Consider the decomposition of α , associated with (5.2): $\alpha = \alpha_+ + \alpha_{inv}$. Since α is of type $(1,1)$ with respect to I , the class α_+ is proportional to the Kähler class ω_I , with positive coefficient. A similar argument leads to the following decomposition for θ :

$$\theta = \omega_I^2 + \omega_J^2 + \omega_K^2 + \sum x_i^2,$$

where $K = I \circ J$ is an induced complex structure, and the classes x_i belong to $H_{inv}^2(M)$. From Corollary 4.10, we obtain that the classes x_i^2 satisfy $\deg_I(x_i^2) = 0$ (here we use $\dim_{\mathbb{H}}(M) > 1$). Thus,

$$\deg_I(\theta) = \deg_I(\omega_I^2 + \omega_J^2 + \omega_K^2) = \deg_I(\omega_I^2) > 0.$$

Similarly one checks that

$$\deg_J(\alpha^2) = \deg_J((\alpha_+ + \alpha_{inv})^2) = \deg_J(\alpha_+^2) = \deg_J(c^2 \omega_I) = 0.$$

This proves Sublemma 5.5. ■

Return to the proof of Lemma 5.4. Since $\deg_I \alpha^2 \neq \deg_J \alpha^2$, the class α^2 is *not* $SU(2)$ -invariant. Since θ is $SU(2)$ -invariant, θ is not collinear with α^2 . The degrees $\deg_I$ of θ and α^2 are non-zero; we have $\deg_I(\theta) = \deg_J(\theta)$, $\deg_I(\alpha^2) \neq \deg_J(\alpha^2)$ for J an induced complex structure, $J \neq \pm I$. By Proposition 4.5, no non-trivial linear combination of θ , α^2 belongs to $\ker p$. By Claim 4.11 (ii), the classes θ , α^2 generate a 2-dimensional subspace in $H^4(M, \mathbb{Q})/\ker r$.

By assumptions of Lemma 5.4, the group $H_I^{2,2}(M) \cap H^4(M, \mathbb{Z})/(\ker r)$ has rank 2. Therefore ω and α^2 generate the space

$$H_I^{2,2}(M) \cap H^4(M, \mathbb{Q})/(\ker r).$$

To prove Lemma 5.4 it suffices to show that for all integer classes

$$\beta = a\alpha^2 + b\theta, \quad a \in \mathbb{Q} \setminus 0, \quad \deg_I \beta \geq \deg_J \beta,$$

we have $|\deg_I \beta| > A \cdot d$, for a constant A depending only on volume, topology and dimension of M . Since $\deg_I \beta \geq |\deg_J \beta|$, and $\deg_J \alpha^2 = 0$ (Sublemma 5.5), we have

$$\deg_I(a\alpha^2 + b\theta) \geq |\deg_J b\theta|.$$

Therefore, either a and b have the same sign, or $\deg_I(a\alpha^2) > 2\deg_I(b\theta)$. In both cases,

$$|\deg_I \beta| \geq \frac{1}{2} \deg_I(a\alpha^2). \quad (5.3)$$

Let $x \in \mathbb{Q}^{>0}$ be the smallest positive rational value of a for which there exists an integer class $\beta = a\alpha^2 + b\theta$. We have an integer lattice L_1 in $H_r^4(M)$ provided by the products of integer classes; the integer lattice $L_2 \supset L_1$ provided by integer cycles might be different from that one. Clearly, x is greater than determinant $\det(L_1/L_2)$ of L_1 over L_2 , and this determinant is determined by the topology of M .

Form the definition of x and (5.3), we have $|\deg_I \beta| > x^2 \deg_I(\alpha^2)$. On the other hand, $\deg_I(\alpha^2) > C \deg_I(\alpha)$, where C is a constant depending on volume and dimension of M . Setting $A := x^2 \cdot C$, we obtain $|\deg_I \beta| > x^2 \cdot C \cdot d$. This proves Lemma 5.4. ■

Consider the maps

$$\widetilde{\deg}: H^2(M, \mathbb{Z}) \setminus H_{inv}^2(M) \longrightarrow \mathbb{R},$$

$$c : H^2(M) \setminus H_{inv}^2(M) \longrightarrow \mathcal{R}/\{\pm 1\}$$

introduced in the beginning of the proof of Proposition 5.3.

Lemma 5.6: In assumptions of Proposition 5.3, let

$$\mathcal{O} \subset H^2(M, \mathbb{R}) \setminus H_{inv}^2(M)$$

be an open subset of $H^2(M, \mathbb{R})$, such that for all $x \in \mathcal{O}$, $k \in \mathbb{R}^{>0}$, we have $k \cdot x \in \mathcal{O}$. Assume that $\mathcal{O}$ contains the Kähler class ω_I for all induced complex structures $I \in \mathcal{R}$. For a positive number $C \in \mathbb{R}^{>0}$, consider the set $X_C \subset \mathcal{O}$

$$X_C := \left\{ \alpha \in \mathcal{O} \cap H^2(M, \mathbb{Z}) \mid \widetilde{\deg}(\alpha) \geq C \right\}.$$

Then $c(X_C)$ is dense in $\mathcal{R}/\{\pm 1\}$ for all $C \in \mathbb{R}^{>0}$.

Proof: The map $\widetilde{\deg}$ can be expressed in the following way. We call an integer cohomology class $\alpha \in H^2(M, \mathbb{Z})$ **indivisible** if its largest divisor is 1, that is, there are no integer classes α' , and numbers $k \in \mathbb{Z}$, $k > 1$, such that $\alpha = k\alpha'$.

Sublemma 5.7: Let $\alpha \in H^2(M)$ be a non- $SU(2)$ -invariant cohomology class and $\alpha = \alpha_{inv} + \alpha_+$ be a decomposition associated with (5.2). Assume that α is indivisible. Then

$$\widetilde{\deg}(\alpha) = C \sqrt{((\alpha_+, \alpha_+)_{\mathcal{B}})}, \quad (5.4)$$

where $(\cdot, \cdot)_{\mathcal{B}}$ is the Bogomolov-Beauville pairing on $H^2(M)$ ([V3-bis]; see also Theorem 4.7), and C a constant depending on $\dim M$, $\text{Vol } M$.

Proof: By Proposition 4.5,

$$\deg_I(\alpha) = \deg_I(\alpha_+)$$

(clearly, $p(\alpha) = \alpha_+$). By definition of $(\cdot, \cdot)_{\mathcal{B}}$, we have

$$\deg_I(\alpha_+) = (\alpha_+, \omega_{c(\alpha)})_{\mathcal{B}}$$

On the other hand, α_+ is collinear with $\omega_{c(\alpha)}$ by definition of the map c . Now (5.4) follows trivially from routine properties of bilinear forms. ■

Let I be an induced complex structure such that the cohomology class ω_I is irrational: $\omega_I \notin H^2(M, \mathbb{Q})$.

To prove Lemma 5.6, we have to produce a sequence $x_i \in \mathcal{O} \cap H^2(M, \mathbb{Z})$ such that

- (i) $c(x_i)$ converges to I ,
 - (ii) and $\lim \widetilde{\deg}(x_i) = \infty$.
- (5.5)

We introduce a metric $(\cdot, \cdot)_{\mathcal{H}}$ on $H^2(M, \mathbb{R})$,

$$(\alpha, \beta)_{\mathcal{H}} := (\alpha_+, \beta_+)_{\mathcal{B}} - (\alpha_{inv}, \beta_{inv})_{\mathcal{B}}.$$

It is easy to check that $(\cdot, \cdot)_{\mathcal{H}}$ is positive definite ([V3]). For every ε , there exists a rational class $\omega_{\varepsilon} \in H^2(M, \mathbb{Q})$ which approximates ω_I with precision

$$(\omega_{\varepsilon} - \omega_I, \omega_{\varepsilon} - \omega_I)_{\mathcal{H}} < \varepsilon.$$

Since $\mathcal{O}$ is open and contains ω_I , we may assume that ω_{ε} belongs to $\mathcal{O}$. Take a sequence ε_i converging to 0, and let $\tilde{x}_i := \omega_{\varepsilon_i}$ be the corresponding sequence of rational cohomology cycles. Let $x_i := \lambda_i \tilde{x}_i$ be the minimal positive integer such that $x_i \in H^2(M, \mathbb{Z})$. We are going to show that the sequence x_i satisfies the conditions of (5.5). First of all, $\tilde{x}_i$ converges to ω_I , and the map

$$c: H^2(M) \setminus H_{inv}^2(M) \longrightarrow \mathcal{R}/\{\pm 1\}$$

is continuous. Therefore, $\lim c(\tilde{x}_i) = c(\omega_I) = I$. By construction of c , c satisfies $c(x) = c(\lambda x)$, and thus, $c(x_i) = c(\tilde{x}_i)$. This proves the condition (i) of (5.5). On the other hand, since ω_I is irrational, the sequence λ_i goes to infinity. Therefore,

$$\lim (x_i, x_i)_{\mathcal{H}} = \infty.$$

It remains to compare $(x_i, x_i)_{\mathcal{H}}$ with $\widetilde{\deg} x_i$. By (5.4),

$$\widetilde{\deg} x_i = \sqrt{((x_i)_+, (x_i)_+)_{\mathcal{B}}}.$$

On the other hand, since $(x_i)_+ \in H_+^2(M)$, we have

$$((x_i)_+, (x_i)_+)_{\mathcal{B}} = ((x_i)_+, (x_i)_+)_{\mathcal{H}}.$$

To prove (5.5) (ii), it remains to show that

$$\lim ((x_i)_+, (x_i)_+)_{\mathcal{H}} = \lim (x_i, x_i)_{\mathcal{H}}.$$

Since the cohomology class $\tilde{x}_i \in H^2(M, \mathbb{Q})$ ε -approximates ω_I , and ω_I belongs to $H_+^2(M)$, we have

$$(\tilde{x}_i - (\tilde{x}_i)_+, \tilde{x}_i - (\tilde{x}_i)_+)_{\mathcal{H}} < \varepsilon_i.$$

Therefore,

$$(x_i - (x_i)_+, x_i - (x_i)_+)_{\mathcal{H}} < \lambda_i \varepsilon_i. \quad (5.6)$$

On the other hand, for i sufficiently big, the cohomology class $\tilde{x}_i$ approaches ω_I , and

$$(x_i, x_i)_{\mathcal{H}} > \frac{1}{2} \lambda_i (\omega_I, \omega_I)_{\mathcal{H}} \quad (5.7)$$

Comparing (5.6) and (5.7) and using the distance property for the distance given by $\sqrt{(\cdot, \cdot)_{\mathcal{H}}}$, we find that

$$\sqrt{(x_i)_+, (x_i)_+} > \sqrt{\frac{1}{2} \lambda_i (\omega_I, \omega_I)_{\mathcal{H}}} - \sqrt{\lambda_i \varepsilon_i} = \sqrt{\lambda_i} \cdot \left(\sqrt{\frac{1}{2} (\omega_I, \omega_I)_{\mathcal{H}}} - \sqrt{\varepsilon_i} \right). \quad (5.8)$$

Since ε_i converges to 0 and λ_i converges to infinity, the right hand side of (5.8) converges to infinity. On the other hand, by (5.4) the left hand side of (5.8) is equal constant times $\deg x_i$, so $\lim \deg x_i = \infty$. This proves the second condition of (5.5). Lemma 5.6 is proven. ■

We use Lemma 5.4 and Lemma 5.6 in order to finish the proof of Proposition 5.3.

Let M be a compact hyperkähler manifold, and $\mathcal{O} \subset H^2(M, \mathbb{R})$ be the set of all Kähler classes for the Kähler metrics compatible with one of induced complex structures. By Lemma 4.2, $\mathcal{O}$ is open in $H^2(M, \mathbb{R})$. Applying Lemma 5.6 to $\mathcal{O}$, we obtain the following. In assumptions of Proposition 5.3, let $Y_C \subset \mathcal{R}$ be the set of all algebraic induced complex structures I with $\deg \alpha > C$, where α is a rational Kähler class, $\alpha \in H^{1,1}(M) \cap H^2(M, \mathbb{Z})$. Then Y_C is dense in $\mathcal{R}$. Now, Lemma 5.4, implies that for all $I \in Y_C$, the induced complex structure I is $A \cdot C$ -restricted, where A is the universal constant of Lemma 5.4. Thus, for all C the set of C -restricted induced complex structures is dense in $\mathcal{R}$. This proves that M admits C -restricted complex structures. We finished the proof of Proposition 5.3. ■

5.2 Hyperkähler structures admitting C -restricted complex structures

Let M be a compact complex manifold admitting a hyperkähler structure $\mathcal{H}$. Assume that $(M, \mathcal{H})$ is a simple hyperkähler manifold of dimension $\dim_{\mathbb{H}} M > 1$. The following definition of (coarse, marked) moduli space for complex and hyperkähler structures on M is standard.

Definition 5.8: Let M_{C^∞} be the M considered as a differential manifold, $\widetilde{Comp}$ be the set of all integrable complex structures, and $\widetilde{Hyp}$ be the set of all hyperkähler structures on M_{C^∞} . The set $\widetilde{Hyp}$ is equipped with a natural topology. Let $\widetilde{Hyp}^0$ be a connected component of $\widetilde{Hyp}$ containing $\mathcal{H}$ and $\widetilde{Comp}^0$ be a set of all complex structures $I \in \widetilde{Comp}$ which are compatible with some hyperkähler structure $\mathcal{H}_1 \in \widetilde{Hyp}^0$. Let $Diff$ be the group of diffeomorphisms of M which act trivially on the cohomology. The coarse, marked moduli Hyp of hyperkähler structures on M is the quotient $Hyp := \widetilde{Hyp}^0 / Diff$ equipped with a natural topology. The coarse, marked moduli $Comp$ of complex structures on M is defined as $Comp := \widetilde{Comp}^0 / Diff$. For a detailed discussion of various aspects of this definition, see [V3].

Consider the variety

$$X \subset \mathbb{P}H^2(M, \mathbb{C}),$$

consisting of all lines $l \in \mathbb{P}H^2(M, \mathbb{C})$ which are isotropic with respect to the Bogomolov-Beauville's pairing:

$$X := \{l \in H^2(M, \mathbb{C}) \mid (l, l)_{\mathcal{B}} = 0\}.$$

Since M is simple, $\dim H^{2,0}(M, I) = 1$ for all induced complex structures. Let $P_c : Comp \rightarrow \mathbb{P}H^2(M, \mathbb{C})$ map I to the line $H_I^{2,0}(M) \subset H^2(M, \mathbb{C})$. The map P_c is called **the period map**. It is well known that $Comp$ is equipped with a natural complex structure. From general properties of the period map it follows that P_c is compatible with this complex structure. Clearly from the definition of Bogomolov-Beauville's form, $P_c(I) \in X$ for all induced complex structures $I \in Comp$ (see [Bea] for details).

Theorem 5.9: [Bes] (Bogomolov) The complex analytic map

$$P_c : Comp \rightarrow X$$

is locally an étale covering.¹

■

It is possible to formulate a similar statement about hyperkähler structures. For a hyperkähler structure $\mathcal{H}$, consider the set $\mathcal{R}_{\mathcal{H}} \subset \text{Comp}$ of all induced complex structures associated with this hyperkähler structure. The subset $\mathcal{R}_{\mathcal{H}} \subset \text{Comp}$ is a complex analytic subvariety, which is isomorphic to $\mathbb{C}P^1$. Let $S := P_c(\mathcal{R}_{\mathcal{H}})$ be the corresponding projective line in X , and $\overline{L}(X)$ be the space of smooth deformations of S in X . The points of $L(X)$ correspond to smooth rational curves of degree 2 in $\mathbb{P}H^2(M, \mathbb{C})$. For every such curve s , there exists a unique 3-dimensional plane $L(s) \subset H^2(M, \mathbb{C})$, such that s is contained in $\mathbb{P}L$. Let Gr be the Grassmanian manifold of all 3-dimensional planes in $H^2(M, \mathbb{C})$ and $Gr_0 \subset Gr$ the set of all planes $L \in Gr$ such that the restriction of the Bogomolov-Beauville form to L is non-degenerate. Let $L(X) \subset \overline{L}(X)$ be the space of all rational curves $s \in \overline{L}(X)$ such that the restriction of the Bogomolov-Beauville form to $L(s)$ is non-degenerate: $L(s) \in Gr_0$. The correspondence $s \longrightarrow L(s)$ gives a map $\kappa : L(X) \longrightarrow Gr_0$.

Lemma 5.10: The map $\kappa : L(X) \longrightarrow Gr_0$ is an isomorphism of complex varieties.

Proof: For every plane $L \in Gr_0$, consider the set $s(L)$ of all isotropic lines $l \in L$, that is, lines satisfying $(l, l)_{\mathcal{B}} = 0$. Since $(\cdot, \cdot)_{\mathcal{B}}|_L$ is non-degenerate, the set $s(L)$ is a rational curve in $\mathbb{P}L$. Clearly, this curve has degree 2. Therefore, $s(L)$ belongs to $X(L)$. The map $L \longrightarrow s(L)$ is inverse to κ . ■

Consider the standard anticomplex involution

$$\iota : H^2(M, \mathbb{C}) \longrightarrow H^2(M, \mathbb{C}), \quad \eta \longrightarrow \overline{\eta}.$$

Clearly, ι is compatible with the Bogomolov-Beauville form. Therefore, ι acts on $L(X)$ as an anticomplex involution. Let $L(X)_{\iota} \subset L(X)$ be the set of all $S \in L(X)$ fixed by ι .

Every hyperkähler structure

$$\mathcal{H} \in \text{Hyp}$$

¹ The space Comp is smooth, as follows from Theorem 5.9. This space is, however, in most cases not separable ([H]). The space Hyp has no natural complex structures, and can be odd-dimensional.

gives a rational curve $\mathcal{R}_{\mathcal{H}} \subset \text{Comp}$ with points corresponding all induced complex structures. Let $P_h(\mathcal{H}) \subset X$ be the line $P_c(\mathcal{R}_{\mathcal{H}})$. Clearly from the definition, $P_h(\mathcal{H})$ belongs to $L(X)_\iota$. We have constructed a map $P_h : \text{Hyp} \longrightarrow L(X)_\iota$. Let $L(\text{Comp})$ be the space of deformations if $\mathcal{R}_{\mathcal{H}}$ in Comp . Denote by

$$\gamma : \text{Hyp} \longrightarrow L(\text{Comp})$$

the map $\mathcal{H} \longrightarrow \mathcal{R}_{\mathcal{H}}$. The following result gives a hyperkähler analogue of Bogomolov's theorem (Theorem 5.9).

Theorem 5.11: The map $\gamma : \text{Hyp} \longrightarrow L(\text{Comp})$ is an embedding. The map $P_h : \text{Hyp} \longrightarrow L(X)_\iota$ is locally a covering.

Proof: The first claim is an immediate consequence of Calabi-Yau Theorem (Theorem 2.4). Now, Theorem 5.11 follows from the Bogomolov's theorem (Theorem 5.9) and dimension count. ■

Let $I \in \text{Comp}$ be a complex structure on M . Consider the groups

$$H_h^2(M, I) := H^{1,1}(M, I) \cap H^2(M, \mathbb{Z})$$

and

$$H_h^2(M, I) := H_r^{2,2}(M, I) \cap H^4(M, \mathbb{Z}).$$

For a general I , $H_h^2(M, I) = 0$ and $H_h^4(M, I) = \mathbb{Z}$ as follows from Proposition 4.12. Therefore, the set of all I with $\text{rk } H_h^2(M, I) = 1$, $\text{rk } H_h^4(M, I) = 2$ is a union of countably many subvarieties of codimension 1 in Comp . Similarly, the set $V \subset \text{Comp}$ of all I with $\text{rk } H_h^2(M, I) > 1$, $\text{rk } H_h^4(M, I) > 2$ is a union of countably many subvarieties of codimension more than 1. Together with Theorem 5.11, this implies the following.

Claim 5.12: Let $U \subset \text{Hyp}$ be the set of all $\mathcal{H} \in \text{Hyp}$ such that $\mathcal{R}_{\mathcal{H}}$ does not intersect V . Then U is dense in Hyp .

Proof: Consider a natural involution i of Comp which is compatible with the involution $\iota : X \longrightarrow X$ under the period map $P_c : \text{Comp} \longrightarrow X$. This involution maps the complex structure I to $-I$.

By Theorem 5.11, Hyp is identified with an open subset in the set $L(X)_\iota$ of real points of $L(\text{Comp})$. (5.9)

Let $L_U \subset L(\text{Comp})$ be the set of all lines which do not intersect V . Since V is a union of subvarieties of codimension at least 2, a general rational line

$l \in L(Comp)$ does not intersect V . Therefore, L_U is dense in $L(Comp)$. Thus, the set of real points of L_U is dense $L(X)_\iota$. Using the identification (5.9), we obtain the statement of Claim 5.12. ■

Claim 5.12 together with Proposition 5.3 imply the following theorem.

Theorem 5.13: Let M be a compact simple hyperkähler manifold, $\dim_{\mathbb{H}} M > 1$, and Hyp its coarse marked moduli of hyperkähler structures. Let $U \subset Hyp$ be the set of all hyperkähler structures which admit C -restricted complex structures (Definition 5.2). Then U is dense in Hyp . ■

5.3 Deformations of coherent sheaves over manifolds with C -restricted complex structures

The following theorem shows that a semistable deformation of a hyperholomorphic sheaf on (M, I) is again hyperholomorphic, provided that I is a C -restricted complex structure and C is sufficiently big.

Theorem 5.14: Let M be a compact hyperkähler manifold, and $\mathcal{F} \in \mathcal{F}(M)$ a polystable hyperholomorphic sheaf on M (Definition 3.28). Let I be a C -restricted induced complex structure, for $C = \deg_I c_2(\mathcal{F})$,² and F' be a semistable torsion-free coherent sheaf on (M, I) with the same rank and Chern classes as $\mathcal{F}$. Then the sheaf F' is hyperholomorphic.

Proof: Let $F_1, \dots, F_n$ be the Jordan-Hölder series for the sheaf F' . Since $\mathcal{F}$ is hyperholomorphic, we have $\text{slope}(\mathcal{F}) = 0$ (Remark 3.12). Therefore, $\text{slope}(F_i) = 0$, and $\deg_I(c_1(F_i)) = 0$. By Definition 5.1 (i), then, the class $c_1(F_i)$ is $SU(2)$ invariant for all i . To prove that F' is hyperholomorphic it remains to show that the classes $c_2(F_i)$, $c_2(F_i^{**})$ are $SU(2)$ -invariant for all i .

Consider an exact sequence

$$0 \longrightarrow F_i \longrightarrow F_i^{**} \longrightarrow F_i/F_i^{**} \longrightarrow 0.$$

Let $[F_i/F_i^{**}] \in H^4(M)$ be the fundamental class of the union of all components of $\text{Sup}(F_i/F_i^{**})$ of complex codimension 2, taken with appropriate

²Clearly, since $\mathcal{F}$ is hyperholomorphic, the class $c_2(\mathcal{F})$ is $SU(2)$ -invariant, and the number $\deg_I c_2(\mathcal{F})$ independent from I .

multiplicities. Clearly, $c_2(F_i) = c_2(F_i^{**}) + [F_i/F_i^{**}]$. Since $[F_i/F_i^{**}]$ is an effective cycle, $\deg_I([F_i/F_i^{**}]) \geq 0$. By the Bogomolov-Miyaoka-Yau inequality (see Corollary 3.24), we have $\deg_I(c_2(F_i^{**})) \geq 0$. Therefore,

$$\deg_I c_2(F_i) \geq \deg_I c_2(F_i^{**}) \geq 0. \quad (5.10)$$

Using the product formula for Chern classes, we obtain

$$c_2(F) = \sum_i c_2(F_i) + \sum_{i,j} c_2(F_i) \wedge c_2(F_j). \quad (5.11)$$

By Corollary 4.10, $\deg_I(\sum_{i,j} c_2(F_i) \wedge c_2(F_j)) = 0$. Since the numbers $\deg_I c_2(F_i)$ are non-negative, we have $\deg_I c_2(F_i) \leq \deg_I c_2(F) = C$. By Theorem 4.15, the classes $c_2(F_i)$, $c_2(F_i^{**})$ are of CA-type. By Definition 5.1 (ii), then, the inequality $\deg_I c_2(F_i) \leq C$ implies that the class $c_2(F_i)$ is $SU(2)$ -invariant. By (5.10), $\deg_I c_2(F_i^{**}) \leq \deg_I c_2(F_i)$, so the class $c_2(F_i^{**})$ is also $SU(2)$ -invariant. Theorem 5.14 is proven. ■

6 Desingularization of hyperholomorphic sheaves

The aim of this section is the following theorem.

Theorem 6.1: Let M be a hyperkähler manifold, not necessarily compact, I an induced complex structure, and F a reflexive coherent sheaf over (M, I) equipped with a hyperholomorphic connection (Definition 3.15). Assume that F has isolated singularities. Let $\widetilde{M} \xrightarrow{\sigma} M$ be a blow-up of (M, I) in the singular set of F , and σ^*F the pullback of F . Then σ^*F is a locally trivial sheaf, that is, a holomorphic vector bundle.

We prove Theorem 6.1 in Subsection 6.4.

The idea of the proof is the following. We apply to F the methods used in the proof of Desingularization Theorem (Theorem 2.16). The main ingredient in the proof of Desingularization Theorem is the existence of a natural $\mathbb{C}^*$ -action on the completion $\hat{\mathcal{O}}_x(M, I)$ of the local ring $\mathcal{O}_x(M, I)$, for all $x \in M$. This $\mathbb{C}^*$ -action identifies $\hat{\mathcal{O}}_x(M, I)$ with a completion of a graded ring. Here we show that a sheaf F is $\mathbb{C}^*$ -equivariant. Therefore, a germ of F at x has a grading, which is compatible with the natural $\mathbb{C}^*$ -action on $\hat{\mathcal{O}}_x(M, I)$. Singularities of such reflexive sheaves can be resolved by a single blow-up.

6.1 Twistor lines and complexification

Further on, we need the following definition.

Definition 6.2: Let X be a real analytic variety, which is embedded to a complex variety $X_{\mathbb{C}}$. Assume that the sheaf of complex-valued real analytic functions on X coincides with the restriction of $\mathcal{O}_{X_{\mathbb{C}}}$ to $X \subset X_{\mathbb{C}}$. Then $X_{\mathbb{C}}$ is called a **complexification of X** .

For more details on complexification, the reader is referred to [GMT]. There are the most important properties.

Claim 6.3: In a neighbourhood of X , the manifold $X_{\mathbb{C}}$ has an anti-complex involution. The variety X is identified with the set of fixed points of this involution, considered as a real analytic variety.

Let Y be a complex variety, and X the underlying real analytic variety. Then the product of Y and its complex conjugate is a complexification of X , with embedding $X \hookrightarrow Y \times \bar{Y}$ given by the diagonal.

The complexification is unique in the following weak sense. For $X_{\mathbb{C}}$, $X'_{\mathbb{C}}$ complexifications of X , the complex manifolds $X_{\mathbb{C}}$, $X'_{\mathbb{C}}$ are naturally identified in a neighbourhood of X .

■

Let M be a hyperkähler manifold, $\text{Tw}(M)$ its twistor space, and $\pi : \text{Tw}(M) \rightarrow \mathbb{C}P^1$ the twistor projection. Let $l \subset \text{Tw}(M)$ be a rational curve, such that the restriction of π to l is an identity. Such a curve gives a section of π , and vice versa, every section of π corresponds to such a curve. The set of sections of the projection π is called **the space of twistor lines**, denoted by Lin , or $\text{Lin}(M)$. This space is equipped with complex structure, by Douady ([Do]).

Let $m \in M$ be a point. Consider a twistor line $s_m : I \rightarrow (I \times m) \in \mathbb{C}P^1 \times M = \text{Tw}$. Then s_m is called a **horizontal twistor line**. The space of horizontal twistor lines is a real analytic subvariety in Lin , denoted by Hor , or $\text{Hor}(M)$. Clearly, the set Hor is naturally identified with M .

Proposition 6.4: (Hitchin, Karlhede, Lindström, Roček) Let M be a hyperkähler manifold, $\text{Tw}(M)$ its twistor space, $I, J \in \mathbb{C}P^1$ induced complex structures, and Lin the space of twistor lines. The complex manifolds (M, I) and (M, J) are naturally embedded to $\text{Tw}(M)$:

$$(M, I) = \pi^{-1}(I), \quad (M, J) = \pi^{-1}(J).$$

Consider a point $s \in \text{Lin}$, $s : \mathbb{CP}^1 \longrightarrow \text{Tw}(M)$. Let

$$ev_{I,J} : \text{Lin}(M) \longrightarrow (M, I) \times (M, J)$$

be the map defined by $ev_{I,J}(s) = (s(I), s(J))$. Assume that $I \neq J$. Then there exists a neighbourhood U of $\text{Hor} \subset \text{Lin}$, such that the restriction of $ev_{I,J}$ to U is an open embedding.

Proof: [HKLR], [V-d3]. ■

Consider the anticomplex involution i of $\mathbb{CP}^1 \cong S^2$ which corresponds to the central symmetry of S^2 . Let $\iota : \text{Tw} \longrightarrow \text{Tw}$ be the corresponding involution of the twistor space $\text{Tw}(M) = \mathbb{CP}^1 \times M$, $(x, m) \longrightarrow (i(x), m)$. It is clear that ι maps holomorphic subvarieties of $\text{Tw}(M)$ to holomorphic subvarieties. Therefore, ι acts on Lin as an anticomplex involution. For $J = -I$, we obtain a local identification of Lin in a neighbourhood of Hor with $(M, I) \times (M, -I)$, that is, with (M, I) times its complex conjugate. Therefore, the space of twistor lines is a complexification of (M, I) . The natural anticomplex involution of Claim 6.3 coincides with ι . This gives an identification of Hor and the real analytic manifold underlying (M, I) .

We shall explain how to construct the natural $\mathbb{C}^*$ -action on a local ring of a hyperkähler manifold, using the machinery of twistor lines.

Fix a point $x_0 \in M$ and induced complex structures I, J , such that $I \neq \pm J$. Let V_1, V_2 be neighbourhoods of $s_{x_0} \in \text{Lin}$, and U_1, U_2 be neighbourhoods of (x_0, x_0) in $(M, I) \times (M, -I)$, $(M, J) \times (M, -J)$, such that the evaluation maps $ev_{I,-I}, ev_{J,-J}$ induce isomorphisms

$$ev_{I,-I} : V_1 \xrightarrow{\sim} U_1, \quad ev_{J,-J} : V_2 \xrightarrow{\sim} U_2.$$

Let B be an open neighbourhood of $x_0 \in M$, such that $(B, I) \times (B, -I) \subset U_1$ and $(B, I) \times (B, -I) \subset U_2$. Denote by $V_I \subset V_1$ be the preimage of $(B, I) \times (B, -I)$ under $ev_{I,-I}$, and by $V_J \subset V_2$ be the preimage of $(B, J) \times (B, -J)$ under $ev_{J,-J}$. Let $p_I : V_I \longrightarrow (B, I)$ be the evaluation, $s \longrightarrow s(I)$, and $e_I : (B, I) \longrightarrow V_I$ the map associating to $x \in B$ the unique twistor line passing through $(x, x_0) \in (B, I) \times (B, -I)$. In the same fashion, we define e_J and p_J . We are interested in the composition

$$\Psi_{I,J} := e_I \circ p_J \circ e_J \circ p_I : (B_0, I) \longrightarrow (B, I)$$

which is defined in a smaller neighbourhood $B_0 \subset B$ of $x_0 \in M$.

The following proposition is the focal point of this Subsection: we explain the map $\Psi_{I,J}$ of [V-d2], [V-d3] in geometric terms (in [V-d2], [V-d3] this map was defined algebraically).

Proposition 6.5: Consider the map $\Psi_{I,J} : (B_0, I) \longrightarrow (B, I)$ defined above. By definition, $\Psi_{I,J}$ preserves the point $x_0 \in B_0 \subset B$. Let $d\Psi_{I,J}$ be the differential of $\Psi_{I,J}$ acting on the tangent space $T_{x_0}B_0$. Assume that $I \neq \pm J$. Then $d\Psi_{I,J}$ is a multiplication by a scalar $\lambda \in \mathbb{C}$, $0 < |\lambda| < 1$.

Proof: The map $\Psi_{I,J}$ was defined in [V-d2], [V-d3] using the identifications between the real analytic varieties underlying (M, I) and (M, J) . We proved that $\Psi_{I,J}$ defined this way acts on $T_{x_0}B_0$ as a multiplication by the scalar $\lambda \in \mathbb{C}$, $0 < |\lambda| < 1$. It remains to show that the map $\Psi_{I,J}$ defined in [V-d2], [V-d3] coincides with $\Psi_{I,J}$ defined above.

Consider the natural identification

$$(B, I) \times (B, -I) \sim (B, J) \times (B, -J),$$

which is defined in a neighbourhood $B_{\mathbb{C}}$ of (x_0, x_0) . There is a natural projection $a_I : B_{\mathbb{C}} \longrightarrow (M, I)$. Consider the embedding $b_I : (B, I) \longrightarrow B_{\mathbb{C}}$, $x \longrightarrow (x, x_0)$, defined in a neighbourhood of $x_0 \in (B, I)$. In a similar way we define a_J , b_J . In [V-d2], [V-d3] we defined $\Psi_{I,J}$ as a composition $b_I \circ a_J \circ b_J \circ a_I$. Earlier in this Subsection, we described a local identification of $(B, I) \times (B, -I)$ and $\text{Lin}(B)$. Clearly, under this identification, the maps a_I , b_I correspond to p_I , e_I . Therefore, the definition of $\Psi_{I,J}$ given in this paper is equivalent to the definition given in [V-d2], [V-d3]. ■

6.2 The automorphism $\Psi_{I,J}$ acting on hyperholomorphic sheaves

In this section, we prove that hyperholomorphic sheaves are equivariant with respect to the map $\Psi_{I,J}$, considered as an automorphism of the local ring $\mathcal{O}_{x_0}(M, I)$.

Theorem 6.6: Let M be a hyperkähler manifold, not necessarily compact, $x_0 \in M$ a point, I an induced complex structure and F a reflexive sheaf over (M, I) equipped with a hyperholomorphic connection. Let $J \neq \pm I$ be another induced complex structure, and B_0 , B the neighbourhoods of $x_0 \in M$ for which the map $\Psi_{I,J} : B_0 \longrightarrow B$ was defined in Proposition

6.5. Assume that $\Psi_{I,J} : B_0 \longrightarrow B$ is an isomorphism. Then there exists a canonical functorial isomorphism of coherent sheaves

$$\Psi_{I,J}^F : F|_{B_0} \longrightarrow \Psi_{I,J}^*(F|_B).$$

Proof: Return to the notation introduced in Subsection 6.1. Let $W := V_I \cap V_J$. By definition of V_I, V_J , the evaluation maps produce open embeddings

$$ev_{I,-I} : \text{Lin}(W) \hookrightarrow (W, I) \times (W, -I),$$

and

$$ev_{J,-J} : \text{Lin}(W) \hookrightarrow (W, J) \times (W, -J),$$

Let $S \subset W$ be the singular set of $F|_W$, $\text{Tw}(S) \subset \text{Tw}(W)$ the corresponding embedding, and $L_0 \subset \text{Lin}(W)$ be the set of all lines $l \in \text{Lin}(W)$ which do not intersect $\text{Tw}(S)$. Consider the maps

$$p_I : L_0 \hookrightarrow (W, I) \setminus S$$

and

$$p_J : L_0 \hookrightarrow (W, J) \setminus S$$

obtained by restricting the evaluation map $p_I : \text{Lin}(M) \longrightarrow (M, I)$ to $L_0 \subset \text{Lin}(M)$. Since F is equipped with a hyperholomorphic connection, the vector bundle $F|_{(M,I) \setminus S}$ has a natural holomorphic structure. Let $\underline{E}_1 := p_I^*(F|_{(M,I) \setminus S})$ and $\underline{E}_2 := p_J^*(F|_{(M,J) \setminus S})$ be the corresponding pull-back sheaves over L_0 , and F_1, F_2 the sheaves on $\text{Lin}(W)$ obtained as direct images of $\underline{E}_1, \underline{E}_2$ under the open embedding $L_0 \hookrightarrow \text{Lin}(W)$.

Lemma 6.7: Under these assumptions, the sheaves F_1, F_2 are coherent reflexive sheaves. Moreover, there exists a natural isomorphism of coherent sheaves $\Psi_{1,2} : F_1 \longrightarrow F_2$.

Proof: The complex codimension of the singular set S in (M, I) is at least 3, because F is reflexive ([OSS], Ch. II, 1.1.10). Since S is trianalytic (Claim 3.16), this codimension is even. Thus, $\text{codim}_{\mathbb{C}}(S, (M, I)) \geq 4$. Therefore,

$$\text{codim}_{\mathbb{C}}(\text{Tw}(S), \text{Tw}(M)) \geq 4.$$

Consider the set L_S of all twistor lines $l \in \text{Lin}(W)$ passing through $\text{Tw}(S)$. For generic points $x, y \in \text{Tw}(W)$, there exists a unique line $l \in \text{Lin}(W)$ passing through x, y . Therefore,

$$\text{codim}_{\mathbb{C}}(L_S, \text{Lin}(W)) = \text{codim}_{\mathbb{C}}(\text{Tw}(S), \text{Tw}(M)) - 1 \geq 3.$$

By definition, $L_0 := \text{Lin}(W) \setminus L_S$. Since F_1, F_2 are direct images of bundles $\underline{F}_1, \underline{F}_2$ over a subvariety L_S of codimension 3, these sheaves are coherent and reflexive ([OSS], Ch. II, 1.1.12; see also Lemma 9.2). To show that they are naturally isomorphic it remains to construct an isomorphism between $\underline{F}_1$ and $\underline{F}_2$.

Let $\mathcal{F}$ be a coherent sheaf on $\text{Tw}(W)$ obtained from $F|_W$ as in the proof of Proposition 3.17. The singular set of $\mathcal{F}$ is $\text{Tw}(S) \subset \text{Tw}(W)$. Therefore, the restriction $\mathcal{F}|_{\text{Tw}(W) \setminus \text{Tw}(S)}$ is a holomorphic vector bundle. For all horizontal twistor lines $l_x \subset \text{Tw}(W) \setminus \text{Tw}(S)$, the restriction $\mathcal{F}|_{l_x}$ is clearly a trivial vector bundle over $l_x \cong \mathbb{CP}^1$. A small deformation of a trivial vector bundle is again trivial. Shrinking W if necessary, we may assume that for all lines $l \in L_0$, the restriction of $\mathcal{F}$ to $l \cong \mathbb{CP}^1$ is a trivial vector bundle.

The isomorphism $\underline{\Psi}_{1,2} : \underline{F}_1 \longrightarrow \underline{F}_2$ is constructed as follows. Let $l \in L_0$ be a twistor line. The restriction $\mathcal{F}|_l$ is trivial. Consider l as a map $l : \mathbb{CP}^1 \longrightarrow \text{Tw}(M)$. We identify $\mathbb{CP}^1$ with the set of induced complex structures on M . By definition, the fiber of F_1 in l is naturally identified with the space $\mathcal{F}|_{\iota(I)}$, and the fiber of F_2 in l identifies with $\mathcal{F}|_{\iota(J)}$. Since $\mathcal{F}|_l$ is trivial, the fibers of the bundle $\mathcal{F}|_l$ are naturally identified. This provides a vector bundle isomorphism $\underline{\Psi}_{1,2} : \underline{F}_1 \longrightarrow \underline{F}_2$ mapping $\underline{F}_1|_l = \mathcal{F}|_{\iota(I)}$ to $\underline{F}_2|_l = \mathcal{F}|_{\iota(J)}$. It remains to show that this isomorphism is compatible with the holomorphic structure.

Since the bundle $\mathcal{F}|_l$ is trivial, we have an identification

$$\mathcal{F}|_{\iota(I)} \cong \mathcal{F}|_{\iota(J)} = \Gamma(\mathcal{F}|_l),$$

where $\Gamma(\mathcal{F}|_l)$ is the space of global sections of $\mathcal{F}|_l$. Thus, $F_i|_l = \Gamma(\mathcal{F}|_l)$, and this identification is clearly holomorphic. This proves Lemma 6.7. ■

We return to the proof of Theorem 6.6. Denote by F_J the restriction of $\mathcal{F}$ to $(M, J) = \pi^{-1}(J) \subset \text{Tw}(M)$. The map $\Psi_{I,J}$ was defined as a composition $e_I \circ p_J \circ e_J \circ p_I$. The sheaf $p_I^* F$ is by definition isomorphic to F_1 , and $p_J^* F_J$ to F_2 . On the other hand, clearly, $e_J^* F_2 = F_J$. Therefore, $(p_J \circ e_J)^* F_2 \cong F_2$. Using the isomorphism $F_1 \cong F_2$, we obtain $(p_J \circ e_J)^* F_1 \cong F_1$. To sum it

up, we have the following isomorphisms:

$$\begin{aligned} p_I^* F &\cong F_1, \\ (p_J \circ e_J)^* F_1 &\cong F_1, \\ e_I^* F_1 &\cong F. \end{aligned}$$

A composition of these isomorphisms gives an isomorphism

$$\Psi_{I,J}^F : F|_{B_0} \longrightarrow \Psi_{I,J}^*(F|_B).$$

This proves Theorem 6.6. ■

6.3 A $\mathbb{C}^*$ -action on a local ring of a hyperkähler manifold

Let M be a hyperkähler manifold, non necessarily compact, $x \in M$ a point and I, J induced complex structures, $I \neq J$. Consider the complete local ring $\mathcal{O}_{x,I} := \hat{\mathcal{O}}_x(M, I)$. Throughout this section we consider the map $\Psi_{I,J}$ (Proposition 6.5) as an automorphism of the ring $\mathcal{O}_{x,I}$. Let $\mathfrak{m}$ be the maximal ideal of $\mathcal{O}_{x,I}$, and $\mathfrak{m}/\mathfrak{m}^2$ the Zariski cotangent space of (M, I) in x .

By Proposition 6.5, $\Psi_{I,J}$ acts on $\mathfrak{m}/\mathfrak{m}^2$ as a multiplication by a number $\lambda \in \mathbb{C}$, $0 < |\lambda| < 1$.

(6.1)

Let V_{λ^n} be the eigenspace corresponding to the eigenvalue λ^n ,

$$V_{\lambda^n} := \{v \in \mathcal{O}_{x,I} \mid \Psi_{I,J}(v) = \lambda^n v\}.$$

Clearly, $\oplus V_{\lambda^i}$ is a graded subring in $\mathcal{O}_{x,I}$. In [V-d2], (see also [V-d3]) we proved that the ring $\oplus V_{\lambda^i}$ is dense in $\mathcal{O}_{x,I}$ with respect to the adic topology. Therefore, the ring $\mathcal{O}_{x,I}$ is identified with the adic completion of $\oplus V_{\lambda^i}$.

Consider an action of $\mathbb{C}^*$ on $\oplus V_{\lambda^i}$, with $z \in \mathbb{C}^*$ acting on V_{λ^i} as a multiplication by z^i . This $\mathbb{C}^*$ -action is clearly continuous, with respect to the adic topology. Therefore, it can be extended to

$$\mathcal{O}_{x,I} = \widehat{\oplus V_{\lambda^i}}.$$

Definition 6.8: Let $M, I, J, x, \mathcal{O}_{x,I}$ be as in the beginning of this Subsection. Consider the $\mathbb{C}^*$ -action

$$\Psi_{I,J}(z) : \mathcal{O}_{x,I} \longrightarrow \mathcal{O}_{x,I}$$

constructed as above. Then $\Psi_{I,J}(z)$ is called **the canonical $\mathbb{C}^*$ -action associated with M, I, J, x** .

In the above notation, consider a reflexive sheaf F on (M, I) equipped with a hyperholomorphic connection. Denote the germ of F at x by F_x , $F_x := F \otimes_{\mathcal{O}_{(M,I)}} \mathcal{O}_{x,I}$. From Theorem 6.6, we obtain an isomorphism $F_x \cong \Psi_{I,J}^* F_x$. This isomorphism can be interpreted as an automorphism

$$\Psi_{I,J}^F : F_x \longrightarrow F_x$$

satisfying

$$\Psi_{I,J}^F(\alpha v) = \Psi_{I,J}(\alpha)v, \quad (6.2)$$

for all $\alpha \in \mathcal{O}_{x,I}$, $v \in F_x$.

By (6.2), the automorphism $\Psi_{I,J}^F$ respects the filtration

$$F_x \supset \mathfrak{m}F_x \supset \mathfrak{m}^2F_x \supset \dots$$

Thus, it makes sense to speak of $\Psi_{I,J}^F$ -action on $\mathfrak{m}^i F_x / \mathfrak{m}^{i+1} F_x$.

Lemma 6.9: The automorphism $\Psi_{I,J}^F$ acts on $\mathfrak{m}^i F_x / \mathfrak{m}^{i+1} F_x$ as a multiplication by λ^i , where $\lambda \in \mathbb{C}$ is the number considered in (6.1).

Proof: By (6.2), it suffices to prove Lemma 6.9 for $i = 0$. In other words, we have to show that $\Psi_{I,J}^F$ acts as identity on $F_x / \mathfrak{m}F_x$. We reduced Lemma 6.9 to the following claim.

Claim 6.10: In the above assumptions, the automorphism $\Psi_{I,J}^F$ acts as identity on $F_x / \mathfrak{m}F_x$.

Proof: In the course of defining the map $\Psi_{I,J}^F$, we identified the space $\text{Lin}(M)$ with a complexification of (M, I) , and defined the maps

$$p_I : \text{Lin}(M) \longrightarrow (M, I), \quad p_J : \text{Lin}(M) \longrightarrow (M, I)$$

(these maps are smooth, in a neighbourhood of $\text{Hor} \subset \text{Lin}(M)$, by Proposition 6.4), and

$$e_I : (B, I) \longrightarrow \text{Lin}(M), \quad e_J : (B, J) \longrightarrow \text{Lin}(M)$$

(these maps are locally closed embeddings). Consider (M, J) as a subvariety of $\text{Tw}(M)$, $(M, J) = \pi^{-1}(J)$. Let $\mathcal{F}$ be the lift of F to $\text{Tw}(M)$ (see the

proof of Proposition 3.17 for details). Denote the completion of $\mathcal{O}_x(M, J)$ by $\mathcal{O}_{x,J}$. Let F_J denote the $\mathcal{O}_{x,J}$ -module $\left(\mathcal{F}\Big|_{(M,J)}\right) \otimes_{\mathcal{O}_{(M,J)}} \hat{\mathcal{O}}_{x,J}$. Consider the horizontal twistor line $l_x \in \text{Lin}(M)$. Let $\text{Lin}_x(M)$ be the spectre of the completion $\mathcal{O}_{x,\text{Lin}}$ of the local ring of holomorphic functions on $\text{Lin}(M)$ in l_x . The maps p_I, p_J, e_I, e_J can be considered as maps of corresponding formal manifolds:

$$\begin{aligned} p_I &: \text{Lin}_x(M) \longrightarrow \text{Spec}(\mathcal{O}_{x,I}), \\ p_J &: \text{Lin}_x(M) \longrightarrow \text{Spec}(\mathcal{O}_{x,J}), \\ e_I &: \text{Spec}(\mathcal{O}_{x,I}) \longrightarrow \text{Lin}_x(M), \\ e_J &: \text{Spec}(\mathcal{O}_{x,J}) \longrightarrow \text{Lin}_x(M), \end{aligned}$$

As in Subsection 6.2, we consider the $\mathcal{O}_{x,\text{Lin}}$ -modules $F_1 := p_I^* F_x$ and $F_2 := p_J^* F_J$. By Lemma 6.7, there exists a natural isomorphism $\Psi_{1,2} : F_1 \longrightarrow F_2$.

Let $\mathfrak{m}_{l_x}$ be the maximal ideal of $\mathcal{O}_{x,\text{Lin}}$. Since the morphism p_I is smooth, the space $F_1/\mathfrak{m}_{l_x} F_1$ is naturally isomorphic to $F_x/\mathfrak{m} F_x$. Similarly, the space $F_2/\mathfrak{m}_{l_x} F_2$ is isomorphic to $F_J/\mathfrak{m}_J F_J$, where $\mathfrak{m}_J$ is the maximal ideal of $\mathcal{O}_{x,J}$. We have a chain of isomorphisms

$$\begin{aligned} F_x/\mathfrak{m} F_x &\xrightarrow{p_I^*} F_1/\mathfrak{m}_{l_x} F_1 \xrightarrow{\Psi_{1,2}} F_2/\mathfrak{m}_{l_x} F_2 \\ &\xrightarrow{e_J^*} F_J/\mathfrak{m}_J F_J \xrightarrow{p_J^*} F_2/\mathfrak{m}_{l_x} F_2 \\ &\xrightarrow{\Psi_{1,2}^{-1}} F_1/\mathfrak{m}_{l_x} F_1 \xrightarrow{e_I^*} F_x/\mathfrak{m} F_x. \end{aligned} \tag{6.3}$$

By definition, for any $f \in F_x/\mathfrak{m} F_x$, the value of $\Psi_{I,J}^F(f)$ is given by the composite map of (6.3) applied to f . The composition

$$F_2/\mathfrak{m}_{l_x} F_2 \xrightarrow{e_J^*} F_J/\mathfrak{m}_J F_J \xrightarrow{p_J^*} F_2/\mathfrak{m}_{l_x} F_2 \tag{6.4}$$

is identity, because the spaces $F_2/\mathfrak{m}_{l_x} F_2$ and $F_J/\mathfrak{m}_J F_J$ are canonically identified, and this identification can be performed via e_J^* or p_J^* . Thus, the map (6.3) is a composition

$$F_x/\mathfrak{m} F_x \xrightarrow{p_I^*} F_1/\mathfrak{m}_{l_x} F_1 \xrightarrow{\Psi_{1,2}} F_2/\mathfrak{m}_{l_x} F_2 \xrightarrow{\Psi_{1,2}^{-1}} F_1/\mathfrak{m}_{l_x} F_1 \xrightarrow{e_I^*} F_x/\mathfrak{m} F_x.$$

This map is clearly equivalent to a composition

$$F_x/\mathfrak{m} F_x \xrightarrow{p_I^*} F_1/\mathfrak{m}_{l_x} F_1 \xrightarrow{e_I^*} F_x/\mathfrak{m} F_x,$$

which is identity according to the same reasoning which proved that (6.4) is identity. We proved Claim 6.10 and Lemma 6.9. ■

Consider the λ^n -eigenspaces F_{λ^n} of F_x . Consider the $\oplus V_{\lambda^n}$ -submodule $\oplus F_{\lambda^n} \subset F_x$, where $\oplus V_{\lambda^n} \subset \mathcal{O}_{x,I}$ is the ring defined in Subsection 6.3. From Claim 6.10 and (6.1) it follows that $\oplus F_{\lambda^n}$ is dense in F_x , with respect to the adic topology on F_x . For $z \in \mathbb{C}^*$, let $\Psi_{I,J}^F(z) : \oplus F_{\lambda^n} \longrightarrow \oplus F_{\lambda^n}$ act on F_{λ^n} as a multiplication by z^n . As in Definition 6.8, we extend $\Psi_{I,J}^F(z)$ to $F_x = \widehat{\oplus F_{\lambda^n}}$. This automorphism makes F_x into a $\mathbb{C}^*$ -equivariant module over $\mathcal{O}_{x,I}$.

Definition 6.11: The constructed above $\mathbb{C}^*$ -equivariant structure on F_x is called **the canonical $\mathbb{C}^*$ -equivariant structure on F_x associated with J** .

6.4 Desingularization of $\mathbb{C}^*$ -equivariant sheaves

Let M be a hyperkähler manifold, I an induced complex structure and F a reflexive sheaf with isolated singularities over (M, I) , equipped with a hyperholomorphic connection. We have shown that the sheaf F admits a $\mathbb{C}^*$ -equivariant structure compatible with the canonical $\mathbb{C}^*$ -action on the local ring of (M, I) . Therefore, Theorem 6.1 is implied by the following proposition.

Proposition 6.12: Let B be a complex manifold, $x \in B$ a point. Assume that there is an action $\Psi(z)$ of $\mathbb{C}^*$ on B which fixes x and acts on $T_x B$ by dilatations. Let F be a reflexive coherent sheaf on B , which is locally trivial outside of x . Assume that the germ F_x of F in x is equipped with a $\mathbb{C}^*$ -equivariant structure, compatible with $\Psi(z)$. Let $\tilde{B}$ be a blow-up of B in x , and $\pi : \tilde{B} \longrightarrow B$ the standard projection. Then the pullback sheaf $\tilde{F} := \pi^* F$ is locally trivial on $\tilde{B}$.

Proof: Let $C := \pi^{-1}(x)$ be the singular locus of π . The sheaf F is locally trivial outside of x . Let d be the rank of $F|_{B \setminus x}$. To prove that $\tilde{F}$ is locally trivial, we need to show that for all points $y \in \tilde{B}$, the fiber $\pi^* F|_y$ is d -dimensional. Therefore, to prove Proposition 6.12 it suffices to show that $\pi^* F|_C$ is a vector bundle of dimension d .

The variety C is naturally identified with the projectivization $\mathbb{P}T_x B$ of

the tangent space $T_x B$. The total space of $T_x B$ is equipped with a natural action of $\mathbb{C}^*$, acting by dilatations. Clearly, coherent sheaves on $\mathbb{P}T_x B$ are in one-to-one correspondence with $\mathbb{C}^*$ -equivariant coherent sheaves on $T_x B$. Consider a local isomorphism $\varphi : T_x B \rightarrow B$ which is compatible with $\mathbb{C}^*$ -action, maps $0 \in T_x B$ to x and acts as identity on the tangent space $T_0(T_x B) = T_x B$. The sheaf $\varphi^* F$ is $\mathbb{C}^*$ -equivariant. Clearly, the corresponding sheaf on $\mathbb{P}T_x B$ is canonically isomorphic with $\pi^* F|_C$. Let $l \in T_x B$ be a line passing through 0, and $l \setminus 0$ its complement to 0. Denote the corresponding point of $\mathbb{P}T_x B$ by y . The restriction $\varphi^* F|_{l \setminus 0}$ is a $\mathbb{C}^*$ -equivariant vector bundle. The $\mathbb{C}^*$ -equivariant structure identifies all the fibers of the bundle $\varphi^* F|_{l \setminus 0}$. Let F_l be one of these fibers. Clearly, the fiber of $\pi^* F|_C$ in y is canonically isomorphic to F_l . Therefore, the fiber of $\pi^* F|_C$ in y is d -dimensional. We proved that $\pi^* F$ is a bundle. ■

7 Twistor transform and quaternionic-Kähler geometry

This Section is a compilation of results known from the literature. Subsection 7.1 is based on [KV] and the results of Subsection 7.2 are implicit in [KV]. Subsection 7.3 is based on [Sal], [N1] and [N2], and Subsection 7.4 is a recapitulation of the results of A. Swann ([Sw]).

7.1 Direct and inverse twistor transform

In this Subsection, we recall the definition and the main properties of the direct and inverse twistor transform for bundles over hyperkähler manifolds ([KV]).

The following definition is a non-Hermitian analogue of the notion of a hyperholomorphic connection.

Definition 7.1: Let M be a hyperkähler manifold, not necessarily compact, and (B, ∇) be a vector bundle with a connection over M , not necessarily Hermitian. Assume that the curvature of ∇ is contained in the space $\Lambda_{inv}^2(M, \text{End}(B))$ of $SU(2)$ -invariant 2-forms with coefficients in $\text{End}(B)$. Then (B, ∇) is called **an autodual bundle**, and ∇ **an autodual connection**.

Let $\mathrm{Tw}(M)$ be the twistor space of M , equipped with the standard maps $\pi : \mathrm{Tw}(M) \longrightarrow \mathbb{C}P^1$, $\sigma : \mathrm{Tw}(M) \longrightarrow M$.

We introduce the direct and inverse twistor transforms which relate autodual bundles on the hyperkähler manifold M and holomorphic bundles on its twistor space $\mathrm{Tw}(M)$.

Let B be a complex vector bundle on M equipped with a connection ∇ . The pullback σ^*B of B to $\mathrm{Tw}(M)$ is equipped with a pullback connection $\sigma^*\nabla$.

Lemma 7.2: ([KV], Lemma 5.1) The connection ∇ is autodual if and only if the connection $\sigma^*\nabla$ has curvature of Hodge type $(1, 1)$.

Proof: Follows from Lemma 2.6. ■

In assumptions of Lemma 7.2, consider the $(0, 1)$ -part $(\sigma^*\nabla)^{0,1}$ of the connection $\sigma^*\nabla$. Since $\sigma^*\nabla$ has curvature of Hodge type $(1, 1)$, we have

$$((\sigma^*\nabla)^{0,1})^2 = 0,$$

and by Proposition 2.19, this connection is integrable. Consider $(\sigma^*\nabla)^{0,1}$ as a holomorphic structure operator on σ^*B .

Let $\mathcal{A}$ be the category of autodual bundles on M , and $\mathcal{C}$ the category of holomorphic vector bundles on $\mathrm{Tw}(M)$. We have constructed a functor

$$(\sigma^*\bullet)^{0,1} : \mathcal{A} \longrightarrow \mathcal{C},$$

$\nabla \longrightarrow (\sigma^*\nabla)^{0,1}$. Let $s \in \mathrm{Hor} \subset \mathrm{Tw}(M)$ be a horizontal twistor line (Subsection 6.1). For any $(B, \nabla) \in \mathcal{A}$, consider corresponding holomorphic vector bundle $(\sigma^*B, (\sigma^*\nabla)^{0,1})$. The restriction of $(\sigma^*B, (\sigma^*\nabla)^{0,1})$ to $s \cong \mathbb{C}P^1$ is a trivial vector bundle. A converse statement is also true. Denote by $\mathcal{C}_0$ the category of holomorphic vector bundles C on $\mathrm{Tw}(M)$, such that the restriction of C to any horizontal twistor line is trivial.

Theorem 7.3: Consider the functor

$$(\sigma^*\bullet)^{0,1} : \mathcal{A} \longrightarrow \mathcal{C}_0$$

constructed above. Then it is an equivalence of categories.

Proof: [KV], Theorem 5.12. ■

Definition 7.4: Let M be a hyperkähler manifold, $\mathrm{Tw}(M)$ its twistor space and $\mathcal{F}$ a holomorphic vector bundle. We say that $\mathcal{F}$ is **compatible with twistor transform** if the restriction of $\mathcal{F}$ to any horizontal twistor line $s \in \mathrm{Tw}(M)$ is a trivial bundle on $s \cong \mathbb{CP}^1$.

Recall that a connection ∇ in a vector bundle over a complex manifold is called $(1,1)$ -connection if its curvature is of Hodge type $(1,1)$.

Remark 7.5: Let $\mathcal{F}$ be a holomorphic bundle over $\mathrm{Tw}(M)$ which is compatible with twistor transform. Then $\mathcal{F}$ is equipped with a natural $(1,1)$ -connection $\nabla_{\mathcal{F}} = \sigma^* \nabla$, where (B, ∇) is the corresponding autodual bundle over M . The connection $\nabla_{\mathcal{F}}$ is not, generally speaking, Hermitian, or compatible with a Hermitian structure.

7.2 Twistor transform and Hermitian structures on vector bundles

Results of this Subsection were implicit in [KV], but in this presentation, they are new.

Let M be a hyperkähler manifold, not necessarily compact, and $\mathrm{Tw}(M)$ its twistor space. In Subsection 7.1, we have shown that certain holomorphic vector bundles over $\mathrm{Tw}(M)$ admit a canonical $(1,1)$ -connection $\nabla_{\mathcal{F}}$ (Remark 7.5). This connection can be non-Hermitian. Here we study the Hermitian structures on $(\mathcal{F}, \nabla_{\mathcal{F}})$ in terms of holomorphic properties of $\mathcal{F}$.

Definition 7.6: Let F be a real analytic complex vector bundle over a real analytic manifold $X_{\mathbb{R}}$, and $h : F \times F \rightarrow \mathbb{C}$ a $\mathcal{O}_{X_{\mathbb{R}}}$ -linear pairing on F . Then h is called **semilinear** if for all $\alpha \in \mathcal{O}_{X_{\mathbb{R}}} \otimes_{\mathbb{R}} \mathbb{C}$, we have

$$h(\alpha x, y) = \alpha \cdot h(x, y), \quad \text{and} \quad h(x, \alpha y) = \overline{\alpha} \cdot h(x, y).$$

For X a complex manifold and F a holomorphic vector bundle, by a semilinear pairing on F we understand a semilinear pairing on the underlying real analytic bundle. Clearly, a real analytic Hermitian metric is always semilinear.

Let X be a complex manifold, $I : TX \rightarrow TX$ the complex structure operator, and $i : X \rightarrow X$ a real analytic map. We say that X is anticomplex if the induced morphism of tangent spaces satisfies $i \circ I = -I \circ i$. For a complex vector bundle F on X , consider the complex adjoint vector bundle

$\overline{F}$, which coincides with F as a real vector bundle, with $\mathbb{C}$ -action which is conjugate to that defined on F . Clearly, for every holomorphic vector bundle F , and any anticomplex map $i : X \longrightarrow X$, the bundle $i^*\overline{F}$ is equipped with a natural holomorphic structure.

Let M be a hyperkähler manifold, and $\text{Tw}(M)$ its twistor space. Recall that $\text{Tw}(M) = \mathbb{C}P^1 \times M$ is equipped with a canonical anticomplex involution ι , which acts as identity on M and as central symmetry $I \longrightarrow -I$ on $\mathbb{C}P^1 = S^2$. For any holomorphic bundle $\mathcal{F}$ on $\text{Tw}(M)$, consider the corresponding holomorphic bundle $\iota^*\overline{\mathcal{F}}$.

Let M be a hyperkähler manifold, I an induced complex structure, F a vector bundle over M , equipped with an autodual connection ∇ , and $\mathcal{F}$ the corresponding holomorphic vector bundle over $\text{Tw}(M)$, equipped with a canonical connection $\nabla_{\mathcal{F}}$. As usually, we identify (M, I) and the fiber $\pi^{-1}(I)$ of the twistor projection $\pi : \text{Tw}(M) \longrightarrow \mathbb{C}P^1$. Let $\nabla_{\mathcal{F}} = \nabla_I^{1,0} + \nabla_I^{0,1}$ be the Hodge decomposition of ∇ with respect to I .

Clearly, the operator $\nabla_I^{1,0}$ can be considered as a holomorphic structure operator on F , considered as a complex vector bundle over $(M, -I)$. (7.1)

Then the holomorphic structure operator on $\mathcal{F}|_{(M, I)}$ is equal to $\nabla_I^{0,1}$, and the holomorphic structure operator on $\mathcal{F}|_{(M, -I)}$ is equal to $\nabla_I^{1,0}$.

Assume that the bundle $(\mathcal{F}, \nabla_{\mathcal{F}})$ is equipped with a non-degenerate semilinear pairing h which is compatible with the connection. Consider the natural connection $\nabla_{\mathcal{F}^*}$ on the dual bundle to $\mathcal{F}$, and its Hodge decomposition (with respect to I)

$$\nabla_{\mathcal{F}^*} = \nabla_{\mathcal{F}^*}^{1,0} + \nabla_{\mathcal{F}^*}^{0,1}.$$

Clearly, the pairing h gives a C^∞ -isomorphism of $\mathcal{F}$ and the complex conjugate of its dual bundle, denoted as $\overline{\mathcal{F}}^*$. Since h is semilinear and compatible with the connection, it maps the holomorphic structure operator $\nabla_I^{0,1}$ to the complex conjugate of $\nabla_{\mathcal{F}^*}^{1,0}$. On the other hand, the operator $\nabla_{\mathcal{F}^*}^{1,0}$ is a holomorphic structure operator in $\mathcal{F}^*|_{(M, -I)}$, as (7.1) claims. We obtain that the map h can be considered as an isomorphism of holomorphic vector bundles

$$h : \mathcal{F} \longrightarrow (\iota^*\overline{\mathcal{F}})^*.$$

This correspondence should be thought of as a (direct) twistor transform for bundles with a semilinear pairing.

Proposition 7.7: (direct and inverse twistor transform for bundles with semilinear pairing) Let M be a hyperkähler manifold, and $\mathcal{C}_{sl}$ the category of autodual bundles over M equipped with a non-degenerate semilinear pairing. Consider the category $\mathcal{C}_{hol,sl}$ of holomorphic vector bundles $\mathcal{F}$ on $\text{Tw}(M)$, compatible with twistor transform and equipped with an isomorphism

$$\mathfrak{h} : \mathcal{F} \longrightarrow (\iota^* \overline{\mathcal{F}})^*.$$

Let $\mathcal{T} : \mathcal{C}_{sl} \longrightarrow \mathcal{C}_{hol,sl}$ be the functor constructed above. Then $\mathcal{T}$ is an isomorphism of categories.

Proof: Given a pair $\mathcal{F}, \mathfrak{h} : \mathcal{F} \longrightarrow (\iota^* \overline{\mathcal{F}})^*$, we need to construct a non-degenerate semilinear pairing h on $\mathcal{F}|_{(M,I)}$, compatible with a connection. Since $\mathcal{F}$ is compatible with twistor transform, it is a pullback of a bundle (F, ∇) on M . This identifies the real analytic bundles $\mathcal{F}|_{(M,I')}$, for all induced complex structures I' . Taking $I' = \pm I$, we obtain an identification of the C^∞ -bundles $\mathcal{F}|_{(M,I)}, \mathcal{F}|_{(M,-I)}$. Thus, $\mathfrak{h}$ can be considered as an isomorphism of $F = \mathcal{F}|_{(M,I)}$ and $(\overline{\mathcal{F}})^*|_{(M,I)}$. This allows one to consider $\mathfrak{h}$ as a semilinear form h on F . We need only to show that h is compatible with the connection ∇ . Since $\nabla_{\mathcal{F}}$ is an invariant of holomorphic structure, the map $\mathfrak{h} : \mathcal{F} \longrightarrow (\iota^* \overline{\mathcal{F}})^*$ is compatible with the connection $\nabla_{\mathcal{F}}$. Thus, the obtained above form h is compatible with the connection $\nabla_{\mathcal{F}}|_{(M,I)} = \nabla$. This proves Proposition 7.7. ■

7.3 B_2 -bundles on quaternionic-Kähler manifolds

Definition 7.8: ([Sal], [Bes]) Let M be a Riemannian manifold. Consider a bundle of algebras $\text{End}(TM)$, where TM is the tangent bundle to M . Assume that $\text{End}(TM)$ contains a 4-dimensional bundle of subalgebras $W \subset \text{End}(TM)$, with fibers isomorphic to a quaternion algebra $\mathbb{H}$. Assume, moreover, that W is closed under the transposition map $\perp : \text{End}(TM) \longrightarrow \text{End}(TM)$ and is preserved by the Levi-Civita connection. Then M is called **quaternionic-Kähler**.

Example 7.9: Consider the quaternionic projective space

$$\mathbb{H}P^n = (\mathbb{H}^n \setminus \{0\})/\mathbb{H}^*.$$

It is easy to see that $\mathbb{H}P^n$ is a quaternionic-Kähler manifold. For more examples of quaternionic-Kähler manifolds, see [Bes].

A quaternionic-Kähler manifold is Einstein ([Bes]), i. e. its Ricci tensor is proportional to the metric: $Ric(M) = c \cdot g$, with $c \in \mathbb{R}$. When $c = 0$, the manifold M is hyperkähler, and its restricted holonomy group is $Sp(n)$; otherwise, the restricted holonomy is $Sp(n) \cdot Sp(1)$. The number c is called **the scalar curvature** of M . Further on, we shall use the term *quaternionic-Kähler manifold* for manifolds with non-zero scalar curvature.

The quaternionic projective space $\mathbb{H}P^n$ has positive scalar curvature.

The quaternionic projective space is the only example of quaternionic-Kähler manifold which we need, in the course of this paper. However, the formalism of quaternionic-Kähler manifolds is very beautiful and significantly simplifies the arguments, so we state the definitions and results for a general quaternionic-Kähler manifold whenever possible.

Let M be a quaternionic-Kähler manifold, and $W \subset \text{End}(TM)$ the corresponding 4-dimensional bundle. For $x \in M$, consider the set $\mathcal{R}_x \subset W|_x$, consisting of all $I \in W|_x$ satisfying $I^2 = -1$. Consider $\mathcal{R}_x$ as a Riemannian submanifold of the total space of $W|_x$. Clearly, $\mathcal{R}_x$ is isomorphic to a 2-dimensional sphere. Let $\mathcal{R} = \cup_x \mathcal{R}_x$ be the corresponding spherical fibration over M , and $\text{Tw}(M)$ its total space. The manifold $\text{Tw}(M)$ is equipped with an almost complex structure, which is defined in the same way as the almost complex structure for the twistor space of a hyperkähler manifold. This almost complex structure is known to be integrable (see [Sal]).

Definition 7.10: ([Sal], [Bes]) Let M be a quaternionic-Kähler manifold. Consider the complex manifold $\text{Tw}(M)$ constructed above. Then $\text{Tw}(M)$ is called **the twistor space of M** .

Note that (unlike in the hyperkähler case) the space $\text{Tw}(M)$ is Kähler. For quaternionic-Kähler manifolds with positive scalar curvature, the anti-canonical bundle of $\text{Tw}(M)$ is ample, so $\text{Tw}(M)$ is a Fano manifold.

Quaternionic-Kähler analogue of a twistor transform was studied by T. Nitta in a serie of papers ([N1], [N2] etc.) It turns out that the picture given in [KV] for Kähler manifolds is very similar to that observed by T. Nitta.

A role of $SU(2)$ -invariant 2-forms is played by so-called B_2 -forms.

Definition 7.11: Let $SO(TM) \subset \text{End}(TM)$ be a group bundle of all orthogonal automorphisms of TM , and $G_M := W \cap SO(TM)$. Clearly, the fibers of G_M are isomorphic to $SU(2)$. Consider the action of G_M on the bundle of 2-forms $\Lambda^2(M)$. Let $\Lambda_{inv}^2(M) \subset \Lambda^2(M)$ be the bundle of G_M -invariants. The bundle $\Lambda_{inv}^2(M)$ is called **the bundle of B_2 -forms**. In a similar fashion we define B_2 -forms with coefficients in a bundle.

Definition 7.12: In the above assumptions, let (B, ∇) be a bundle with connection over M . The bundle B is called **a B_2 -bundle**, and ∇ is called **a B_2 -connection**, if its curvature is a B_2 -form.

Consider the natural projection $\sigma : \text{Tw}(M) \longrightarrow M$. The proof of the following claim is completely analogous to the proof of Lemma 2.6 and Lemma 7.2.

Claim 7.13:

- (i) Let ω be a 2-form on M . The pullback $\sigma^*\omega$ is of type $(1, 1)$ on $\text{Tw}(M)$ if and only if ω is a B_2 -form on M .
- (ii) Let B be a complex vector bundle on M equipped with a connection ∇ , not necessarily Hermitian. The pullback σ^*B of B to $\text{Tw}(M)$ is equipped with a pullback connection $\sigma^*\nabla$. Then, ∇ is a B_2 -connection if and only if $\sigma^*\nabla$ has curvature of Hodge type $(1, 1)$.

■

There exists an analogue of direct and inverse twistor transform as well.

Theorem 7.14: For any B_2 -connection (B, ∇) , consider the corresponding holomorphic vector bundle

$$(\sigma^*B, (\sigma^*\nabla)^{0,1}).$$

The restriction of $(\sigma^*B, (\sigma^*\nabla)^{0,1})$ to a line $\sigma^{-1}(m) \cong \mathbb{C}P^1$ is a trivial vector bundle, for any point $m \in M$. Denote by $\mathcal{C}_0$ the category of holomorphic vector bundles C on $\text{Tw}(M)$, such that the restriction of C to $\sigma^{-1}(m)$ is trivial, for all $m \in M$, and by $\mathcal{A}$ the category of B_2 -bundles (not necessarily Hermitian). Consider the functor

$$(\sigma^*\bullet)^{0,1} : \mathcal{A} \longrightarrow \mathcal{C}_0$$

constructed above. Then it is an equivalence of categories.

Proof: It is easy to modify the proof of the direct and inverse twistor transform theorem from [KV] to work in quaternionic-Kähler situation. ■

We will not use Theorem 7.14, except for its consequence, which was proven in [N1].

Corollary 7.15: Consider the functor

$$(\sigma^* \bullet)^{0,1} : \mathcal{A} \longrightarrow \mathcal{C}_0$$

constructed in Theorem 7.14. Then $(\sigma^* \bullet)^{0,1}$ gives an injection κ from the set of equivalence classes of Hermitian B_2 -connections over M to the set of equivalence classes of holomorphic connections over $\text{Tw}(M)$. ■

Let M be a quaternionic-Kähler manifold. The space $\text{Tw}(M)$ has a natural Kähler metric g , such that the standard map $\sigma : \text{Tw}(M) \longrightarrow M$ is a Riemannian submersion, and the restriction of g to the fibers $\sigma^{-1}(m)$ of σ is a metric of constant curvature on $\sigma^{-1}(m) = \mathbb{C}P^1$ ([Sal], [Bes]).

Example 7.16: In the case $M = \mathbb{H}P^n$, we have $\text{Tw}(M) = \mathbb{C}P^{2n+1}$, and the Kähler metric g is proportional to the Fubini-Study metric on $\mathbb{C}P^{2n+1}$.

Theorem 7.17: (T. Nitta) Let M be a quaternionic-Kähler manifold of positive scalar curvature, $\text{Tw}(M)$ its twistor space, equipped with a natural Kähler structure, and B a Hermitian B_2 -bundle on M . Consider the pull-back σ^*B , equipped with a Hermitian connection. Then σ^*B is a Yang-Mills bundle on $\text{Tw}(M)$, and $\deg c_1(\sigma^*B) = 0$.

Proof: [N2]. ■

Let κ be the map considered in Corollary 7.15. Assume that M is a compact manifold. In [N2], T. Nitta defined the moduli space of Hermitian B_2 -bundles. By Uhlenbeck-Yau theorem, Yang-Mills bundles are polystable. Then the map κ provides an embedding from the moduli of non-decomposable Hermitian B_2 -bundles to the moduli $\mathcal{M}$ of stable bundles on $\text{Tw}(M)$. The image of κ is a totally real subvariety in $\mathcal{M}$ ([N2]).

7.4 Hyperkähler manifolds with special $\mathbb{H}^*$ -action and quaternionic-Kähler manifolds of positive scalar curvature

Further on, we shall need the following definition.

Definition 7.18: An **almost hypercomplex manifold** is a smooth manifold M with an action of quaternion algebra in its tangent bundle. For each $L \in \mathbb{H}$, $L^2 = -1$, L gives an almost complex structure on M . The manifold M is called **hypercomplex** if the almost complex structure L is integrable, for all possible choices $L \in \mathbb{H}$.

The twistor space for a hypercomplex manifold is defined in the same way as for hyperkähler manifolds. It is also a complex manifold ([K]). The formalism of direct and inverse twistor transform can be repeated for hypercomplex manifolds verbatim.

Let $\mathbb{H}^*$ be the group of non-zero quaternions. Consider an embedding $SU(2) \hookrightarrow \mathbb{H}^*$. Clearly, every quaternion $h \in \mathbb{H}^*$ can be uniquely represented as $h = |h| \cdot g_h$, where $g_h \in SU(2) \subset \mathbb{H}^*$. This gives a natural decomposition $\mathbb{H}^* = SU(2) \times \mathbb{R}^{>0}$. Recall that $SU(2)$ acts naturally on the set of induced complex structures on a hyperkähler manifold.

Definition 7.19: Let M be a hyperkähler manifold equipped with a free smooth action ρ of the group $\mathbb{H}^* = SU(2) \times \mathbb{R}^{>0}$. The action ρ is called **special** if the following conditions hold.

- (i) The subgroup $SU(2) \subset \mathbb{H}^*$ acts on M by isometries.
- (ii) For $\lambda \in \mathbb{R}^{>0}$, the corresponding action $\rho(\lambda) : M \rightarrow M$ is compatible with the hyperholomorphic structure (which is a fancy way of saying that $\rho(\lambda)$ is holomorphic with respect to any of induced complex structures).
- (iii) Consider the smooth $\mathbb{H}^*$ -action $\rho_e : \mathbb{H}^* \times \text{End}(TM) \rightarrow \text{End}(TM)$ induced on $\text{End}(TM)$ by ρ . For any $x \in M$ and any induced complex structure I , the restriction $I|_x$ can be considered as a point in the total space of $\text{End}(TM)$. Then, for all induced complex structures I , all $g \in SU(2) \subset \mathbb{H}^*$, and all $x \in M$, the map $\rho_e(g)$ maps $I|_x$ to $g(I)|_{\rho_e(g)(x)}$.

Speaking informally, this can be stated as “ $\mathbb{H}^*$ -action interchanges the induced complex structures”.

- (iv) Consider the automorphism of S^2T^*M induced by $\rho(\lambda)$, where $\lambda \in \mathbb{R}^{>0}$. Then $\rho(\lambda)$ maps the Riemannian metric tensor $s \in S^2T^*M$ to $\lambda^2 s$.

Example 7.20: Consider the flat hyperkähler manifold $M_{\mathbb{H}} = \mathbb{H}^n \setminus 0$. There is a natural action of $\mathbb{H}^*$ on $\mathbb{H}^n \setminus 0$. This gives a special action of $\mathbb{H}^*$ on $M_{\mathbb{H}}$.

The case of a flat manifold $M_{\mathbb{H}} = \mathbb{H}^n \setminus 0$ is the only case where we apply the results of this section. However, the general statements are just as difficult to prove, and much easier to comprehend.

Definition 7.21: Let M be a hyperkähler manifold with a special action ρ of $\mathbb{H}^*$. Assume that $\rho(-1)$ acts non-trivially on M . Then $M/\rho(\pm 1)$ is also a hyperkähler manifold with a special action of $\mathbb{H}^*$. We say that the manifolds (M, ρ) and $(M/\rho(\pm 1), \rho)$ are **hyperkähler manifolds with special action of $\mathbb{H}^*$ which are special equivalent**. Denote by H_{sp} the category of hyperkähler manifolds with a special action of $\mathbb{H}^*$ defined up to special equivalence.

A. Swann ([Sw]) developed an equivalence between the category of quaternionic-Kähler manifolds of positive scalar curvature and the category H_{sp} . The purpose of this Subsection is to give an exposition of Swann’s formalism.

Let Q be a quaternionic-Kähler manifold. The restricted holonomy group of Q is $Sp(n) \cdot Sp(1)$, that is, $(Sp(n) \times Sp(1))/\{\pm 1\}$. Consider the principal bundle $\mathcal{G}$ with the fiber $Sp(1)/\{\pm 1\}$, corresponding to the subgroup

$$Sp(1)/\{\pm 1\} \subset (Sp(n) \times Sp(1))/\{\pm 1\}.$$

of the holonomy. There is a natural $Sp(1)/\{\pm 1\}$ -action on the space $\mathbb{H}^*/\{\pm 1\}$. Let

$$\mathcal{U}(Q) := \mathcal{G} \times_{Sp(1)/\{\pm 1\}} \mathbb{H}^*/\{\pm 1\}.$$

Clearly, $\mathcal{U}(Q)$ is fibered over Q , with fibers which are isomorphic to $\mathbb{H}^*/\{\pm 1\}$. We are going to show that the manifold $\mathcal{U}(Q)$ is equipped with a natural hypercomplex structure.

There is a natural smooth decomposition $\mathcal{U}(Q) \cong \mathcal{G} \times \mathbb{R}^{>0}$ which comes from the isomorphism $\mathbb{H}^* \cong Sp(1) \times \mathbb{R}^{>0}$.

Consider the standard 4-dimensional bundle W on Q . Let $x \in Q$ be a point. The fiber $W|_q$ is isomorphic to $\mathbb{H}$, in a non-canonical way. The choices of isomorphism $W|_q \cong \mathbb{H}$ are called **quaternion frames in q** . The set of quaternion frames gives a fibration over Q , with a fiber $\text{Aut}(\mathbb{H}) \cong Sp(1)/\{\pm 1\}$. Clearly, this fibration coincides with the principal bundle $\mathcal{G}$ constructed above. Since $\mathcal{U}(Q) \cong \mathcal{G} \times \mathbb{R}^{>0}$, a choice of $u \in \mathcal{U}(Q)|_q$ determines an isomorphism $W|_q \cong \mathbb{H}$.

Let (q, u) be the point of $\mathcal{U}(Q)$, with $q \in Q$, $u \in \mathcal{U}(Q)|_q$. The natural connection in $\mathcal{U}(Q)$ gives a decomposition

$$T_{(q,u)}\mathcal{U}(Q) = T_u\left(\mathcal{U}(Q)|_q\right) \oplus T_qQ.$$

The space $\mathcal{U}(Q)|_q \cong \mathbb{H}^*/\{\pm 1\}$ is equipped with a natural hypercomplex structure. This gives a quaternion action on $T_u\left(\mathcal{U}(Q)|_q\right)$. The choice of $u \in \mathcal{U}(Q)|_q$ determines a quaternion action on T_qQ , as we have seen above. We obtain that the total space of $\mathcal{U}(Q)$ is an almost hypercomplex manifold.

Proposition 7.22: (A. Swann) Let Q be a quaternionic-Kähler manifold. Consider the manifold $\mathcal{U}(Q)$ constructed as above, and equipped with a quaternion algebra action in its tangent space. Then $\mathcal{U}(Q)$ is a hypercomplex manifold.

Proof: Clearly, the manifold $\mathcal{U}(Q)$ is equipped with a $\mathbb{H}^*$ -action, which is related with the almost hypercomplex structure as prescribed by Definition 7.19 (ii)-(iii). Pick an induced complex structure $I \in \mathbb{H}$. This gives an algebra embedding $\mathbb{C} \rightarrow \mathbb{H}$. Consider the corresponding $\mathbb{C}^*$ -action ρ_I on an almost complex manifold $(\mathcal{U}(Q), I)$. This $\mathbb{C}^*$ -action is compatible with the almost complex structure. The quotient $\mathcal{U}(Q)/\rho(I)$ is an almost complex manifold, which is naturally isomorphic to the twistor space $\text{Tw}(Q)$. Let L^* be a complex vector bundle of all $(1, 0)$ -vectors $v \in T(\text{Tw}(Q))$ tangent to the fibers of the standard projection $\sigma : \text{Tw}(Q) \rightarrow Q$, and L be the

dual vector bundle. Denote by $Tot_{\neq 0}(L)$ the complement $Tot(L) \setminus N$, where $N = Tw(Q) \subset Tot(L)$ is the zero section of L . Using the natural connection in L , we obtain an almost complex structure on $Tot(L)$.

Consider the natural projection $\varphi : Tot_{\neq 0}(L) \longrightarrow Q$. The fibers $\varphi^{-1}(q)$ of φ are identified with the space of non-zero vectors in the total space of the cotangent bundle $T^*\sigma^{-1}(q) \cong T^*(\mathbb{C}P^1)$. This space is naturally isomorphic to

$$\mathcal{G}|_q \times \mathbb{R}^{>0} = \mathcal{U}(Q)|_q \cong \mathbb{H}^*/\{\pm 1\}.$$

This gives a canonical isomorphism of almost complex manifolds

$$(\mathcal{U}(Q), I) \longrightarrow Tot_{\neq 0}(L).$$

Therefore, to prove that $(\mathcal{U}(Q), I)$ is a complex manifold, it suffices to show that the natural almost complex structure on $Tot_{\neq 0}(L) \subset Tot(L)$ is integrable. Consider the natural connection ∇_L on L . To prove that $Tot(L)$ is a complex manifold, it suffices to show that ∇_L is a holomorphic connection. The bundle L is known under the name of **holomorphic contact bundle**, and it is known to be holomorphic ([Sal], [Bes]). ■

Remark 7.23: The result of Proposition 7.22 is well known. We have given its proof because we shall need the natural identification $Tot_{\neq 0}(L) \cong \mathcal{U}(Q)$ further on in this paper.

Theorem 7.24: Let Q be a quaternionic-Kähler manifold of positive scalar curvature, and $\mathcal{U}(Q)$ the hypercomplex manifold constructed above. Then $\mathcal{U}(Q)$ admits a unique (up to a scaling) hyperkähler metric compatible with the hypercomplex structure.

Proof: [Sw]. ■

Consider the action of $\mathbb{H}^*$ on $\mathcal{U}(M)$ defined in the proof of Proposition 7.22. This action satisfies the conditions (ii) and (iii) of Definition 7.19. The conditions (i) and (iv) of Definition 7.19 are easy to check (see [Sw] for details). This gives a functor from the category $\mathcal{C}$ of quaternionic-Kähler manifolds of positive scalar curvature to the category H_{sp} of Definition 7.21.

Theorem 7.25: The functor $Q \longrightarrow \mathcal{U}(Q)$ from $\mathcal{C}$ to H_{sp} is an equivalence of categories.

Proof: [Sw]. ■

The inverse functor from H_{sp} to C is constructed by taking a quotient of M by the action of $\mathbb{H}^*$. Using the technique of quaternionic-Kähler reduction and hyperkähler potentials ([Sw]), one can equip the quotient $M/\mathbb{H}^*$ with a natural quaternionic-Kähler structure.

8 $\mathbb{C}^*$ -equivariant twistor spaces

In Section 7, we gave an exposition of the twistor transform, B_2 -bundles and Swann's formalism. In the present Section, we give a synthesis of these theories, obtaining a construction which should be thought of as Swann's formalism for vector bundles.

Consider the equivalence of categories $Q \longrightarrow \mathcal{U}(Q)$ constructed in Theorem 7.25 (we call this equivalence "Swann's formalism"). We show that B_2 -bundles on Q are in functorial bijective correspondence with $\mathbb{C}^*$ -equivariant holomorphic bundles on $\text{Tw}(\mathcal{U}(Q))$ (Theorem 8.5).

In Subsection 8.4, this equivalence is applied to the vector bundle $\pi^*(F)$ of Theorem 6.1. We use it to construct a canonical Yang-Mills connection on $\pi^*(F)|_C$, where C is a special fiber of $\pi : \widetilde{M} \longrightarrow (M, I)$ (see Theorem 6.1 for details and notation). This implies that the holomorphic bundle $\pi^*(F)|_C$ is polystable (Theorem 8.15).

8.1 B_2 -bundles on quaternionic-Kähler manifolds and $\mathbb{C}^*$ -equivariant holomorphic bundles over twistor spaces

For the duration of this Subsection, we fix a hyperkähler manifold M , equipped with a special $\mathbb{H}^*$ -action ρ , and the corresponding quaternionic-Kähler manifold $Q = M/\mathbb{H}^*$. Denote the natural quotient map by $\varphi : M \longrightarrow Q$.

Lemma 8.1: Let ω be a 2-form over Q , and $\varphi^*\omega$ its pullback to M . Then the following conditions are equivalent

- (i) ω is a B_2 -form
- (ii) $\varphi^*\omega$ is of Hodge type $(1, 1)$ with respect to some induced complex structure I on M
- (iii) $\varphi^*\omega$ is $SU(2)$ -invariant.

Proof: Let I be an induced complex structure on M . As we have shown in the proof of Proposition 7.22, the complex manifold (M, I) is identified with an open subset of the total space $\text{Tot}(L)$ of a holomorphic line bundle L over $\text{Tw}(Q)$. The map φ is represented as a composition of the projections $h : \text{Tot}(L) \rightarrow \text{Tw}(Q)$ and $\sigma_Q : \text{Tw}(Q) \rightarrow Q$. Since the map h is smooth and holomorphic, the form $\varphi^*\omega$ is of Hodge type $(1, 1)$ if and only if $\sigma_Q^*\omega$ is of type $(1, 1)$. By Claim 7.13 (i), this happens if and only if ω is a B_2 -form. This proves an equivalence (i) $\Leftrightarrow$ (ii). Since the choice of I is arbitrary, the pullback $\varphi^*\omega$ of a B_2 -form is of Hodge type $(1, 1)$ with respect to all induced complex structures. By Lemma 2.6, this proves the implication (i) $\Rightarrow$ (iii). The implication (iii) $\Rightarrow$ (ii) is clear. ■

Proposition 8.2: Let (B, ∇) be a complex vector bundle with connection over Q , and $(\varphi^*B, \varphi^*\nabla)$ its pullback to M . Then the following conditions are equivalent

- (i) (B, ∇) is a B_2 -form
- (ii) The curvature of $(\varphi^*B, \varphi^*\nabla)$ is of Hodge type $(1, 1)$ with respect to some induced complex structure I on M
- (iii) The bundle $(\varphi^*B, \varphi^*\nabla)$ is autodual

Proof: Follows from Lemma 8.1 applied to $\omega = \nabla^2$. ■

For any point $I \in \mathbb{C}P^1$, consider the corresponding algebra embedding $\mathbb{C} \xrightarrow{c_I} \mathbb{H}$. Let ρ_I be the action of $\mathbb{C}^*$ on (M, I) obtained as a restriction of ρ to $c_I(\mathbb{C}^*) \subset \mathbb{H}^*$. Clearly from Definition 7.19 (ii), ρ_I acts on (M, I) by holomorphic automorphisms.

Consider $\text{Tw}(M)$ as a union

$$\text{Tw}(M) = \bigcup_{I \in \mathbb{C}P^1} \pi^{-1}(I), \quad \pi^{-1}(I) = (M, I)$$

Gluing $\rho(I)$ together, we obtain a smooth $\mathbb{C}^*$ -action $\rho_{\mathbb{C}}$ on $\text{Tw}(M)$.

Claim 8.3: Consider the action $\rho_{\mathbb{C}} : \mathbb{C}^* \times \text{Tw}(M) \rightarrow \text{Tw}(M)$ constructed above. Then $\rho_{\mathbb{C}}$ is holomorphic.

Proof: It is obvious from construction that $\rho_{\mathbb{C}}$ is compatible with the complex structure on $\text{Tw}(M)$. ■

Example 8.4: Let $M = \mathbb{H}^n \setminus 0$. Since $\mathrm{Tw}(\mathbb{H}^n)$ is canonically isomorphic to a total space of the bundle $\mathcal{O}(1)^n$ over $\mathbb{C}P^1$, the twistor space $\mathrm{Tw}(M)$ is $\mathrm{Tot}(\mathcal{O}(1)^n)$ without zero section. The group $\mathbb{C}^*$ acts on $\mathrm{Tot}(\mathcal{O}(1)^n)$ by dilatation, and the restriction of this action to $\mathrm{Tw}(M)$ coincides with $\rho_{\mathbb{C}}$.

Consider the map $\sigma : \mathrm{Tw}(M) \longrightarrow M$. Let (B, ∇) be a B_2 -bundle over Q . Since the bundle $(\varphi^*B, \varphi^*\nabla)$ is autodual, the curvature of $\sigma^*\varphi^*\nabla$ has type $(1, 1)$. Let $(\sigma^*\varphi^*B, (\sigma^*\varphi^*\nabla)^{0,1})$ be the holomorphic bundle obtained from $(\varphi^*B, \varphi^*\nabla)$ by twistor transform. Clearly, this bundle is $\mathbb{C}^*$ -equivariant, with respect to the natural $\mathbb{C}^*$ -action on $\mathrm{Tw}(M)$. It turns out that any $\mathbb{C}^*$ -equivariant bundle $\mathcal{F}$ on $\mathrm{Tw}(M)$ can be obtained this way, assuming that $\mathcal{F}$ is compatible with twistor transform.

Theorem 8.5: In the above assumptions, let $\mathcal{C}_{B_2}$ be the category of B_2 -bundles on Q , and $\mathcal{C}_{\mathrm{Tw}, \mathbb{C}^*}$ the category of $\mathbb{C}^*$ -equivariant holomorphic bundles on $\mathrm{Tw}(M)$ which are compatible with the twistor transform. Consider the functor

$$(\sigma^*\varphi^*)^{0,1} : \mathcal{C}_{B_2} \longrightarrow \mathcal{C}_{\mathrm{Tw}, \mathbb{C}^*},$$

$(B, \nabla) \longrightarrow (\sigma^*\varphi^*B, (\sigma^*\varphi^*\nabla)^{0,1})$, constructed above. Then $(\sigma^*\varphi^*)^{0,1}$ establishes an equivalence of categories.

We prove Theorem 8.5 in Subsection 8.3.

Remark 8.6: Let Q be an arbitrary quaternionic-Kähler manifold, and $M = \mathcal{U}(Q)$ the corresponding fibration. Then M is hypercomplex, and its twistor space is equipped with a natural holomorphic action of $\mathbb{C}^*$. This gives necessary ingredients needed to state Theorem 8.5 for Q with negative scalar curvature. The proof which we give for Q with positive scalar curvature will in fact work for all quaternionic-Kähler manifolds.

Question 8.7: What happens with this construction when Q is a hyperkähler manifold?

In this paper, we need Theorem 8.5 only in the case $Q = \mathbb{H}P^n$, $M = \mathbb{H}^n \setminus 0$, but the general proof is just as difficult.

8.2 $\mathbb{C}^*$ -equivariant bundles and twistor transform

Let M be a hyperkähler manifold, and $\text{Tw}(M)$ its twistor space. Recall that $\text{Tw}(M) = \mathbb{C}P^1 \times M$ is equipped with a canonical anticomplex involution ι , which acts as identity on M and as central symmetry $I \longrightarrow -I$ on $\mathbb{C}P^1 = S^2$.

Proposition 8.8: Let M be a hyperkähler manifold, and $\text{Tw}(M)$ its twistor space. Assume that $\text{Tw}(M)$ is equipped with a free holomorphic action $\rho(z) : \text{Tw}(M) \longrightarrow \text{Tw}(M)$ of $\mathbb{C}^*$, acting along the fibers of $\pi : \text{Tw}(M) \longrightarrow \mathbb{C}P^1$. Assume, moreover, that $\iota \circ \rho(z) = \rho(\bar{z}) \circ \iota$, where ι is the natural anticomplex involution of $\text{Tw}(M)$.¹ Let $\mathcal{F}$ be a $\mathbb{C}^*$ -equivariant holomorphic vector bundle on $\text{Tw}(M)$. Assume that $\mathcal{F}$ is compatible with the twistor transform. Let $\nabla_{\mathcal{F}}$ be the natural connection on $\mathcal{F}$ (Remark 7.5). Then $\nabla_{\mathcal{F}}$ is flat along the leaves of ρ .

Proof: First of all, let us recall the construction of the natural connection $\nabla_{\mathcal{F}}$. Let $\mathcal{F}$ be an arbitrary bundle compatible with the twistor transform. We construct $\nabla_{\mathcal{F}}$ in terms of the isomorphism $\Psi_{1,2}$ defined in Lemma 6.7.

Consider an induced complex structure I . Let F_I be the restriction of $\mathcal{F}$ to $(M, I) = \pi^{-1}(I) \subset \text{Tw}(M)$. Consider the evaluation map

$$p_I : \text{Lin}(M) \longrightarrow (M, I)$$

(Subsection 6.1). In a similar way we define the holomorphic vector bundle F_{-I} on $(M, -I)$ and the map $p_{-I} : \text{Lin}(M) \longrightarrow (M, -I)$. Denote by F_1, F_{-1} the sheaves $p_I^*(F_I), p_{-I}^*(F_{-I})$. In Lemma 6.7, we constructed an isomorphism $\Psi_{1,-1} : F_1 \longrightarrow F_{-1}$.

Let us identify $\text{Lin}(M)$ with $(M, I) \times (M, I)$ (this identification is naturally defined in a neighbourhood of $\text{Hor} \subset \text{Lin}(M)$ – see Proposition 6.4). Then the maps p_I, p_{-I} became projections to the relevant components. Let

$$\begin{aligned} \bar{\partial} : F_1 &\longrightarrow F_1 \otimes p_I^* \Omega^1(M, -I), \\ \partial : F_{-1} &\longrightarrow F_{-1} \otimes p_{-I}^* \Omega^1(M, I), \end{aligned}$$

be the sheaf maps obtained as pullbacks of de Rham differentials (the tensor product is taken in the category of coherent sheaves over $\text{Lin}(M)$). Twisting

¹These assumptions are automatically satisfied when M is equipped with a special $\mathbb{H}^*$ -action, and $\rho(z)$ is the corresponding $\mathbb{C}^*$ -action on $\text{Tw}(M)$.

∂ by an isomorphism $\Psi_{1,-1} : F_1 \longrightarrow F_{-1}$, we obtain a map

$$\partial^\Psi : F_1 \longrightarrow F_1 \otimes p_I^* \Omega^1(M, I).$$

Adding $\bar{\partial}$ and ∂^Ψ , we obtain

$$\nabla : F_1 \longrightarrow F_1 \otimes \left(p_I^* \Omega^1(M, I) \oplus p_I^* \Omega^1(M, -I) \right).$$

Clearly, ∇ satisfies the Leibniz rule. Moreover, the sheaf $p_I^* \Omega^1(M, I) \oplus p_I^* \Omega^1(M, -I)$ is naturally isomorphic to the sheaf of differentials over

$$\text{Lin}(M) = (M, I) \times (M, -I).$$

Therefore, ∇ can be considered as a connection in F_1 , or as a real analytic connection in a real analytic complex vector bundle underlying F_I . From the definition of $\nabla_{\mathcal{F}}$ ([KV]), it is clear that $\nabla_{\mathcal{F}}|_{(M, I)}$ equals ∇ .

Return to the proof of Proposition 8.8. Consider a $\mathbb{C}^*$ -action $\rho_I(z)$ on (M, I) , $(M, -I)$ induced from the natural embeddings $(M, I) \hookrightarrow \text{Tw}(M)$, $(M, -I) \hookrightarrow \text{Tw}(M)$. Then F_I is a $\mathbb{C}^*$ -equivariant bundle. Since $\iota \circ \rho(z) = \rho(\bar{z}) \circ \iota$, the identification $\text{Lin}(M) = (M, I) \times (M, I)$ is compatible with $\mathbb{C}^*$ -action. Let $\mathbf{r} = \frac{d}{dr}$ be the holomorphic vector field on (M, I) corresponding to the $\mathbb{C}^*$ -action. To prove Proposition 8.8, we have to show that the operator

$$[\nabla_{\mathbf{r}}, \nabla_{\bar{\mathbf{r}}}] : F_I \longrightarrow F_I \otimes \Lambda^{1,1}(M, I)$$

vanishes.

Consider the equivariant structure operator

$$\rho(z)^F : \rho_I(z)^* F_I \longrightarrow F_I.$$

Let U be a $\mathbb{C}^*$ -invariant Stein subset of (M, I) . Consider $\rho(z)^F$ as an endomorphism of the space of global holomorphic sections $\Gamma_U(F_I)$. Let

$$D_r(f) := \lim_{\varepsilon \rightarrow 0} \frac{\rho_I(1 + \varepsilon)}{\varepsilon},$$

for $f \in \Gamma_U(F_I)$. Clearly, D_r is a well defined sheaf endomorphism of F_I , satisfying

$$D_r(\alpha \cdot f) = \frac{d}{dr} \alpha \cdot f + \alpha \cdot D_r(f),$$

for all $\alpha \in \mathcal{O}_{(M,I)}$. We say that a holomorphic section f of F_I is $\mathbb{C}^*$ -**invariant** if $D_r(f) = 0$. Clearly, the $\mathcal{O}_{(M,I)}$ -sheaf F_I is generated by $\mathbb{C}^*$ -invariant sections. Therefore, it suffices to check the equality

$$[\nabla_{\mathbf{r}}, \nabla_{\bar{\mathbf{r}}}] (f) = 0$$

for holomorphic $\mathbb{C}^*$ -invariant $f \in F_I$.

Since f is holomorphic, we have $\nabla_{\bar{\mathbf{r}}} f = 0$. Thus,

$$[\nabla_{\mathbf{r}}, \nabla_{\bar{\mathbf{r}}}] (f) = \nabla_{\bar{\mathbf{r}}} \nabla_{\mathbf{r}} (f).$$

We obtain that Proposition 8.8 is implied by the following lemma.

Lemma 8.9: In the above assumptions, let f be a $\mathbb{C}^*$ -invariant section of F_I . Then $\nabla_{\mathbf{r}}(f) = 0$.

Proof: Return to the notation we used in the beginning of the proof of Proposition 8.8. Then, $\nabla(f) = \bar{\partial}(f) + \partial^\Psi(f)$. Since f is holomorphic, $\bar{\partial}(f) = 0$, so we need to show that $\partial^\Psi(f)(\mathbf{r}) = 0$. By definition of ∂^Ψ , this is equivalent to proving that

$$\partial \Psi_{1,-1}(f)(\mathbf{r}) = 0.$$

Consider the $\mathbb{C}^*$ -action on $\text{Lin}(M)$ which is induced by the $\mathbb{C}^*$ -action on $\text{Tw}(M)$. Since the maps p_I, p_{-I} are compatible with the $\mathbb{C}^*$ -action, the sheaves F_1, F_{-1} are $\mathbb{C}^*$ -equivariant. We can repeat the construction of the operator D_r for the sheaf F_{-I} . This allows one to speak of holomorphic $\mathbb{C}^*$ -invariant sections of F_{-I} . Pick a $\mathbb{C}^*$ -invariant Stein subset $U \subset (M, -I)$. Since the statement of Lemma 8.9 is local, we may assume that $M = U$. Let $g_1, \dots, g_n$ be a set of $\mathbb{C}^*$ -invariant sections of F_I which generated F_I . Then, the sections $p_{-I}^*(g_1), \dots, p_{-I}^*(g_n)$ generate F_{-1} . Consider the section $\Psi_{1,-1}(f)$ of F_{-1} . Clearly, $\Psi_{1,-1}$ commutes with the natural $\mathbb{C}^*$ -action. Therefore, the section $\Psi_{1,-1}(f)$ is $\mathbb{C}^*$ -invariant, and can be written as

$$\Psi_{1,-1}(f) = \sum \alpha_i p_{-I}^*(g_i),$$

where the functions α_i are $\mathbb{C}^*$ -invariant. By definition of ∂ we have

$$\partial \left(\sum \alpha_i p_{-I}^*(g_i) \right) = \sum \partial(\alpha_i p_{-I}^*(g_i)) + \sum \alpha_i \partial(p_{-I}^*(g_i)).$$

On the other hand, g_i is a holomorphic section of F_{-I} , so $\partial p_{-I}^*(g_i) = 0$. We obtain

$$\partial \left(\sum \alpha_i p_{-I}^*(g_i) \right) = \sum \partial \alpha_i p_{-I}^*(g_i).$$

Thus,

$$\partial\Psi_{1,-1}(f)(\mathbf{r}) = \sum \frac{\partial\alpha_i}{\partial r} p_{-I}^*(g_i),$$

but since the functions α_i are $\mathbb{C}^*$ -invariant, their derivatives along $\mathbf{r}$ vanish. We obtain $\partial\Psi_{1,-1}(f)(\mathbf{r}) = 0$. This proves Lemma 8.9. Proposition 8.8 is proven. ■

8.3 Twistor transform and the $\mathbb{H}^*$ -action

For the duration of this Subsection, we fix a hyperkähler manifold M , equipped with a special $\mathbb{H}^*$ -action ρ , and the corresponding quaternionic-Kähler manifold $Q = M/\mathbb{H}^*$. Denote the natural quotient map by $\varphi : M \rightarrow Q$. Clearly, Theorem 8.5 is an immediate consequence of the following theorem.

Theorem 8.10: Let $\mathcal{F}$ be a $\mathbb{C}^*$ -equivariant holomorphic bundle over $\text{Tw}(M)$, which is compatible with the twistor transform. Consider the natural connection $\nabla_{\mathcal{F}}$ on $\mathcal{F}$. Then $\nabla_{\mathcal{F}}$ is flat along the leaves of $\mathbb{H}^*$ -action.

Proof: The leaves of $\mathbb{H}^*$ -action are parametrized by the points of Q . Consider such a leaf $M_q := \varphi^{-1}(q) \subset M$. Clearly, M_q is a hyperkähler submanifold in M , equipped with a special action of $\mathbb{H}^*$. Moreover, the restriction of $\mathcal{F}$ to $\text{Tw}(M_q) \subset \text{Tw}(M)$ satisfies assumptions of Theorem 8.10. To prove that $\nabla_{\mathcal{F}}$ is flat along the leaves of $\mathbb{H}^*$ -action, we have to show that $\mathcal{F}|_{\text{Tw}(M_q)}$ is flat, for all q . Therefore, it suffices to prove Theorem 8.10 for $\dim_{\mathbb{H}} M = 1$.

Lemma 8.11: We work in notation and assumptions of Theorem 8.10. Assume that $\dim_{\mathbb{H}} M = 1$. Then the connection $\nabla_{\mathcal{F}}$ is flat.

Proof: Let I be an induced complex structure, and $F_I := F|_{(M,I)}$ the corresponding holomorphic bundle on (M, I) . Denote by z_I the vector field corresponding to the $\mathbb{C}^*$ -action ρ_I on (M, I) . By definition, the connection $\nabla|_{F_I}$ has $SU(2)$ -invariant curvature Θ_I . On the other hand, $\Theta_I(z_I, \bar{z}_I) = 0$ by Proposition 8.8. Since $\nabla_{\mathcal{F}} = \sigma^*\nabla$ is a pullback of an autodual connection ∇ on M , its curvature is a pullback of Θ_I . In particular, $\Theta = \Theta_I$ is independent from the choice of induced complex structure I . We obtain that $\Theta(z_I, \bar{z}_I) = 0$ for all induced complex structures I on M .

Now Lemma 8.11 is implied by the following linear-algebraic claim.

Claim 8.12: Let M be a hyperkähler manifold equipped with a special $\mathbb{H}^*$ -action, $\dim_{\mathbb{H}} M = 1$. Consider the vectors $z_I, \bar{z}_I$ defined above. Let Θ be a smooth $SU(2)$ -invariant 2-form, such that for all induced complex structures, I , we have $\Theta(z_I, \bar{z}_I) = 0$. Then $\Theta = 0$.

Proof: The proof of Claim 8.12 is an elementary calculation. Fix a point $m_0 \in M$. Consider the flat hyperkähler manifold $\mathbb{H} \setminus 0$, equipped with a natural special action of $\mathbb{H}^*$. From the definition of a special action, it is clear that the map ρ defines a covering $\mathbb{H} \setminus 0 \longrightarrow M$, $h \longrightarrow \rho(h)m_0$ of hyperkähler manifolds, and this covering is compatible with the special action. Therefore, the hyperkähler manifold M is flat, and the $\mathbb{H}^*$ -action is linear in the flat coordinates.

Let

$$\Lambda^2(M) = \Lambda^+(M) \oplus \Lambda^-(M)$$

be the standard decomposition of $\Lambda^2(M)$ according to the eigenvalues of the Hodge $*$ operator. Consider the natural Hermitian metric on $\Lambda^2(M)$. Then $\Lambda^-(M)$ is the bundle of $SU(2)$ -invariant 2-forms (see, e. g., [V1]), and $\Lambda^+(M)$ is its orthogonal complement. Consider the corresponding orthogonal projection $\Pi : \Lambda^2(M) \longrightarrow \Lambda^-(M)$. Denote by $dz_I \wedge d\bar{z}_I$ the differential form which is dual to the bivector $z_I \wedge \bar{z}_I$. Let $R \subset \Lambda^-(M)$ be the $C^\infty(M)$ -subsheaf of $\Lambda^-(M)$ generated by $\Pi(dz_I \wedge d\bar{z}_I)$, for all induced complex structures I on M . Clearly, $\Theta \in \Lambda^-(M)$ and Θ is orthogonal to $R \subset \Lambda^-(M)$. Therefore, to prove that $\Theta = 0$ it suffices to show that $R = \Lambda^-(M)$. Since M is covered by $\mathbb{H} \setminus 0$, we may prove $R = \Lambda^-(M)$ in assumption $M = \mathbb{H} \setminus 0$.

Let γ be the real vector field corresponding to dilatations of $M = \mathbb{H} \setminus 0$, and $d\gamma$ the dual 1-form. Clearly,

$$dz_I \wedge d\bar{z}_I = 2\sqrt{-1} d\gamma \wedge I(d\gamma).$$

Averaging $d\gamma \wedge I(d\gamma)$ by $SU(2)$, we obtain

$$\Pi(dz_I \wedge d\bar{z}_I) = \sqrt{-1} \left(d\gamma \wedge I(d\gamma) - J(d\gamma) \wedge K(d\gamma) \right)$$

where I, J, K is the standard triple of generators for quaternion algebra. Similarly,

$$\Pi(dz_J \wedge d\bar{z}_J) = \sqrt{-1} \left(d\gamma \wedge J(d\gamma) + K(d\gamma) \wedge I(d\gamma) \right)$$

and

$$\Pi(dz_K \wedge d\bar{z}_K) = \sqrt{-1} \left(d\gamma \wedge K(d\gamma) + I(d\gamma) \wedge J(d\gamma) \right)$$

Thus, $\Pi(R)$ is a 3-dimensional sub-bundle of $\Lambda^-(M)$. Since $\dim \Lambda^-(M) = 3$, we have $\Pi(R) = \Lambda^-(M)$. This proves Claim 8.12. Lemma 8.11 and Theorem 8.10 is proven. ■

8.4 Hyperholomorphic sheaves and $\mathbb{C}^*$ -equivariant bundles over $M_{\mathbb{H}}$

Let M be a hyperkähler manifold, I an induced complex structure and F a reflexive sheaf over (M, I) , equipped with a hyperholomorphic connection. Assume that F has an isolated singularity in $x \in M$. Consider the sheaf $\mathcal{F}$ on $\text{Tw}(M)$ corresponding to $\mathcal{F}$ as in the proof of Proposition 3.17. Let $s_x \subset \text{Tw}(M)$ be the horizontal twistor line corresponding to x , and $\mathfrak{m}$ its ideal. Consider the associated graded sheaf of $\mathfrak{m}$. Denote by Tw^{gr} the spectre of this associated graded sheaf. Clearly, Tw^{gr} is naturally isomorphic to $\text{Tw}(T_x M)$, where $T_x M$ is the flat hyperkähler manifold corresponding to the space $T_x M$ with induced quaternion action. Consider the natural $\mathbb{H}^*$ -action on $T_x M$. This provides the hyperkähler manifold $T_x M \setminus 0$ with a special $\mathbb{H}^*$ -action.

Let $s_0 \subset \text{Tw}^{gr}$ be the horizontal twistor line corresponding to s_x . The space $\text{Tw}^{gr} \setminus s_0$ is equipped with a holomorphic $\mathbb{C}^*$ -action (Claim 8.3). Denote by $\mathcal{F}^{gr}$ the sheaf on Tw^{gr} associated with $\mathcal{F}$. Clearly, $\mathcal{F}^{gr}$ is $\mathbb{C}^*$ -equivariant. In order to be able to apply Theorem 8.5 and Theorem 8.10 to $\mathcal{F}^{gr}|_{\text{Tw}^{gr} \setminus s_0}$, we need only to show that $\mathcal{F}^{gr}$ is compatible with twistor transform.

Proposition 8.13: Let M be a hyperkähler manifold, I an induced complex structure and F a reflexive sheaf over (M, I) , equipped with a hyperholomorphic connection. Assume that $\mathcal{F}$ has an isolated singularity in $x \in M$. Let $\mathcal{F}^{gr}$ be the $\mathbb{C}^*$ -equivariant bundle on $\text{Tw}^{gr} \setminus s_0$ constructed above. Then

- (i) the bundle $\mathcal{F}^{gr}$ is compatible with twistor transform.
- (ii) Moreover, the natural connection $\nabla_{\mathcal{F}^{gr}}$ (Remark 7.5) is Hermitian.

Proof: The argument is clear, but cumbersome, and essentially hinges on taking associate graded quotients everywhere and checking that all equa-

tions remain true. We give a simplified version of the proof, which omits some details and notation.

Consider the bundle $\mathcal{F}|_{M \setminus s_x}$. This bundle is compatible with twistor transform, and therefore, is equipped with a natural connection $\nabla_{\mathcal{F}}$. This connection is constructed using the isomorphism $\Psi_{1,-1} : F_1 \longrightarrow F_{-1}$ (see the proof of Proposition 8.8). We apply the same consideration to $\mathcal{F}^{gr}|_{(T_x M, I)}$, and show that the resulting connection $\nabla_{\mathcal{F}^{gr}}$ is hyperholomorphic. This implies that $\mathcal{F}^{gr}$ admits a $(1, 1)$ -connection which is a pullback of some connection on $\mathcal{F}^{gr}|_{(T_x M, I)}$. This argument is used to prove that $\mathcal{F}^{gr}$ is compatible with the twistor transform.

We use the notation introduced in the proof of Proposition 8.8. Let Lin^{gr} be the space of twistor maps in Tw^{gr} . Consider the maps $p_{\pm I}^{gr} : \text{Lin}^{gr} \longrightarrow (T_x M, \pm I)$ and the sheaves $F_{\pm 1}^{gr} := (p_{\pm I}^{gr})^* \mathcal{F}_{\pm I}^{gr}$ obtained in the same way as the maps $p_{\pm I}$ and the sheaves $F_{\pm 1}$ from the corresponding associated graded objects. Taking the associated graded of $\Psi_{1,-1}$ gives an isomorphism $\Psi_{1,-1}^{gr} : F_1^{gr} \longrightarrow F_{-1}^{gr}$. Using the same construction as in the proof of Proposition 8.8, we obtain a connection operator

$$\bar{\partial}^{gr} + \partial^{\Psi^{gr}} = \nabla_I^{gr} : F_1^{gr} \longrightarrow F_1^{gr} \otimes \left((p_{-I}^{gr})^* \Omega^1(T_x M, I) \oplus (p_I^{gr})^* \Omega^1(T_x M, -I) \right).$$

Since $(\bar{\partial}^{gr})^2 = (\partial^{\Psi^{gr}})^2 = 0$, the curvature of ∇_I^{gr} has Hodge type $(1, 1)$ with respect to I . To prove that ∇_I^{gr} is hyperholomorphic, we need to show that the curvature of ∇_I^{gr} has type $(1, 1)$ with respect to every induced complex structure. Starting from another induced complex structure J , we obtain a connection ∇_J^{gr} , with the curvature of type $(1, 1)$ with respect to J . To prove that ∇_J^{gr} is hyperholomorphic it remains to show that $\nabla_J^{gr} = \nabla_I^{gr}$.

Let ∇_I, ∇_J be the corresponding operators on F_1 . From the construction, it is clear that $\nabla_I^{gr}, \nabla_J^{gr}$ are obtained from ∇_I, ∇_J by taking the associated graded quotients. On the other hand, $\nabla_I = \nabla_J$. Therefore, the connections ∇_I^{gr} and ∇_J^{gr} are equal. We proved that the bundle $\mathcal{F}^{gr}|_{\text{Tw}^{gr} \setminus s_0}$ is compatible with the twistor structure. To prove Proposition 8.13, it remains to show that the natural connection on $\mathcal{F}^{gr}$ is Hermitian.

The bundle $\mathcal{F}|_{\text{Tw}(M \setminus x_0)}$ is by definition Hermitian. Consider the corresponding isomorphism $\mathcal{F} \longrightarrow (\iota^* \bar{\mathcal{F}})^*$ (Proposition 7.7). Taking an associated graded map, we obtain an isomorphism

$$\mathcal{F}^{gr} \xrightarrow{\sim} (\iota^* \bar{\mathcal{F}}^{gr})^*.$$

This gives a non-degenerate semilinear form h^{gr} on $\mathcal{F}^{gr}$. It remains only to show that h^{gr} is pseudo-Hermitian (i. e. satisfies $h(x, y) = \overline{h(y, x)}$) and positive definite.

Let $M_{\mathbb{C}}^{gr}$ be a complexification of $M^{gr} = T_x M$, $M_{\mathbb{C}}^{gr} = \text{Lin}(M^{gr})$. Consider the corresponding complex vector bundle $\mathcal{F}_{\mathbb{C}}^{gr}$ over $M_{\mathbb{C}}^{gr}$ underlying $\mathcal{F}^{gr}$. The metric h^{gr} can be considered as a semilinear form $\mathcal{F}_{\mathbb{C}}^{gr} \times \mathcal{F}_{\mathbb{C}}^{gr} \rightarrow \mathcal{O}_{M_{\mathbb{C}}^{gr}}$. This semilinear form is obtained from the corresponding form h on $\mathcal{F}$ by taking the associate graded quotients. Since h is Hermitian, the form h^{gr} is pseudo-Hermitian. To prove that h^{gr} is positive semidefinite, we need to show that for all $f \in \mathcal{F}_{\mathbb{C}}^{gr}$, the function $h^{gr}(f, \bar{f})$ belongs to $\mathcal{O}_{M_{\mathbb{C}}^{gr}} \cdot \overline{\mathcal{O}_{M_{\mathbb{C}}^{gr}}}$, where $\mathcal{O}_{M_{\mathbb{C}}^{gr}} \cdot \overline{\mathcal{O}_{M_{\mathbb{C}}^{gr}}}$ denotes the $\mathbb{R}^{>0}$ -semigroup of $\mathcal{O}_{M_{\mathbb{C}}^{gr}}$ generated by $x \cdot \bar{x}$, for all $x \in \mathcal{O}_{M_{\mathbb{C}}^{gr}}$. A similar property for h holds, because h is positive definite. Clearly, taking associated graded quotient of the semigroup $\mathcal{O}_{M_{\mathbb{C}}} \cdot \overline{\mathcal{O}_{M_{\mathbb{C}}}}$, we obtain $\mathcal{O}_{M_{\mathbb{C}}^{gr}} \cdot \overline{\mathcal{O}_{M_{\mathbb{C}}^{gr}}}$. Thus,

$$h^{gr}(f, \bar{f}) \in (\mathcal{O}_{M_{\mathbb{C}}} \cdot \overline{\mathcal{O}_{M_{\mathbb{C}}}})^{gr} = \mathcal{O}_{M_{\mathbb{C}}^{gr}} \cdot \overline{\mathcal{O}_{M_{\mathbb{C}}^{gr}}}$$

This proves that h^{gr} is positive semidefinite. Since h^{gr} is non-degenerate, this form is positive definite. Proposition 8.13 is proven. ■

Remark 8.14: Return to the notations of Theorem 6.1. Consider the bundle $\pi^* F|_C$, where $C = \mathbb{P}T_x M$ is the blow-up divisor. Clearly, this bundle corresponds to the graded sheaf $F_I^{gr} = \mathcal{F}^{gr}|_{(M, I)}$ on $(T_x M, I)$. By Proposition 8.13 (see also Theorem 8.10), the bundle $\pi^* F|_C$ is equipped with a natural $\mathbb{H}^*$ -invariant connection and Hermitian structure.² The sheaf $\pi^* F|_{\widetilde{M} \setminus C}$ is a hyperholomorphic bundle over $\widetilde{M} \setminus C \cong M \setminus x_0$. Therefore, $\pi^* F|_{\widetilde{M} \setminus C}$ is equipped with a natural metric and a hyperholomorphic connection. It is expected that the natural connection and metric on $\pi^* F|_{\widetilde{M} \setminus C}$ can be extended to $\pi^* F$, and the restriction of the resulting connection and metric to $\pi^* F|_C$ coincides with that given by Proposition 8.13 and Theorem 8.10. This will give an alternative proof of Proposition 8.13 (ii), because a continuous extension of a positive definite Hermitian metric is a positive semidefinite Hermitian metric.

²As usually, coherent sheaves over projective variety X correspond to finitely generated graded modules over the graded ring $\oplus \Gamma(\mathcal{O}_X(i))$.

8.5 Hyperholomorphic sheaves and stable bundles on $\mathbb{C}P^{2n+1}$

The purpose of the current Section was to prove the following result, which is a consequence of Proposition 8.13 and Theorem 8.10.

Theorem 8.15: Let M be a hyperkähler manifold, I an induced complex structure and F a reflexive sheaf on (M, I) admitting a hyperholomorphic connection. Assume that F has an isolated singularity in $x \in M$, and is locally trivial outside of x . Let $\pi : \widetilde{M} \longrightarrow (M, I)$ be the blow-up of (M, I) in x . Consider the holomorphic vector bundle π^*F on $\widetilde{M}$ (Theorem 6.1). Let $C \subset (M, I)$ be the blow-up divisor, $C = \mathbb{P}T_x M$. Then the holomorphic bundle $\pi^*F|_C$ admits a natural Hermitian connection ∇ which is flat along the leaves of the natural $\mathbb{H}^*$ -action on $\mathbb{P}T_x M$. Moreover, the connection ∇ is Yang-Mills, with respect to the Fubini-Study metric on $C = \mathbb{P}T_x M$, the degree $\deg c_1(\pi^*F|_C)$ vanishes, and the holomorphic vector bundle $\pi^*F|_C$ is polystable.

Proof: By definition, coherent sheaves on $C = \mathbb{P}T_x M$ correspond bijectively to $\mathbb{C}^*$ -equivariant sheaves on $T_x M \setminus 0$. Let F^{gr} be the associated graded sheaf of F (Subsection 8.4). Consider F^{gr} as a bundle on $T_x M \setminus 0$. In the notation of Proposition 8.13, $F^{gr} = \mathcal{F}^{gr}|_{(M, I)}$. By Proposition 8.13, the sheaf $\mathcal{F}^{gr}$ is $\mathbb{C}^*$ -equivariant and compatible with the twistor structure. According to the Swann's formalism for bundles (Theorem 8.10), the bundle $F^{gr}|_{T_x M \setminus 0}$ is equipped with a natural Hermitian connection $\nabla_{F^{gr}}$ which is flat along the leaves of $\mathbb{H}^*$ -action. Let $(B, \nabla_{\mathbb{H}})$ be the corresponding B_2 -bundle on

$$\mathbb{P}_{\mathbb{H}}T_x M := (T_x M \setminus 0)/\mathbb{H}^* \cong \mathbb{H}P^n.$$

Then $\pi^*F|_C$ is a holomorphic bundle over the corresponding twistor space $C = \text{Tw}(\mathbb{P}_{\mathbb{H}}T_x M)$, obtained as a pullback of $(B, \nabla_{\mathbb{H}})$ as in Claim 7.13. The natural Kähler metric on the twistor space $C = \text{Tw}(\mathbb{P}_{\mathbb{H}}T_x M)$ is the Fubini-Study metric (Example 7.16). By Theorem 7.17, the bundle $\pi^*F|_C$ is Yang-Mills and has $\deg c_1(\pi^*F|_C) = 0$. Finally, by Uhlenbeck-Yau theorem (Theorem 2.24), the bundle $\pi^*F|_C$ is polystable. ■

9 Moduli spaces of hyperholomorphic sheaves and bundles

9.1 Deformation of hyperholomorphic sheaves with isolated singularities

The following theorem is an elementary consequence of Theorem 8.15. The proof uses well known results on stability and reflexization (see, for instance, [OSS]). The main idea of the proof is the following. Given a family of hyperholomorphic sheaves with an isolated singularity, we blow-up this singularity and restrict the obtained family to a blow-up divisor. We obtain a family of coherent sheaves $\mathfrak{V}_s$, $s \in S$ over $\mathbb{C}P^{2n+1}$, with fibers semistable of slope zero. Assume that for all $s \in S$, $s \neq s_0$, the sheaf $\mathfrak{V}_s$ is trivial. Then the family $\mathfrak{V}$ is also trivial, up to a reflexization.

We use the following property of reflexive sheaves.

Definition 9.1: Let X be a complex manifold, and F a torsion-free coherent sheaf. We say that F is **normal** if for all open subvarieties $U \subset X$, and all closed subvarieties $Y \subset U$ of codimension 2, the restriction

$$\Gamma_U(F) \longrightarrow \Gamma_{U \setminus Y}(F)$$

is an isomorphism.

Lemma 9.2: Let X be a complex manifold, and F a torsion-free coherent sheaf. Then F is reflexive if and only if F is normal.

Proof: [OSS], Lemma 1.1.12. ■

Theorem 9.3: Let M be a hyperkähler manifold, I an induced complex structure, S a complex variety and $\mathfrak{F}$ a family of coherent sheaves over $(M, I) \times S$. Consider the sheaf $F_{s_0} := \mathfrak{F}|_{(M, I) \times \{s_0\}}$. Assume that the sheaf F_{s_0} is equipped with a filtration ξ . Let F_i , $i = 1, \dots, m$ denote the associated graded components of ξ , and F_i^{**} denote their reflexizations. Assume that $\mathfrak{F}$ is locally trivial outside of $(x_0, s_0) \in (M, I) \times S$. Assume, moreover, that all sheaves F_i^{**} , $i = 1, \dots, m$ admit a hyperholomorphic connection. Then the reflexization $\mathfrak{F}^{**}$ is locally trivial.

Proof: Clearly, it suffices to prove Theorem 9.3 for $\mathfrak{F}$ reflexive. Let $\tilde{X}$ be the blow-up of $(M, I) \times S$ in $\{x_0\} \times S$, and $\tilde{\mathfrak{F}}$ the pullback of $\mathfrak{F}$ to $\tilde{X}$. Clearly,

$\tilde{X} = \tilde{M} \times S$, where $\tilde{M}$ is a blow-up of (M, I) in x_0 . Denote by $C \subset \tilde{M}$ the blow-up divisor of $\tilde{M}$. Taking S sufficiently small, we may assume that the bundle $\mathfrak{F}|_{\{x_0\} \times (S \setminus \{s_0\})}$ is trivial. Thus, the bundle $\tilde{\mathfrak{F}}|_{(C \times S) \setminus (C \times \{s_0\})}$, which is a pullback of $\mathfrak{F}|_{\{x_0\} \times (S \setminus \{s_0\})}$ under the natural projection $(C \times S) \setminus (C \times \{s_0\}) \longrightarrow \{x_0\} \times (S \setminus \{s_0\})$ is trivial. To prove that $\mathfrak{F}$ is locally trivial, we have to show that $\tilde{\mathfrak{F}}$ is locally trivial, and that the restriction of $\tilde{\mathfrak{F}}$ to $C \times S$ is trivial along the fibers of the natural projection $C \times S \longrightarrow S$. Clearly, to show that $\tilde{\mathfrak{F}}$ is locally trivial we need only to prove that the fiber $\tilde{\mathfrak{F}}|_z$ has constant dimension for all $z \in C \times S$. Thus, $\tilde{\mathfrak{F}}$ is locally trivial if and only if $\tilde{\mathfrak{F}}|_{C \times S}$ is locally trivial. This sheaf is reflexive, since it corresponds to an associate graded sheaf of a reflexive sheaf, in the sense of Footnote to Remark 8.14. It is non-singular in codimension 2, because all reflexive sheaves are non-singular in codimension 2 ([OSS], Ch. II, Lemma 1.1.10). By Theorem 8.15, the sheaf $\tilde{\mathfrak{F}}|_{C \times \{s\}}$ is semistable of slope zero. Theorem 9.3 is implied by the following lemma, applied to the sheaf $\tilde{\mathfrak{F}}|_{C \times S}$.

Lemma 9.4: Let C be a complex projective space, S a complex variety and $\mathfrak{F}$ a torsion-free sheaf over $C \times S$. Consider an open set $U \xrightarrow{j} C \times S$, which is a complement of $C \times \{s_0\} \subset C \times S$. Assume that the sheaf $\mathfrak{F}|_U$ is trivial: $\mathfrak{F}|_U \cong \mathcal{O}_U^n$. Assume, moreover, that $\mathfrak{F}$ is non-singular in codimension 2, the sheaf $(\mathfrak{F}|_{C \times \{s_0\}})^{**}$ is semistable of slope zero and

$$\mathrm{rk} \mathfrak{F} = \mathrm{rk} \mathfrak{F}|_{C \times \{s_0\}}.$$

Then the reflexization $\mathfrak{F}^{**}$ of $\mathfrak{F}$ is a trivial bundle.

Proof: Using induction, it suffices to prove Lemma 9.4 assuming that it is proven for all $\mathfrak{F}'$ with $\mathrm{rk} \mathfrak{F}' < \mathrm{rk} \mathfrak{F}$. We may also assume that S is Stein, smooth and 1-dimensional.

Step 1: *We construct an exact sequence*

$$0 \longrightarrow \mathfrak{F}_2 \longrightarrow \mathfrak{F} \longrightarrow \mathrm{im} p_{O_1} \longrightarrow 0$$

of sheaves of positive rank, which, as we prove in Step 3, satisfy assumptions of Lemma 9.4.

Consider the pushforward sheaf $j_*\mathcal{O}_U^n$. From the definition of j_* , we obtain a canonical map

$$\mathfrak{F} \longrightarrow j_*\mathcal{O}_U^n, \quad (9.1)$$

and the kernel of this map is a torsion subsheaf in $\mathfrak{F}$.

Let f be a coordinate function on S , which vanishes in $s_0 \in S$. Clearly,

$$j_*\mathcal{O}_U^n \cong \mathcal{O}_{C \times S}^n \left[\frac{1}{f} \right].$$

On the other hand, the sheaf $\mathcal{O}_{C \times S} \left[\frac{1}{f} \right]$ is a direct limit of the following diagram:

$$\mathcal{O}_{C \times S}^n \xrightarrow{\cdot f} \mathcal{O}_{C \times S}^n \xrightarrow{\cdot f} \mathcal{O}_{C \times S}^n \xrightarrow{\cdot f} \dots,$$

where $\cdot f$ is the injection given by the multiplication by f . Thus, the map (9.1) gives an embedding

$$\mathfrak{F} \xhookrightarrow{p} \mathcal{O}_{C \times S}^n,$$

which is identity outside of (x_0, s_0) . Multiplying p by $\frac{1}{f}$ if necessary, we may assume that the restriction $p|_{C \times \{s_0\}}$ is non-trivial. Thus, p gives a map

$$\mathfrak{F}|_{C \times \{s_0\}} \longrightarrow \mathcal{O}_{C \times \{s_0\}}^n. \quad (9.2)$$

with image of positive rank. Since both sides of (9.2) are semistable of slope zero, and $\mathcal{O}_{C \times \{s_0\}}^n$ is polystable, the map (9.2) satisfies the following conditions. (see [OSS], Ch. II, Lemma 1.2.8 for details).

Let $F_1 := \text{im } p|_{C \times \{s_0\}}$, and $F_2 := \ker p|_{C \times \{s_0\}}$. Then the reflexization of F_1 is a trivial bundle $\mathcal{O}_{C \times \{s_0\}}^k$, and p maps F_1 to the direct summand $\mathcal{O}'_1 = \mathcal{O}_{C \times \{s_0\}}^k \subset \mathcal{O}_{C \times \{s_0\}}^n$.

Let $\mathcal{O}_1 = \mathcal{O}_{C \times S}^k \subset \mathcal{O}_{C \times S}^n$ be the corresponding free subsheaf of $\mathcal{O}_{C \times S}^n$. Consider the natural projection $\pi_{\mathcal{O}_1}$ of $\mathcal{O}_{C \times S}^n$ to $\mathcal{O}_1$. Let $p_{\mathcal{O}_1}$ be the composition of p and $\pi_{\mathcal{O}_1}$, $\mathfrak{F}_1$ the image of $p_{\mathcal{O}_1}$, and $\mathfrak{F}_2$ the kernel of $p_{\mathcal{O}_1}$.

Step 2: *We show that the sheaves $\mathfrak{F}_2$ and $\mathfrak{F}_1$ are non-singular in codimension 2.*

Consider the exact sequence

$$Tor^1(\mathcal{O}_{C \times \{s_0\}}, \mathfrak{F}_1) \longrightarrow \mathfrak{F}_2|_{C \times \{s_0\}} \longrightarrow \mathfrak{F}|_{C \times \{s_0\}} \longrightarrow \mathfrak{F}_1|_{C \times \{s_0\}} \longrightarrow 0$$

obtained by tensoring the sequence

$$0 \longrightarrow \mathfrak{F}_2 \longrightarrow \mathfrak{F} \longrightarrow \mathfrak{F}_1 \longrightarrow 0$$

with $\mathcal{O}_{C \times \{s_0\}}$. From this sequence, we obtain an isomorphism $\mathfrak{F}_1|_{C \times \{s_0\}} \cong F_1$.

A torsion-free coherent sheaf over a smooth manifold is non-singular in codimension 1 ([OSS], Ch. II, Corollary 1.1.8).

Since $\mathfrak{F}$ is non-singular in codimension 2, the restriction $\mathfrak{F}|_{C \times \{s_0\}}$ is non-singular in codimension 1. Therefore, the torsion of $\mathfrak{F}|_{C \times \{s_0\}}$ has support of codimension at least 2 in $C \times \{s_0\}$. Since the sheaf F_2 is a subsheaf of $\mathfrak{F}|_{C \times \{s_0\}}$, its torsion has support of codimension at least 2. Therefore, the singular set of F_2 has codimension at least 2 in $C \times \{s_0\}$. The rank of F_2 is by definition equal to $n - k$.

Since F_1 has rank k , the singular set of $\mathfrak{F}_1$ coincides with the singular set of F_1 . Since the restriction $\mathfrak{F}_1|_{C \times \{s_0\}} = F_1$, is a subsheaf of a trivial bundle of dimension k on $C \times \{s_0\}$, it is torsion-free. Therefore, the singularities of $\mathfrak{F}_1$ have codimension at least 2 in $C \times \{s_0\}$.

We obtain that the support of $Tor^1(\mathcal{O}_{C \times \{s_0\}}, \mathfrak{F}_1)$ has codimension at least 2 in $C \times \{s_0\}$. Since the quotient sheaf

$$\mathfrak{F}_2|_{C \times \{s_0\}} / Tor^1(\mathcal{O}_{C \times \{s_0\}}, \mathfrak{F}_1) \cong F_2 \quad (9.3)$$

is isomorphic to the sheaf F_2 , this quotient is non-singular in codimension 1. Since we proved that F_2 is non-singular in codimension 1, the sheaf $\mathfrak{F}_2|_{C \times \{s_0\}}$ is also non-singular in codimension 1, and its rank is equal to the rank of F_2 .

Let R be the union of singular sets of the sheaves $\mathfrak{F}_2, \mathfrak{F}, \mathfrak{F}_1$. Clearly, R is contained in $C \times \{s_0\}$, and R coincides with the set of all $x \in C \times \{s_0\}$ where the dimension of the fiber of the sheaves $\mathfrak{F}_2, \mathfrak{F}, \mathfrak{F}_1$ is not equal to $n - k, n, k$. We have seen that the restrictions of $\mathfrak{F}_2, \mathfrak{F}_1$ to $C \times \{s_0\}$ have ranks $n - k, k$. Therefore, the singular sets of $\mathfrak{F}_2, \mathfrak{F}_1$ coincide with the singular sets of $\mathfrak{F}_2|_{C \times \{s_0\}}, \mathfrak{F}_1|_{C \times \{s_0\}}$. We have shown that these singular

sets have codimension at least 2 in $C \times \{s_0\}$. On the other hand, $\mathfrak{F}$ is non-singular in codimension 2, by the conditions of Lemma 9.4. Therefore, R has codimension at least 3 in $C \times S$.

Step 3: *We check the assumptions of Lemma 9.4 applied to the sheaves $\mathfrak{F}_2, \mathfrak{F}_1$.*

Since the singular set of $\mathfrak{F}_1$ has codimension 2 in $C \times \{s_0\}$, the $\mathcal{O}_{C \times \{s_0\}}$ -sheaf $Tor^1(\mathcal{O}_{C \times \{s_0\}}, \mathfrak{F}_1)$ is a torsion sheaf with support of codimension 2 in $C \times \{s_0\}$. By (9.3), the reflexization of $\mathfrak{F}_2|_{C \times \{s_0\}}$ coincides with the reflexization of F_2 . Thus, the sheaf $\left(\mathfrak{F}_2|_{C \times \{s_0\}}\right)^{**}$ is semistable. On the other hand, outside of $C \times \{s_0\}$, the sheaf $\mathfrak{F}_2$ is a trivial bundle. Thus, $\mathfrak{F}_2$ satisfies assumptions of Lemma 9.4. Similarly, the sheaf $\mathfrak{F}_1$ is non-singular in codimension 2, its restriction to $C \times \{s_0\}$ has trivial reflexization, and it is free outside of $C \times \{s_0\}$.

Step 4: *We apply induction and prove Lemma 9.4.*

By induction assumption, the reflexization of $\mathfrak{F}_2$ is isomorphic to a trivial bundle $\mathcal{O}_{C \times S}^{n-k}$, and reflexization of $\mathfrak{F}_1$ is $\mathcal{O}_{C \times S}^k$. We obtain an exact sequence

$$0 \longrightarrow \mathfrak{F}_2 \longrightarrow \mathfrak{F} \longrightarrow \mathfrak{F}_1 \longrightarrow 0, \quad (9.4)$$

where the sheaves $\mathfrak{F}_2$ and $\mathfrak{F}_1$ have trivial reflexizations.

Let $V := C \times S \setminus R$. Restricting the exact sequence (9.4) to V , we obtain an exact sequence

$$0 \longrightarrow \mathcal{O}_V^{n-k} \xrightarrow{a} \mathfrak{F}|_V \xrightarrow{b} \mathcal{O}_V^k \longrightarrow 0. \quad (9.5)$$

Since V is a complement of a codimension-3 complex subvariety in a smooth Stein domain, the first cohomology of a trivial sheaf on V vanish. Therefore, the sequence (9.5) splits, and the sheaf $\mathfrak{F}|_V$ is a trivial bundle. Consider the pushforward $\zeta_* \mathfrak{F}|_V$, where $\zeta : V \longrightarrow C \times S$ is the standard map. Then $\zeta_* \mathfrak{F}|_V$ is a reflexization of $\mathfrak{F}$ (a pushforward of a reflexive sheaf over a subvariety of codimension 2 or more is reflexive – see Lemma 9.2). On the other hand, since the sheaf $\mathfrak{F}|_V$ is a trivial bundle, its push-forward over a subvariety of codimension at least 2 is also a trivial bundle over $C \times S$. We proved that the sheaf $\mathfrak{F}^{**} = \zeta_* \mathfrak{F}|_V$ is a trivial bundle over $C \times S$. The push-forward

$\zeta_* \mathfrak{F}|_V$ coincides with reflexization of $\mathfrak{F}$, by Lemma 9.2. This proves Lemma 9.4 and Theorem 9.3. ■

9.2 The Maruyama moduli space of coherent sheaves

This Subsection is a compilation of results of Gieseker and Maruyama on the moduli of coherent sheaves over projective manifolds. We follow [OSS], [Ma2].

To study the moduli spaces of holomorphic bundles and coherent sheaves, we consider the following definition of stability.

Definition 9.5: (Gieseker–Maruyama stability) ([Gi], [OSS]) Let X be a projective variety, $\mathcal{O}(1)$ the standard line bundle and F a torsion-free coherent sheaf. The sheaf F is called **Gieseker–Maruyama stable** (resp. Gieseker–Maruyama semistable) if for all coherent subsheaves $E \subset F$ with $0 < \text{rk } E < \text{rk } F$, we have

$$p_F(k) < p_E(k) \quad (\text{resp.}, \quad p_F(k) \leq p_E(k))$$

for all sufficiently large numbers $k \in \mathbb{Z}$. Here

$$p_F(k) = \frac{\dim \Gamma_X(F \otimes \mathcal{O}(k))}{\text{rk } F}.$$

Clearly, Gieseker–Maruyama stability is weaker than the Mumford–Takemoto stability. Every Gieseker–Maruyama semistable sheaf F has a so-called Jordan–Hölder filtration $F_0 \subset F_1 \subset \dots \subset F$ with Gieseker–Maruyama stable successive quotients F_i/F_{i-1} . The corresponding associated graded sheaf

$$\oplus F_i/F_{i-1}$$

is independent from a choice of a filtration. It is called **the associate graded quotient of the Jordan–Hölder filtration on F** .

Definition 9.6: Let F, G be Gieseker–Maruyama semistable sheaves on X . Then F, G are called **S -equivalent** if the corresponding associate graded quotients $\oplus F_i/F_{i-1}, \oplus G_i/G_{i-1}$ are isomorphic.

Definition 9.7: Let X be a complex manifold, F a torsion-free sheaf on X , and Y a complex variety. Consider a sheaf $\mathcal{F}$ on $X \times Y$ which is flat over Y . Assume that for some point $s_0 \in Y$, the sheaf $\mathcal{F}|_{X \times \{s_0\}}$ is isomorphic

to F . Then $\mathcal{F}$ is called a **deformation of F parametrized by Y** . We say that a sheaf F' on X is **deformationally equivalent** to F if for some $s \in Y$, the restriction $\mathcal{F}|_{X \times \{s\}}$ is isomorphic to F' . Slightly less formally, such sheaves are called **deformations of F** . If F' is a (semi-)stable bundle, it is called a **(semi-)stable bundle deformation of F** .

Remark 9.8: Clearly, the Chern classes of deformationally equivalent sheaves are equal.

Definition 9.9: Let X be a complex manifold, and F a torsion-free sheaf on X , and $\mathcal{M}_{mar}$ a complex variety. We say that $\mathcal{M}_{mar}$ is a **coarse moduli space of deformations of F** if the following conditions hold.

- (i) The points of $s \in \mathcal{M}_{mar}$ are in bijective correspondence with S -equivalence classes of coherent sheaves F_s which are deformationally equivalent to F .
- (ii) For any flat deformation $\mathcal{F}$ of F parametrized by Y , there exists a unique morphism $\varphi : Y \rightarrow \mathcal{M}_{mar}$ such that for all $s \in Y$, the restriction $\mathcal{F}|_{X \times \{s\}}$ is S -equivalent to the sheaf $F_{\varphi(s)}$ corresponding to $\varphi(s) \in \mathcal{M}_{mar}$.

Clearly, the coarse moduli space is unique. By Remark 9.8, the Chern classes of F_s are equal for all $s \in \mathcal{M}_{mar}$.

It is clear how to define other kinds of moduli spaces. For instance, replacing the word *sheaf* by the word *bundle* throughout Definition 9.9, we obtain a definition of **the coarse moduli space of semistable bundle deformations of F** . Further on, we shall usually omit the word “coarse” and say “moduli space” instead.

Theorem 9.10: (Maruyama) Let X be a projective manifold and F a coherent sheaf over X . Then the Maruyama moduli space $\mathcal{M}_{mar}$ of deformations of F exists and is compact.

Proof: See, e. g., [Ma2]. ■

9.3 Moduli of hyperholomorphic sheaves and C -restricted complex structures

Usually, the moduli space of semistable bundle deformations of a bundle F is not compact. To compactify this moduli space, Maruyama adds points corresponding to the deformations of F which are singular (these deformations can be non-reflexive and can have singular reflexizations). Using the desingularization theorems for hyperholomorphic sheaves, we were able to obtain Theorem 9.3, which states (roughly speaking) that a deformation of a semistable hyperholomorphic bundle is again a semistable bundle, assuming that all its singularities are isolated. In Section 5, we showed that under certain conditions, a deformation of a hyperholomorphic sheaf is again hyperholomorphic (Theorem 5.14). This makes it possible to prove that a deformation of a semistable hyperholomorphic bundle is locally trivial.

In [V5], we have shown that a Hilbert scheme of a K3 surface has no non-trivial trianalytic subvarieties, for a general hyperkähler structure.

Theorem 9.11: Let M be a compact hyperkähler manifold without non-trivial trianalytic subvarieties, $\dim_{\mathbb{H}} \geq 2$, and I an induced complex structure. Consider a hyperholomorphic bundle F on (M, I) (Definition 3.11). Assume that I is a C -restricted complex structure, $C = \deg_I c_2(F)$. Let $\mathcal{M}$ be the moduli space of semistable bundle deformations of F over (M, I) . Then $\mathcal{M}$ is compact.

Proof: The complex structure I is by definition algebraic, with unique polarization. This makes it possible to speak of Gieseker–Maruyama stability on (M, I) . Denote by $\mathcal{M}_{mar}$ the Maruyama moduli of deformations of F . Then $\mathcal{M}$ is naturally an open subset of $\mathcal{M}_{mar}$. Let $s \in \mathcal{M}_{mar}$ be an arbitrary point and F_s the corresponding coherent sheaf on (M, I) , defined up to S -equivalence. According to Remark 9.8, the Chern classes of F and F_s are equal. Thus, by Theorem 5.14, the sheaf F_s is hyperholomorphic. Therefore, F_s admits a filtration with hyperholomorphic stable quotient sheaves F_i , $i = 1, \dots, m$. By Claim 3.16, the singular set S of F_s is trianalytic. Since M has no proper trianalytic subvarieties, S is a collection of points. We obtain that F_s has isolated singularities. Let $\mathfrak{F}$ be a family of deformations of F , parametrized by Y . The points $y \in Y$ correspond to deformations F_y of F_s . Assume that for all $y \in Y$, $y \neq s$, the sheaf F_y is a bundle. Since $\mathcal{M}$ is open in $\mathcal{M}_{mar}$, such a deformation always exists.

The sheaf F_s has isolated singularities and admits a filtration with hy-

perholomorphic stable quotient sheaves. This implies that the family $\mathfrak{F}$ satisfies the conditions of Theorem 9.3. By Theorem 9.3, the reflexization $\mathfrak{F}^{**}$ is locally trivial. To prove that $\mathcal{M} = \mathcal{M}_{mar}$, we have to show that for all $s \in \mathcal{M}_{mar}$, the corresponding coherent sheaf F_s is locally trivial. Therefore, to finish the proof of Theorem 9.11, it remains to prove the following algebro-geometric claim.

Claim 9.12: Let X be a compact complex manifold, $\dim_{\mathbb{C}} X > 2$, and $\mathfrak{F}$ a torsion-free coherent sheaf over $X \times Y$ which is flat over Y . Assume that the reflexization of $\mathfrak{F}$ is locally trivial, $\mathfrak{F}$ has isolated singularities, and for some point $s \in Y$, the restriction of $\mathfrak{F}$ to the complement $(X \times Y) \setminus (X \times \{s\})$ is locally trivial. Then the reflexization $(\mathfrak{F}|_{X \times \{s\}})^{**}$ is locally trivial.

Remark 9.13: We say that a kernel of a map from a bundle to an Artinian sheaf is a **bundle with holes**. In slightly more intuitive terms, Claim 9.12 states that a flat deformation of a bundle with holes is again a bundle with holes, and cannot be smooth, assuming that $\dim_{\mathbb{C}} X > 2$.

Proof of Claim 9.12: Claim 9.12 is well known. Here we give a sketch of a proof. Consider a coherent sheaf $F_s = \mathfrak{F}|_{X \times \{s\}}$, and an exact sequence

$$0 \longrightarrow F_s \longrightarrow F_s^{**} \longrightarrow k \longrightarrow 0,$$

where k is an Artinian sheaf. By definition, the sheaf F_s^{**} is locally trivial. The flat deformations of F_s are infinitesimally classified by $Ext^1(F_s, F_s)$. Replacing F_s by a quasi-isomorphic complex of sheaves $F_s^{**} \longrightarrow k$, we obtain a spectral sequence converging to $Ext^{\bullet}(F_s, F_s)$. In the E_2 -term of this sequence, we observe the group

$$Ext^1(F_s^{**}, F_s^{**}) \oplus Ext^1(k, k) \oplus Ext^2(k, F_s^{**}) \oplus Ext^0(F_s^{**}, k).$$

which is responsible for $Ext^1(F_s, F_s)$. The term $Ext^1(F_s^{**}, F_s^{**})$ is responsible for deformations of the bundle F_s^{**} , the term $Ext^0(F_s^{**}, k)$ for the deformations of the map $F_s^{**} \longrightarrow k$, and the term $Ext^1(k, k)$ for the deformations of the Artinian sheaf k . Thus, the term $Ext^2(k, F_s^{**})$ is responsible for the deformations of F_s which change the dimension of the cokernel of the embedding $F_s \longrightarrow F_s^{**}$. We obtain that whenever $Ext^2(k, F_s^{**}) = 0$, all deformations of F_s are singular. On the other hand, $Ext^2(k, F_s^{**}) = 0$, because the i -th Ext from the skyscraper to a free sheaf on a manifold of dimension more than i vanishes (this is a basic result of Grothendieck's duality, [H-Gro]). ■

10 New examples of hyperkähler manifolds

10.1 Twistor paths

This Subsection contains an exposition and further elaboration of the results of [V3-bis] concerning the twistor curves in the moduli space of complex structures on a complex manifold of hyperkähler type.

Let M be a compact manifold admitting a hyperkähler structure. In Definition 5.8, we defined the coarse, marked moduli space of complex structures on M , denoted by $Comp$. For the duration of this section, we fix a compact simple hyperkähler manifold M , and its moduli $Comp$.

Further on, we shall need the following fact.

Claim 10.1: Let M be a hyperkähler manifold, I an induced complex structure of general type, and B a holomorphic vector bundle over (M, I) . Then B is stable if and only if B is simple.¹

Proof: By Lemma 2.26, for all $\omega \in Pic(M, I)$, we have $\deg_I(\omega) = 0$. Therefore, every subsheaf of B is destabilising. ■

Remark 10.2: In assumptions of Claim 10.1, all stable bundles are hyperholomorphic (Theorem 2.27). Therefore, Claim 10.1 implies that B is hyperholomorphic if it is simple.

In Subsection 5.2, we have shown that every hyperkähler structure $\mathcal{H}$ corresponds to a holomorphic embedding

$$\kappa(\mathcal{H}) : \mathbb{CP}^1 \longrightarrow Comp, \quad L \longrightarrow (M, L).$$

Definition 10.3: A projective line $C \subset Comp$ is called a **twistor curve** if $C = \kappa(\mathcal{H})$ for some hyperkähler structure $\mathcal{H}$ on M .

The following theorem was proven in [V3-bis].

Theorem 10.4: ([V3-bis], Theorem 3.1) Let $I_1, I_2 \in Comp$. Then there exist a sequence of intersecting twistor curves which connect I_1 with I_2 . ■

Definition 10.5: Let $P_0, \dots, P_n \subset Comp$ be a sequence of twistor curves, supplied with an intersection point $x_{i+1} \in P_i \cap P_{i+1}$ for each i . We say that

¹Simple sheaves are coherent sheaves which have no proper subsheaves

$\gamma = P_0, \dots, P_n, x_1, \dots, x_n$ is a **twistor path**. Let $I, I' \in \text{Comp}$. We say that γ is a **twistor path connecting I to I'** if $I \in P_0$ and $I' \in P_n$. The lines P_i are called **the edges**, and the points x_i **the vertices** of a twistor path.

Recall that in Definition 2.13, we defined induced complex structures which are generic with respect to a hyperkähler structure.

Given a twistor curve P , the corresponding hyperkähler structure $\mathcal{H}$ is unique (Theorem 5.11). We say that a point $x \in P$ is **of general type**, or **generic with respect to P** if the corresponding complex structure is generic with respect to $\mathcal{H}$.

Definition 10.6: Let $I, J \in \text{Comp}$ and $\gamma = P_0, \dots, P_n$ be a twistor path from I to J , which corresponds to the hyperkähler structures $\mathcal{H}_0, \dots, \mathcal{H}_n$. We say that γ is **admissible** if all vertices of γ are of general type with respect to the corresponding edges.

Remark 10.7: In [V3-bis], admissible twistor paths were defined slightly differently. In addition to the conditions above, we required that I, J are of general type with respect to $\mathcal{H}_0, \mathcal{H}_n$.

Theorem 10.4 proves that every two points I, I' in Comp are connected with a twistor path. Clearly, each twistor path induces a diffeomorphism $\mu_\gamma : (M, I) \longrightarrow (M, I')$. In [V3-bis], Subsection 5.2, we studied algebro-geometrical properties of this diffeomorphism.

Theorem 10.8: Let $I, J \in \text{Comp}$, and γ be an admissible twistor path from I to J . Then

- (i) There exists a natural isomorphism of tensor categories

$$\Phi_\gamma : \text{Bun}_I(\mathcal{H}_0) \longrightarrow \text{Bun}_J(\mathcal{H}_n),$$

where $\text{Bun}_I(\mathcal{H}_0), \text{Bun}_J(\mathcal{H}_n)$ are the categories of polystable hyperholomorphic vector bundles on $(M, I), (M, J)$, taken with respect to $\mathcal{H}_0, \mathcal{H}_n$ respectively.

- (ii) Let $B \in \text{Bun}_I(\mathcal{H}_0)$ be a stable hyperholomorphic bundle, and

$$\mathcal{M}_{I, \mathcal{H}_0}(B)$$

the moduli of stable deformations of B , where stability is taken with respect to the Kähler metric induced by $\mathcal{H}_0$. Then Φ_γ maps stable bundles which are deformationally equivalent to B to the stable bundles which are deformationally equivalent to $\Phi_\gamma(B)$. Moreover, obtained this way bijection

$$\Phi_\gamma : \mathcal{M}_{I, \mathcal{H}_0}(B) \longrightarrow \mathcal{M}_{J, \mathcal{H}_n}(\Phi_\gamma(B))$$

induces a real analytic isomorphism of deformation spaces.

Proof: Theorem 10.8 (i) is a consequence of [V3-bis], Corollary 5.1. Here we give a sketch of its proof.

Let I be an induced complex structure of general type. By Claim 10.1, a bundle B over (M, I) is stable if and only if it is simple. Thus, the category $Bun_I(\mathcal{H})$ is independent from the choice of $\mathcal{H}$ (Claim 10.1).

In Theorem 3.27, we constructed the equivalence of categories $\Phi_{I, J}$, which gives the functor Φ_γ for twistor path which consists of a single twistor curve. This proves Theorem 10.8 (i) for $n = 1$. A composition of isomorphisms $\Phi_{I, J} \circ \Phi_{J, J'}$ is well defined, because the category $Bun_I(\mathcal{H})$ is independent from the choice of $\mathcal{H}$. Taking successive compositions of the maps $\Phi_{I, J}$, we obtain an isomorphism Φ_γ . This proves Theorem 10.8 (i).

The variety $\mathcal{M}_{I, \mathcal{H}}(B)$ is singular hyperkähler ([V1]), and the variety $\mathcal{M}_{J, \mathcal{H}}(B)$ is the same singular hyperkähler variety, taken with another induced complex structure. By definition of singular hyperkähler varieties, this implies that $\mathcal{M}_{I, \mathcal{H}}(B)$, $\mathcal{M}_{J, \mathcal{H}}(B)$ are real analytic equivalent, with equivalence provided by $\Phi_{I, J}$. This proves Theorem 10.8 (ii). ■

For $I \in Comp$, denote by $Pic(M, I)$ the group $H^{1,1}(M, I) \cap H^2(M, \mathbb{Z})$, and by $Pic(I, \mathbb{Q})$ the space $H^{1,1}(M, I) \cap H^2(M, \mathbb{Q}) \subset H^2(M)$. Let $Q \subset H^2(M, \mathbb{Q})$ be a subspace of $H^2(M, \mathbb{Q})$, and

$$Comp_Q := \{I \in Comp \mid Pic(I, \mathbb{Q}) = Q\}.$$

Theorem 10.9: Let $\mathcal{H}, \mathcal{H}'$ be hyperkähler structures, and I, I' be complex structures of general type to and induced by $\mathcal{H}, \mathcal{H}'$. Assume that $Pic(I, \mathbb{Q}) = Pic(I', \mathbb{Q}) = Q$, and I, I' lie in the same connected component of $Comp_Q$. Then I, I' can be connected by an admissible path.

Proof: This is [V3-bis], Theorem 5.2. ■

For a general Q , we have no control over the number of connected components of $Comp_Q$ (unless global Torelli theorem is proven), and therefore

we cannot directly apply Theorem 10.9 to obtain results from algebraic geometry.² However, when $Q = 0$, $Comp_Q$ is clearly connected and dense in $Comp$. This is used to prove the following corollary.

Corollary 10.10: Let $I, I' \in Comp_0$. Then I can be connected to I' by an admissible twistor path.

Proof This is [V3-bis], Corollary 5.2. ■

Definition 10.11: Let $I \in Comp$ be a complex structure, ω be a Kähler form on (M, I) , and $\mathcal{H}$ the corresponding hyperkähler metric, which exists by Calabi-Yau theorem. Then ω is called a **generic polarization** if any of the following conditions hold

- (i) For all $a \in Pic(M, I)$, the degree $\deg_\omega(a) \neq 0$, unless $a = 0$.
- (ii) For all $SU(2)$ -invariant integer classes $a \in H^2(M, \mathbb{Z})$, we have $a = 0$.

The conditions (i) and (ii) are equivalent by Lemma 2.26.

Claim 10.12: Let $I \in Comp$ be a complex structure, ω be a Kähler form on (M, I) , and $\mathcal{H}$ the corresponding hyperkähler structure, which exists by Calabi-Yau theorem. Then ω is generic if and only if for all integer classes $a \in H^{1,1}(M, I)$, the class a is not orthogonal to ω with respect to the Bogomolov-Beauville pairing.

Proof: Clearly, the map $\deg_\omega : H^2(M) \rightarrow \mathbb{R}$ is equal (up to a scalar multiplier) to the orthogonal projection onto the line $\mathbb{R} \cdot \omega$. Then, Claim 10.12 is equivalent to Definition 10.11, (i). ■

From Claim 10.12 it is clear that the set of generic polarizations is a complement to a countable union of hyperplanes. Thus, generic polarizations are dense in the Kähler cone of (M, I) , for all I .

Claim 10.13: Let $I, J \in Comp$, and a, b be generic polarizations on (M, I) . Consider the corresponding hyperkähler structures $\mathcal{H}_0$ and $\mathcal{H}_n$ inducing I and J . Then there exists an admissible twistor path starting from $I, \mathcal{H}_0$ and ending with $\mathcal{H}_n, J$.

²Exception is a K3 surface, where Torelli holds. For K3, $Comp_Q$ is connected for all $Q \subset H^2(M, \mathbb{Q})$.

Proof: Consider the twistor curves P_0, P_n corresponding to $\mathcal{H}_0, \mathcal{H}_n$. Since a, b are generic, the curves P_0, P_n intersect with $Comp_0$. Applying Corollary 10.10, we connect the curves P_0 and P_n by an admissible path. ■

Putting together Claim 10.13 and Theorem 10.8, we obtain the following result.

Theorem 10.14: Let $I, J \in Comp$ be complex structures, and a, b be generic polarizations on $(M, I), (M, J)$. Then

- (i) There exist an isomorphism of tensor categories

$$\Phi_\gamma : Bun_I(a) \longrightarrow Bun_J(a),$$

where $Bun_I(a), Bun_J(b)$ are the categories of polystable hyperholomorphic vector bundles on $(M, I), (M, J)$, taken with respect to the hyperkähler structures defined by the Kähler classes a, b as in Theorem 2.4.

- (ii) Let $B \in Bun_I(a)$ be a stable hyperholomorphic bundle, and

$$\mathcal{M}_{I,a}(B)$$

the moduli of stable deformations of B , where stability is taken with respect to the polarization a . Then Φ_γ maps stable bundles which are deformationally equivalent to B to the stable bundles which are deformationally equivalent to $\Phi_\gamma(B)$. Moreover, obtained this way bijection

$$\Phi_\gamma : \mathcal{M}_{I,a}(B) \longrightarrow \mathcal{M}_{J,b}(\Phi_\gamma(B))$$

induces a real analytic isomorphism of deformation spaces.

■

Lemma 10.15: In assumptions of Theorem 10.8, let B be a holomorphic tangent bundle of (M, I) . Then $\Phi_\gamma(B)$ is a holomorphic tangent bundle of (M, J) .

Proof: Clear. ■

Corollary 10.16: Let $I, J \in Comp$ be complex structures, and a, b generic polarizations on $(M, I), (M, J)$. Assume that the moduli of stable deformations $\mathcal{M}_{I,a}(T(M, I))$ of the holomorphic tangent bundle $T^{1,0}(M, I)$ is compact. Then the space $\mathcal{M}_{J,b}(T(M, J))$ is also compact.

Proof: Let γ be the twistor path of Claim 10.13. By Lemma 10.15, $\Phi_\gamma(T(M, I)) = T(M, J)$. Applying Theorem 10.8, we obtain a real analytic equivalence from $\mathcal{M}_{I,a}(T(M, I))$ to $\mathcal{M}_{J,b}(T(M, J))$. ■

10.2 New examples of hyperkähler manifolds

Theorem 10.17: Let M be a compact hyperkähler manifold without non-trivial trianalytic subvarieties, $\dim_{\mathbb{H}} M \geq 2$, and I an induced complex structure. Consider a hyperholomorphic bundle F on M (Definition 3.28). Let F_I be the corresponding holomorphic bundle over (M, I) . Assume that I is a C -restricted complex structure, $C = \deg_I c_2(F)$. Assume, moreover, that all semistable bundle deformations of F_I are stable.³ Denote by $\mathcal{M}_F^I$ the moduli of stable bundle deformations of F_I over (M, I) . Then

- (i) the normalization $\widetilde{\mathcal{M}}_F^I$ is a compact and smooth complex manifold equipped with a natural hyperkähler structure.
- (ii) Moreover, for all induced complex structures J on M , the variety $\mathcal{M}_F^J$ is compact, and has a smooth normalization $\widetilde{\mathcal{M}}_F^J$, which is also equipped with a natural hyperkähler structure.
- (iii) Finally, the hyperkähler manifolds $\widetilde{\mathcal{M}}_F^J, \widetilde{\mathcal{M}}_F^I$ are naturally isomorphic.

Proof: The variety $\mathcal{M}_F^I$ is compact by Theorem 9.11. In [V1], it was proven that the space $\mathcal{M}_F^I$ of stable deformations of F is a singular hyperkähler variety (see also [KV] for an explicit construction of the twistor space of $\mathcal{M}_F^I$). Then Theorem 10.17 is a consequence of the Desingularization Theorem for singular hyperkähler varieties (Theorem 2.16). ■

The assumptions of Theorem 10.17 are quite restrictive. Using the technique of twistor paths, developed in Subsection 10.1, it is possible to prove a more accessible form of Theorem 10.17.

Let M be a hyperkähler manifold, and I, J induced complex structures. Given an admissible twistor path from I to J , we obtain an equivalence Φ_γ between the category of hyperholomorphic bundles on (M, I) and (M, J) .

Theorem 10.18: Let M be a compact simple hyperkähler manifold, $\dim_{\mathbb{H}} M > 1$, and I a complex structure on M . Consider a generic polarization a on (M, I) . Let $\mathcal{H}$ be the corresponding hyperkähler structure, and

³This may happen, for instance, when $\text{rk } F = \dim_{\mathbb{C}} M = n$, and the number $c_n(F)$ is prime.

F a hyperholomorphic bundle on (M, I) . Fix a hyperkähler structure $\mathcal{H}'$ on M admitting C -restricted complex structures, such that M has no tri-analytic subvarieties with respect to $\mathcal{H}'$. Assume that for some C -restricted complex structure J induced by $\mathcal{H}'$, $C = \deg_I c_2(F)$, all admissible twistor paths γ from I to J , and all semistable bundles F' which are deformationally equivalent to $\Phi_\gamma(F)$, the bundle F' is stable. Then the space of stable deformations of F is compact.

Remark 10.19: The space of stable deformations of F is singular hyperkähler ([V1]) and its normalization is smooth and hyperkähler (Theorem 2.16).

Proof of Theorem 10.18: Clearly, F' satisfies assumptions of Theorem 10.17, and the moduli space of its stable deformations is compact. Since Φ_γ induces a homeomorphism of moduli spaces (Theorem 10.8), the space of stable deformations of F is also compact. ■

Applying Theorem 10.18 to the holomorphic tangent bundle $T(M, I)$, we obtain the following corollary.

Theorem 10.20: Let M be a compact simple hyperkähler manifold, $\dim_{\mathbb{H}}(M) > 1$. Assume that for a generic hyperkähler structure $\mathcal{H}$ on M , this manifold admits no tri-analytic subvarieties.⁴ Assume, moreover, that for some C -restricted induced complex structure I , all semistable bundle deformations of $T(M, I)$ are stable, for $C > \deg_I c_2(M)$. Then, for all complex structures J on M and all generic polarizations ω on (M, J) , the deformation space $\mathcal{M}_{J, \omega}(T(M, J))$ is compact.

Proof: Follows from Theorem 10.18 and Corollary 10.16. ■

10.3 How to check that we obtained new examples of hyperkähler manifolds?

A. Beauville [Bea] described two families of compact hyperkähler manifolds, one obtained as the Hilbert scheme of points on a K3-surface, another obtained as the Hilbert scheme of a 2-dimensional torus factorized by the free torus action.

⁴This assumption holds for a Hilbert scheme of points on a K3 surface.

Conjecture 10.21: There exist compact simple hyperkähler manifolds which are not isomorphic to deformations of these two fundamental examples.

Here we explain our strategy of a proof of Conjecture 10.21 using results on compactness of the moduli space of hyperholomorphic bundles.

The results of this subsection are still in writing, so all statements below this line should be considered as conjectures. We give an idea of a proof for each result and label it as “proof”, but these “proofs” are merely sketches.

First of all, it is possible to prove the following theorem.

Theorem 10.22: Let M be a complex K3 surface without automorphisms. Assume that M is Mumford-Tate generic with respect to some hyperkähler structure. Consider the Hilbert scheme $M^{[n]}$ of points on M , $n > 1$. Pick a hyperkähler structure $\mathcal{H}$ on $M^{[n]}$ which is compatible with the complex structure. Let B be a hyperholomorphic bundle on $(M^{[n]}, \mathcal{H})$, $\text{rk } B = 2$. Then B is a trivial bundle.

Proof: The proof of Theorem 10.22 is based on the same ideas as the proof of Theorem 2.17. ■

For a compact complex manifold X of hyperkähler type, denote its coarse, marked moduli space (Definition 5.8) by $\text{Comp}(X)$.

Corollary 10.23: Let M be a K3 surface, $I \in \text{Comp}(X)$ an arbitrary complex structure on $X = M^{[n]}$, $n > 1$, and a a generic polarization on (X, I) . Consider the hyperkähler structure $\mathcal{H}$ which corresponds to (I, a) as in Theorem 2.4. Let B , $\text{rk } B = 2$ be a hyperholomorphic bundle over $(X, \mathcal{H})$. Then B is trivial.

Proof: Follows from Theorem 10.22 and Theorem 10.14. ■

Corollary 10.24: Let M be a K3 surface, $I \in \text{Comp}(X)$ an arbitrary complex structure on $X = M^{[n]}$, $n > 1$, and a a generic polarization on (X, I) . Consider the hyperkähler structure $\mathcal{H}$ which corresponds to (I, a) (Theorem 2.4). Let B , $\text{rk } B \leq 6$ be a stable hyperholomorphic bundle on $(X, \mathcal{H})$. Assume that the Chern class $c_{\text{rk } B}(B)$ is non-zero. Assume, moreover, that I is C -restricted, $C = \deg_I(c_2(B))$. Let B' be a semistable deformation of B over (X, I) . Then B' is stable.

Proof: Consider the Jordan–Hölder series for B' . Let $Q_1 \oplus Q_2 \oplus \dots$ be the associated graded sheaf. By Theorem 5.14, the stable bundles Q_i are hyperholomorphic. Since $c_{\text{rk } B}(B) \neq 0$, we have $c_{\text{rk } Q_i}(Q_i) \neq 0$. Therefore, the bundles Q_i are non-trivial. By Corollary 10.23, $\text{rk } Q_i > 2$. Since all the Chern classes of the bundles Q_i are $SU(2)$ -invariant, the odd Chern classes of Q_i vanish (Lemma 2.6). Therefore, $\text{rk } Q_i \geq 4$ for all i . Since $\text{rk } B \leq 6$, we have $i = 1$ and the bundle B' is stable. ■

Let M be a K3 surface, $X = M^{[i]}$, $i = 2, 3$ be its second or third Hilbert scheme of points, $I \in \text{Comp}(X)$ arbitrary complex structure on X , and a a generic polarization on (X, I) . Consider the hyperkähler structure $\mathcal{H}$ which corresponds to J and a by Calabi–Yau theorem (Theorem 2.4). Denote by TX the tangent bundle of X , considered as a hyperholomorphic bundle. Let $\text{Def}(TX)$ denote the hyperkähler desingularization of the moduli of stable deformations of TX . By Theorem 10.14, the real analytic subvariety underlying $\text{Def}(TX)$ is independent from the choice of I . Therefore, its dimension is also independent from the choice of I . The dimension of the deformation space $\text{Def}(TX)$ can be estimated by a direct computation, for X a Hilbert scheme. We obtain that $\dim \text{Def}(TX) > 40$.

Claim 10.25: In these assumptions, the space $\text{Def}(TX)$ is a compact hyperkähler manifold.

Proof: By Corollary 10.24, all semistable bundle deformations of TX are stable. Then Claim 10.25 is implied by Theorem 10.20. ■

Clearly, deforming the complex structure on X , we obtain a deformation of complex structures on $\text{Def}(TX)$. This gives a map

$$\text{Comp}(X) \longrightarrow \text{Comp}(\text{Def}(TX)). \quad (10.1)$$

It is easy to check that the map (10.1) is complex analytic, and maps twistor curves to twistor curves.

Claim 10.26: Let X, Y be hyperkähler manifolds, and

$$\varphi : \text{Comp}(X) \longrightarrow \text{Comp}(Y)$$

be a holomorphic map of corresponding moduli spaces which maps twistor curves to twistor curves. Then φ is locally an embedding.

Proof: An elementary argument using the period maps, in the spirit of Subsection 5.2. ■

The following result, along with Theorem 10.22, is the major stumbling block on the way to proving Conjecture 10.21. The other results of this Subsection are elementary or routinely proven, but the complete proof of Theorem 10.22 and Theorem 10.27 seems to be difficult.

Theorem 10.27: Let X be a simple hyperkähler manifold without proper trianalytic subvarieties, B a hyperholomorphic bundle over X , and I an induced complex structure. Denote the corresponding holomorphic bundle over (X, I) by B_I . Assume that the space $\mathcal{M}$ of stable bundle deformations of B is compact. Let $\text{Def}(B)$ be the hyperkähler desingularization of $\mathcal{M}$. Then $\text{Def}(M)$ is a simple hyperkähler manifold.

Proof: Given a decomposition $\text{Def}(M) = M_1 \times M_2$, we obtain a parallel 2-form on Ω_1 on $\text{Def}(B)$, which is a pullback of the holomorphic symplectic form on M_1 . Consider the space $\mathcal{A}$ of connections on B , which is an infinitely-dimensional complex analytic Banach manifold. Then Ω_1 corresponds to a holomorphic 2-form $\tilde{\Omega}_1$ on $\mathcal{A}$. Since Ω_1 is parallel with respect to the natural connection on $\text{Def}(B)$, the form $\tilde{\Omega}_1$ is also a parallel 2-form on the tangent space to $\mathcal{A}$, which is identified with $\Omega^1(X, \text{End}(B))$. It is possible to prove that this 2-form is obtained as

$$A, B \longrightarrow \int_Y \Theta \left(A|_Y, B|_Y \right) \text{Vol}(Y)$$

where

$$\Theta : \Omega^1(Y, \text{End}(B)) \times \Omega^1(Y, \text{End}(B)) \longrightarrow \mathcal{O}_Y$$

is a certain holomorphic pairing on the bundle $\Omega^1(Y, \text{End}(B))$, and Y is a trianalytic subvariety of X . Since X has no trianalytic subvarieties, $\tilde{\Omega}_1$ is obtained from a $\mathcal{O}_X$ -linear pairing

$$\Omega^1(X, \text{End}(B)) \times \Omega^1(X, \text{End}(B)) \longrightarrow \mathcal{O}_X.$$

Using stability of B , it is possible to show that such a pairing is unique, and thus, Ω_1 coincides with the holomorphic symplectic form on $\text{Def}(B)$. Therefore, $\text{Def}(B) = M_1$, and this manifold is simple. ■

Return to the deformations of tangent bundles on $X = M^{[i]}$, $i = 2, 3$. Recall that the second Betti number of a Hilbert scheme of points on a K3 surface is equal to 23, and that of the generalized Kummer variety is 7

([Bea]). Consider the map (10.1). By Theorem 10.27, the manifold $\text{Def}(TX)$ is simple. By Bogomolov's theorem (Theorem 5.9), we have

$$\dim \text{Comp}(\text{Def}(TX)) = \dim H^2(\text{Def}(TX)) - 2.$$

Therefore, either $\dim H^2(\text{Def}(TX)) > \dim H^2(X) = 23$, or the map (10.1) is étale. In the first case, the second Betti number of $\text{Def}(TX)$ is bigger than that of known simple hyperkähler manifolds, and thus, $\text{Def}(TX)$ is a new example of a simple hyperkähler manifold; this proves Conjecture 10.21. Therefore, to prove Conjecture 10.21, we may assume that $\dim H^2(\text{Def}(TX)) = 23$, the map (10.1) is étale, and $\text{Def}(TX)$ is a deformation of a Hilbert scheme of points on a K3 surface.

Consider the universal bundle $\tilde{B}$ over $X \times \text{Def}(TX)$. Restricting $\tilde{B}$ to $\{x\} \times \text{Def}(TX)$, we obtain a bundle B on $\text{Def}(TX)$. Let $\text{Def}(B)$ be the hyperkähler desingularization of the moduli space of stable deformations of B . Clearly, the manifold $\text{Def}(B)$ is independent from the choice of $x \in X$. Taking the generic hyperkähler structure on X , we may assume that the hyperkähler structure $\mathcal{H}$ on $\text{Def}(TX)$ is also generic. Thus, $(\text{Def}(TX), \mathcal{H})$ admits C -restricted complex structures and has no trianalytic subvarieties. In this situation, Corollary 10.24 implies that the hyperkähler manifold $\text{Def}(B)$ is compact. Applying Claim 10.26 again, we obtain a sequence of maps

$$\text{Comp}(X) \longrightarrow \text{Comp}(\text{Def}(TX)) \longrightarrow \text{Comp}(\text{Def}(B))$$

which are locally closed embeddings. By the same argument as above, we may assume that the composition $\text{Comp}(X) \longrightarrow \text{Comp}(\text{Def}(B))$ is étale, and the manifold $\text{Def}(B)$ is a deformation of a Hilbert scheme of points on K3. Using Mukai's version of Fourier transform ([O], [BBR]), we obtain an embedding of the corresponding derived categories of coherent sheaves,

$$D(X) \longrightarrow D(\text{Def}(TX)) \longrightarrow D(\text{Def}(B)).$$

Using this approach, it is easy to prove that

$$\dim X \leq \dim \text{Def}(TX) \leq \dim \text{Def}(B).$$

Let $x \in X$ be an arbitrary point. Consider the complex $C_x \in D(\text{Def}(B))$ of coherent sheaves on $\text{Def}(B)$, obtained as a composition of the Fourier-Mukai transform maps. It is easy to check that the lowest non-trivial cohomology sheaf of C_x is a skyscraper sheaf in a point $F(x) \in \text{Def}(B)$. This gives an embedding

$$F : X \longrightarrow \text{Def}(B).$$

The map F is complex analytic for all induced complex structure. We obtained the following result.

Lemma 10.28: In the above assumptions, the embedding

$$F : X \longrightarrow \mathrm{Def}(B)$$

is compatible with the hyperkähler structure.

■

By Lemma 10.28, the manifold $\mathrm{Def}(B)$ has a trianalytic subvariety $F(X)$, of dimension $0 < \dim F(X) < 40 < \dim \mathrm{Def}(B)$. On the other hand, for a hyperkähler structure on X generic, the corresponding hyperkähler structure on $\mathrm{Def}(B)$ is also generic, so this manifold has no trianalytic subvarieties. We obtained a contradiction. Therefore, either $\mathrm{Def}(TX)$ or $\mathrm{Def}(B)$ is a new example of a simple hyperkähler manifold. This proves Conjecture 10.21.

Acknowledgments: I am grateful to V. Batyrev, A. Beilinson, P. Deligne, D. Gaitsgory, D. Kaledin, D. Kazhdan, M. Konsevich and T. Pantev for valuable discussions. My gratitude to D. Kaledin, who explained to me the results of [Sw]. This paper uses many ideas of our joint work on direct and inverse twistor transform ([KV]).

References

- [BS] Bando, S., Siu, Y.-T, *Stable sheaves and Einstein-Hermitian metrics*, In: Geometry and Analysis on Complex Manifolds, Festschrift for Professor S. Kobayashi's 60th Birthday, ed. T. Mabuchi, J. Noguchi, T. Ochiai, World Scientific, 1994, pp. 39-50.
- [BBR] Bartocci, C., Bruzzo, U., Hernandez Ruiperez, D., *A Fourier-Mukai transform for stable bundles on K3 surfaces*, J. Reine Angew. Math. **486**, pp. 1-16 (1997).
- [Bea] Beauville, A. *Varieties Kähleriennes dont la première classe de Chern est nulle*. J. Diff. Geom. **18**, pp. 755-782 (1983).
- [Bes] Besse, A., *Einstein Manifolds*, Springer-Verlag, New York (1987)
- [Bo] Bogomolov, F.A., *Hamiltonian Kaehler manifolds*, (English. Russian original) Sov. Math., Dokl. **19**, 1462-1465 (1978); translation from Dokl. Akad. Nauk SSSR **243**, 1101-1104 (1978).

- [C] Calabi, E., *Metriques kähleriennes et fibrès holomorphes*, Ann. Ecol. Norm. Sup. **12** (1979), 269-294.
- [D] Deligne, P., *Categories tannakiennes*, The Grothendieck Festschrift, Collect. Artic. in Honor of the 60th Birthday of A. Grothendieck. Vol. II, Prog. Math. **87**, 111-195 (1990).
- [Do] Douady, A., *Le probleme des modules pour les sous-espaces analitiques compacts d'un espace analitique donné*, Ann. Inst. Fourier, **16** (1966), vol. 1, pp. 1-95.
- [O'G] O'Grady, K. G., *Desingularized moduli spaces of sheaves on a K3*, alg-geom/9708009, 41 pages, preliminary version.
- [Gi] Gieseker, D. *On the moduli of vector bundles on an algebraic surface*, Ann. Math., II. Ser. **106**, 45-60 (1977).
- [GH] Griffiths, Ph., Harris, J., *Principles of Algebraic Geometry*, Wiley-Interscience, New York, 1978.
- [GMT] Guaraldo, F., Macri, P., Tancredi, A., *Topics on real analytic spaces*, Advanced lectures in mathematics, Braunschweig: F. Vieweg, 1986.
- [F] Fujiki, A., *On primitively symplectic compact Kähler V-manifolds of dimension four*, in Classification of algebraic and analytic manifolds, (Katata, 1982), 71-250, Progr. Math., 39, Birkhäuser Boston, 1983.
- [H-Gro] Hartshorne, R., Local cohomology; a seminar given by A. Grothendieck, Harvard University, Fall, 1961. // Lecture notes in mathematics, vol. 41, Springer-Verlag, 1967.
- [HKLR] N.J. Hitchin, A. Karlhede, U. Lindström, M. Roček, *Hyperkähler metrics and supersymmetry*, Comm. Math. Phys (1987).
- [H] Huybrechts, D., Birational symplectic manifolds and their deformations, alg-geom/9601015.
- [K] Kaledin, D., *Integrability of the twistor space for a hypercomplex manifold*, alg-geom/9612016, 9 pages, Latex2e.
- [KV] Kaledin, D., Verbitsky, M., *Non-Hermitian Yang-Mills connections*, // alg-geom 9606019 (1996), 48 pages, LaTeX 2e.
- [Ko] Kobayashi S., *Differential geometry of complex vector bundles*, // Princeton University Press, 1987.
- [Ma1] Maruyama, Masaki *On boundedness of families of torsion free sheaves*, J. Math. Kyoto Univ. **21**, 673-701 (1981)
- [Ma2] Maruyama, Masaki, *Construction of moduli spaces of stable sheaves via Simpson's idea*, Lect. Notes Pure Appl. Math. **179**, 147-187 (1996).

- [LY] Li, Jun, and Yau, S.-T., *Hermitian Yang-Mills connections on non-Kähler manifolds*, in “Mathematical aspects of string theory” (S.T. Yau ed.), World Scientific Publ., London, 1987, pp. 560-573.
- [N1] Nitta, Takashi, *Vector bundles over quaternionic Kähler manifolds*, Tohoku Math. J., II. Ser. 40, No.3, 425-440 (1988).
- [N2] Nitta, Takashi, *Moduli spaces of Yang-Mills connections over quaternionic Kähler manifolds*, Osaka J. Math. **26**, No.2, 287-298 (1989).
- [OSS] Christian Okonek, Michael Schneider, Heinz Spindler, *Vector bundles on complex projective spaces*. Progress in mathematics, vol. 3, Birkhauser, 1980.
- [O] Orlov, D. **Equivalences of derived categories and K3 surfaces**, alg-geom/9606006.
- [Sal] Salamon, S., *Quaternionic Kähler manifolds*, Inv. Math. **67** (1982), 143-171.
- [Sw] Swann, A., *HyperKähler and quaternionic Kähler geometry*, Math. Ann. **289**, No. 3, 421-450 (1991).
- [UY] Uhlenbeck K., Yau S. T., *On the existence of Hermitian Yang-Mills connections in stable vector bundles*, // Comm. on Pure and Appl. Math., 39, p. S257-S293 (1986).
- [V0] M. Verbitsky, *On the action of the Lie algebra $\mathfrak{so}(5)$ on the cohomology of a hyperkähler manifold*, Func. Anal. and Appl. **24** (1990), 70-71.
- [V1] Verbitsky M., *Hyperholomorphic bundles over a hyperkähler manifold*, alg-geom electronic preprint 9307008 (1993), 43 pages, LaTeX, also published in: Journ. of Alg. Geom., **5** no. 4 (1996) pp. 633-669.
- [V2] Verbitsky M., *Hyperkähler and holomorphic symplectic geometry I*, Journ. of Alg. Geom., **5** no. 3 (1996) pp. 401-415. A different version appeared as alg-geom/9307009
- [V2(II)] Verbitsky M., *Hyperkähler embeddings and holomorphic symplectic geometry II*, alg-geom electronic preprint 9403006 (1994), 14 pages, LaTeX, also published in: GAFA **5** no. 1 (1995), 92-104.
- [V3] Verbitsky, M., *Cohomology of compact hyperkähler manifolds*. alg-geom electronic preprint 9501001, 89 pages, LaTeX.
- [V3-bis] Verbitsky, M., *Cohomology of compact hyperkähler manifolds and its applications*, alg-geom electronic preprint 9511009, 12 pages, LaTeX, also published in: GAFA vol. 6 (4) pp. 601-612 (1996).
- [V4] Verbitsky M., *Deformations of trianalytic subvarieties of hyperkähler manifolds*, alg-geom electronic preprint 9610010 (1996), 51 pages, LaTeX2e.
- [V5] Verbitsky M., *Trianalytic subvarieties of the Hilbert scheme of points on a K3 surface*, alg-geom/9705004.

- [V-a] Verbitsky M., *Algebraic structures on hyperkähler manifolds*, alg-geom electronic preprint 9609011, also published in: Math. Res. Lett., **3**, 763-767 (1996).
- [V-d] Verbitsky M., *Desingularization of singular hyperkähler varieties I*, electronic preprint 9611015 (1996), 13 pages, LaTeX 2e, also published in: Math. Res. Lett. **4** (1997), no. 2-3, pp. 259–271.
- [V-d2] Verbitsky M., *Desingularization of singular hyperkähler varieties II*, alg-geom/9612013 (1996), 15 pages, LaTeX 2e.
- [V-d3] Verbitsky M., *Hypercomplex Varieties*, alg-geom/9703016 (1997), 40 pages, LaTeX (to appear in Comm. in An. and Geom.)
- [Y] Yau, S. T., *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation I.* // Comm. on Pure and Appl. Math. **31**, 339-411 (1978).